

BIBLIOGRAFIA

-
- Adler DA, Quaderi NA, Brown SD, Chapman VM, Moore J, Tate P, Disteché CM (1995) The X-linked methylated DNA binding protein, MeCP2, is subject to X inactivation in the mouse. *Mamm Genome*. 6:491-492
 - Akbarian S, Chen RZ, Gribnau J, Rasmussen TP, Fong H, Jaenisch R, Jones EG (2001) Expression pattern of the Rett syndrome gene MeCP2 in primate prefrontal cortex. *Neurobiol Dis* 8:784-791
 - Akesson HO, Hagberg B, Wahlstrom J, Engerstrom IW (1992) Rett syndrome: a search for gene sources. *Am J Med Genet* 42:104-8
 - Akesson HO, Wahlstrom J, Engerstrom IW, Hagberg B (1995) Rett syndrome: potential gene sources: phenotypical variability. *Clin Genet* 48:169-172
 - Akesson HO, Hagberg B, Wahlstrom J (1996) Rett syndrome, classical and atypical: genealogical support for common origin. *J Med Genet* 33:764-766
 - Amir RE, Van den Veyver IB, Wan M, Tran CQ, Franke U, Zoghbi HY (1999) Rett syndrome is caused by mutations in X-linked MECP2, encoding methyl-CpG-binding protein 2. *Nature Genet* 23:185-188
 - Amir RE, Van den Veyver IB, Schultz R, Malicki DM, Tran CQ, Dahle EJ, Philippi A, Timar L, Percy AK, Motil KJ, Lichtarge O, Smith EO, Glaze DG, Zoghbi HY (2000) Influence of mutation type and X chromosome inactivation on Rett syndrome phenotypes. *Ann Neurol* 47:670-679
 - Anvret M, Wahlstrom J, Skogserberg P, Hagberg B (1990) Segregation analysis of the X-chromosome in a family with Rett syndrome in two generations. *Am J Med Genet* 37:31-35
 - Anvret M & Wahlstrom J (1994) Rett Syndrome: random X chromosome inactivation. *Clin Genet* 45:274-275
 - Archidiacono N, Lerone M, Rocchi M, Anvret M, Ozcelik T, Francke U, Romero G (1991) Rett syndrome: exclusion mapping following the hypothesis of germinal mosaicism for new X-linked mutations. *Hum Genet* 86:604-606
 - Armstrong DD (1992) The neuropathology of the Rett syndrome. *Brain Dev* 14 Suppl:S89-S98
 - Armstrong D, Dunn JK, Antalffy B, Trivedi R (1995) Selective dendritic alternations in the cortex of Rett syndrome. *J Neuropathol Exp Neurol* 54:195-201
 - Armstrong DD, Dunn K, Antalffy B (1998) Decreased dendritic branching in frontal, motor and limbic cortex in Rett syndrome compared with trisomy 21. *J Neuropathol Exp Neurol* 57:1013-1017
 - Armstrong DD, Dunn JK, Schultz RJ, Herbert DA, Glaze DG, Motil KJ (1999) Organ growth in Rett syndrome: a postmortem examination analysis. *Pediatr Neurol* 20:125-129

- Armstrong DD (2001) Rett syndrome neuropathology review 2000. *Brain Dev* 23 Suppl 1:S72-S76
- Armstrong J, Pineda M, Monrós E (2000) Mutation analysis of 16SrRNA in patients with Rett syndrome. *Pediatr Neurol* 23:85-87
- Armstrong J, Poo P, Pineda M, Aibar E, Geán E, Català V, Monrós E (2001) Classic Rett syndrome in a boy as a result of somatic mosaicism for a *MECP2* mutation. *Ann Neurol* 5:692
- Armstrong J, Aibar E, Pineda M, Pérez MM, Géan E, Carrera M, Casas C, Martínez F, Monrós E (2002) Prenatal diagnosis in Rett syndrome. *Fetal Diagnostic and Therapy* 17:200-204.
- Armstrong J, Aibar E, González E, Garrido C, Jabalera M, Campistol J, Poo P, Vernet A, Pérez MM, Géan E, Pineda M, Monrós E. Genetic and clinical correlation analysis of Rett syndrome in Spain. (*submitted*).
- Ars E, Serra A, García J, Kruyer H, Gaona A, Lázaro C, Estivill X (2001) Mutations affecting mRNA splicing are the most common molecular defects in patients with neurofibromatosis type 1. *Hum Mol Genet* 9:237-247
- Auranen M, Vanhala R, Vosman M, Levander M, Varilo T, Hietala M, Riikonen R, Peltonen L, Jarvela I (2001) *MECP2* gene analysis in classical Rett syndrome and in patients with Rett-like features. *Neurology* 56:611-617B
- Ballestar E & Wolffe AP (2001) Methyl-CpG-binding proteins: Targeting specific gene repression. *Eur J Biochem FEBS* 268:1-6
- Ballestar E, Yusufzai TM, Wolffe AP (2000) Effects of Rett syndrome mutations of the methyl-CpG binding domain of the transcriptional repressor MeCP2 on selectivity for association with methylated DNA. *Biochemistry* 39:7100-7106
- Balmer D, Arredondo J, Samaco RC, LaSalle JM (2002) *MECP2* mutations in Rett syndrome adversely affect lymphocyte growth, but do not affect imprinted gene expression in blood or brain. *Hum Genet* 110(6):545-52
- Balmer D, Goldstine J, Rao YM, LaSalle JM (2003) Elevated methyl-CpG-binding protein 2 expression is acquired during postnatal human brain development and is correlated with alternative polyadenylation. *J Mol Med* 81:61-68
- Bauman ML, Kemper TL, Arin DM (1995) Pervasive neuroanatomic abnormalities of the brain in three cases of Rett's syndrome. *Neurology* 45:1581-1586
- Belinchenko PV, Oldfors A, Hagberg B, Dahstrom A (1994) Rett syndrome: 3-D confocal microscopy of cortical pyramidal dendrites and afferents. *Neuroreport* 5:1509-1513
- Belknap JK (1992) Empirical estimates of Bonferroni corrections for use in chromosome mapping studies with the BXD recombinant inbred strains. *Behav Genet* 22:677-684

-
- Benedetti L, Munnich A, Melki J, Tardieu M, Turleau C (1992) Parental origin of the X chromosome in Rett syndrome. *Am J Med Genet* 44:121-122
 - Beyer KS, Blasi F, Bacchelli E, Klauck SM, Maestrini E, Poustka A (2002) Mutation analysis of the coding sequence of the MECP2 gene in infantile autism. *Hum Genet* 111:305-309
 - Bienvenu T, Carrié A, de Roux N, Vinet MC, Jonveaux P, Couvert P, Villard L, Arzimanoglou A, Beldjord C, Fontes M, Tardieu M, Chelly J (2000) MECP2 mutations account for most cases of typical forms of Rett syndrome. *Hum Mol Genet* 9:1377-1384
 - Boeke J, Ammerpohl O, Kegel S, Moehren U, Renkawitz R (2000) The minimal repression domain of MBD2b overlaps with the methyl-CpG-binding domain and binds directly to Sin3A. *J Biol Chem* 275:34963-34967
 - Bourdon V, Philippe C, Labrune O, Amsallem D, Arnould C, Jonveaux P (2001) A detailed analysis of the MECP2 gene: prevalence of recurrent mutations and gross DNA rearrangements in Rett syndrome patients. *Hum Genet* 108:43-50
 - Buyse IM, Fang P, Hoon KT, Amir RE, Zoghbi HY, Roa BB (2000) Diagnostic testing for Rett syndrome by DHPLC and direct sequencing analysis of the MECP2 gene: identification of several novel mutations and polymorphisms. *Am J Hum Genet* 67:1428-1436
 - Camus P, Abbadì N, Perrier MC, Chery M, Gilgenkrantz S (1996) X chromosome inactivation in 30 girls with Rett syndrome. *Hum Genet* 97:247-250
 - Cardioli E, Dotti MT, Hayek G, Zappella M, Federico AJ (1999) Studies on mitochondrial pathogenesis of Rett syndrome: ultrastructural data from skin and muscle biopsies and mutational analysis at mtDNA nucleotides 10463 and 2835. *Submicrosc Cytol Pathol* 31:301-304
 - Chandler SP, Guschin D, Landsberger N, Wolffe AP (1999) The methyl-CpG binding transcriptional repressor MeCP2 stably associates with nucleosomal DNA. *Biochemistry* 38:7008-7018
 - Cheadle JP, Gill H, Fleming N, Maylanrd J, Kerr A, Leonard H, Krawczak M, Cooper DN, Lynch S, Thomas NST, Hugues HE, Hulten M, Ravine D, Clarke A (2000) Long-read sequence analysis of the MECP2 gene in Rett syndrome patients: correlation of disease severity with mutation type and location. *Hum Mol Genet* 9:1119-1129
 - Chen RZ, Akbarian S, Tudor M, Jaenisch R (2001) Deficiency of methyl-CpG binding protein 2 in CNS neurons results in a Rett-like phenotype in mice. *Nat Genet* 27:327-331
 - Christy RJ, Kaestber KH, Geiman DE, Lane MD (1991) CCAAT/enhancer binding protein gene promoter: binding of nuclear factor during differentiation of 3T3-L1 preadipocytes. *Proc Natl Acad Sci* 88:2593-7
 - Clark A (1996) Rett syndrome. *J Med Genet* 33:693-699
-

- Clayton-Smith J, Watson P, Ramsden S, Black GC (2000) Somatic mutation in *MECP2* as a non-fatal neurodevelopmental disorder in males. *Lancet* 356:830-832
- Coker SB & Melnyk AR (1991) Rett syndrome and mitochondrial enzyme deficiencies. *J Child Neurol* 6:164-166
- Conforti FL, Massei R, Magariello A, Patitucci A, Gabriele AL, Muglia M, Quattrone A, Fiumara A, Barone R, Pavone L, Nisticò R, Mangone L (2003) Mutation analysis of the *MECP2* gene in patients with Rett syndrome. *Am J Med Genet* 117A:184-187
- Couvert P, Bienvenu T, Aquaviva C, Poirier K, Moraine C, Gendrot C, Verloes A, Andres C, Le Fevre AC, Souville I, Steffann J, des Portes V, ROperes HH, Yntema HG, Fryns JP, Briault S, Chelly J, Cherif B (2001) *MECP2* is highly mutated in X-linked mental retardation. *Hum Mol Genet* 10: 941-946
- Coy JF, Sedlacek Z, Bachner D, Celius H, Poustka A (1999). A complex pattern of evolutionary conservation and alternative polyadenylation within the long 3'-untranslated region of the methyl-CpG-Binding protein 2 gene (*MeCP2*) suggests a regulatory role in gene expression. *Hum Mol Genet* 8:1253-1262
- Cross SH, Meehan RR, Nan X, Bird A (1997) A component of the transcriptional repressor *MeCP1* shares a motif with DNA methyltransferase and HRX proteins. *Nat Genet* 16:256-259
- Cummings CJ, Dahle EJ, Zoghbi HY (1998) Analysis of the genomic structure of the human glycine receptor alpha 2 subunit gene and exclusion of this gene as a candidate for Rett syndrome. *Am J Med Genet* 78:176-178
- Curtis AR, Headland S, Lindsay S, Thomas NS, Boye E, Kamakari S, Roustan P, Anvret M, Wahlstrom J, McCarthy G (1993) X chromosome linkage studies in familial Rett syndrome. *Hum Genet* 90:551-555
- D'Esposito M, Quaderi NA, Ciccodicola A, Bruni P, Esposito T, D'Urso M, Brown SDM (1996) Isolation, physical mapping and Northern analysis of the X-linked human gene encoding methyl-CpG-binding protein, *MeCP2*. *Mamm Genome* 7:533-535
- Daniel JM & Reynolds AB (1999) The catenin p120(cnt) interacts with Kaiso, a novel BTB/POZ domain zinc finger transcription factor. *Mol Cell Biol* 19:3614-3623
- De Bona C, Zappella M, Hayeck G, Meloni I, Vitelli F, Bruttini M, Cusano R, Loffredo P, Longo I, Ranieri A (2000) Preserved speech variant is allelic of classic Rett syndrome. *Eur J Hum Genet* 8:325-330
- Deguchi K, Anatlffy BA, Twohill LJ, Chakraborty S, Glaze DG, Armstrong DD (2000) Substance P immunoreactivity in Rett syndrome. *Pediatr Neurol* 22:259-266
- DeZazzo J, Sandstrom D, de Bell S, Velinzon K, Smith P, Grady L, DelVecchio M, Ramaswami M, Tully T. (2000) *nayot*, a mutation of the *Drosophila* myb-related *Adf1* transcription factor, disrupts synapse formation and olfactory memory. *Neuron* 27:145-58

-
- Diagnostic criteria for Rett syndrome (1988) The Rett Syndrome Diagnostic Criteria Work Group. *Ann Neurol* 23:425-428
 - Dotti MT, Manneschi L, Malandrini A, De Stefano N, Caznerale F, Federico A (1993) Mitochondrial dysfunction in Rett syndrome: An ultrastructural and biochemical study. *Brain Dev* 15:103-106
 - Dotti MT, Orrico A, De Stefano N, Battisti C, Sicurelli F, Severi S, Lam CW, Galli L, Sorrentino V, Federico A (2002) A Rett syndrome MECP2 mutation that causes mental retardation in men. *Neurology* 58:226-30
 - Dragich J, Houwink-Manville I, Schanen C (2000) Rett syndrome: a surprising result of mutation in MECP2. *Hum Mol Genet* 9:2365-2375
 - Ellison KA, Fill CP, Terwilliger J, DeGennari LJ, Martin-Gallardo A, Anvret M, Percy AK, Ott J, Zoghbi U (1992) Examination of X chromosome markers in Rett syndrome: exclusion mapping with a novel variation on multilocus linkage analysis. *Am J Hum Genet* 50:278-287
 - Everitt BS (1997) Annotation: correspondence analysis. *J Child Psychol Psychiatry* 38:737-745
 - Fañanás L, Moral P, Panadero MA, Bertranpetit J (1992) Complement genetic markers in schizophrenia: C3, BF and C6 polymorphisms. *Hum Hered* 42:162-167
 - Foulds GJ & Etzkorn FA (1998) A capillary eletrophoresis mobility shift assay for protein-DNA binding affinities free in solution. *Nucleic Acids Res* 26:4304-4305.
 - Fraga MF, Rodriguez R, Canal MJ (2000) Rapid quantification of DNA methylation by high performance capillary electrophoresis. *Electrophoresis* 21:2990-2994.
 - Fraga MF, Uriol E, Borja Diego J, Berdasco M, Esteller M, Canal MJ, Rodriguez R (2002) High-performance capillary electrophoresis method for the quantification of 5-methyl 2'deoxyctidine in genomic DNA: application to plant, animal and human cancer tissues. *Electrophoresis* 23:1677-81.
 - Free A, Wakefield RI, Smith BO, Dryden DT, Barlow PN, Bird AP (2001) DNA recognition by the methyl-CpG binding domain of MeCP2. *J Biol Chem* 276:3353-3360
 - Gale RE & Linch DC (1994) Interpretation of X-chromosome inactivation patterns. *Blood* 84:2376-2378
 - Garcia-Giralt N, Nogués X, Enjuanes A, Puig J, Mellibovsky L, Bay-Jensen A, Carreras R, Balcells S, Díez-Pérez A, Grinberg D (2002) Two new single-nucleotide polymorphisms in the COL1A1 upstream regulatory region and their relationship to Bone Mineral Density. *J Bone Miner Res* 17:384-393
 - Gartler SM & Riggs AD (1983) Mammalian X-chromosome inactivation. *Annu Rev Genet* 17:155-190
-

- Geerdink N, Rotteveed JJ, Lammens M, Sistermans EA, Heikens GT, Gabreels FJ, Mullaart RA, Hamel BC (2002) *MECP2* mutation in a boy with severe neonatal encephalopathy: clinical, neuropathological and molecular findings. *Neuropediatrics* 33:33-36
- Gill H, Cheadle JP, Maylanrd J, Fleming N, Whatley S, Cranston T, Thompson EM, Leonard H, Davis M, Christoloulou, Skjeldal O, Hanefeld F, Kerr A, Tandy A, Ravine D, Clarke A (2003) Mutation analysis in the *MECP2* gene and genetic counseling for Rett syndrome. *J Med Genet* 40:380-384
- Goto T & Monk M (1998) Regulation of X-chromosome inactivation in development in mice and humans. *Microbiol Mol Biol Rev* 62:362-78
- Gregory RI, Randall TE, Johnson CA, Khosla S, Hatada I, O'Neill LP, Turner BM, Feil R (2001) DNA methylation is linked to deacetylation of histone H3, but not H4, on the imprinted genes *Snrpn* and *U2f1-rs1*. *Mol Cell Biol* 21:5426-5436
- Guy J, Hendrich B, Holmes M, Martin JE, Bird A (2001) A mouse *Mecp2*-null mutation causes neurological symptoms that mimic Rett syndrome. *Nat Genet* 27:322-326
- Haas RH, Dixon SD, Sartoris DJ, Hennessy MJ (1997) Osteopenia in Rett syndrome. *J Pediatr* 131:771-774
- Hagberg B, Aicardi J, Dias K, Ramos O (1983) A progressive syndrome of autism, dementia, ataxia and loss of purposeful hand use in girls: Rett's syndrome: report of 35 cases. *Ann Neurol* 14:471-479
- Hagberg B (1985) Rett's syndrome: prevalence and impact on progressive severe mental retardation in girls. *Acta Paediatr Scand* 74:405-408
- Hagberg B & Witt-Engerstom I (1986) Rett syndrome: a suggested staging system for describing impairment profile with increasing towards adolescence. *Am J Med Genet Suppl* 1:47-59
- Hagberg BA & Skjeldal OH (1994) Rett variants: a suggested model for inclusion criteria. *Pediatr Neurol* 11:5-11
- Hagberg B, Hanefeld F, Percy A, Skjeldal O (2002) An update on clinically applicable diagnostic criteria in Rett syndrome. Comments to Rett Syndrome clinical criteria consensus panel satellite to European Paediatric Neurology Society Meeting, Baden Baden, Germany, 11th September 2001. *Eur J Paediatr Neurol* 6:293-297
- Harley HG, Brook JD, Rundle SA, Crow S, Brook JD, Harper PS, Shaw DJ (1992) Expansion of unstable DNA region and phenotype variation in myotonic dystrophy. *Nature* 335:545-546
- Hassig CA, Fleischer TC, Billini AN, Schreiber SL, Ayer DE (1997) Histone deacetylase activity is required for full transcriptional repression by mSin3A. *Cell* 89:341-347.

-
- Heidary G, Hampton LL, Schanen NC, Rivkin MJ, Darras BT, Battey J, Francke U (1998) Exclusion of the gastrin-releasin peptide receptor (GRPR) locus as a candidate gene for Rett syndrome. *Am J Med Genet* 78:173-175
 - Heinzl T, Lavinsky RM, Mullen TM, Soderstrom M, Laherty CD, Torchia J, Yang WM, Brard G, Ngo SD, Davie JR, Seto E, Eisenman RN, Rose DW, Glass CK, Rosenfeld MG (1997) A complex containing N-CoR, mSin3 and histone deacetylase mediates transcriptional repression. *Nature* 387:43-48
 - Hendrich B & Bird A (1998) Identification and characterization of a family of mammalian methyl-CpG binding proteins. *Mol Cell Biol* 18:6538-47
 - Hendrich B, Abbott C, McQueen H, Chambers D, Cross S, Bird A. (1999) Genomic structure and chromosomal mapping of the murine and human Mbd1, Mbd2, Mbd3, and Mbd4 genes. *Mamm Genome* 10:906-12.
 - Herbst D & Miller J (1980) Nonsepecific X-linked mental retardation II: the frequency in British Columbia. *Am J Med Genet* 7:461-469
 - Hoffbuhr K, Devaney J, LaFleur B, Sirianni N, Scacheri C, Giron J, Schuette J, Innis J, Marino M, Philippart M, Narayanan V, Umansky R, Kronn, Hoffman EP, Naidu S (2001) Genotype and Phenotype evaluation of MeCP2 mutations in Rett syndrome. *Neurology* 56:1486-1495
 - Hofferbert S, Schanen NC, Budden SS, Francke U (1997) Is Rett syndrome caused by triplet repeat expansion? *Neuropediatrics* 28:179-183
 - Holm VA (1986) Physical growth and development in patients with Rett syndrome. *Am J Med Genet Suppl* 1:119-126
 - Huppke P, Held M, Hanefeld F, Engel W, Laccone F (2002) Influence of mutation type and location on phenotype in 123 patients with Rett syndrome. *Neuropediatrics*. 33:63-68
 - Huppke P, Laccone F, Kramer N, Engel W, Hanefeld F (2000) Rett syndrome: analysis of MECP2 and clinical characterization of 31 patients. *Hum Mol Genet* 22:1369-1375
 - Imessaoudene B, Bonnefont JP, Royer G, Cormier-Daire V, Lyonner s, Lyon G, Munnich A, Amiel J (2001) MECP2 mutation in non-fatal, non progressive encephalopathy in a male. *J Med Genet* 38:171-174
 - Inui K, Akagi M, Ono J, Tsukamoto H, Shimono K, Mano T, Imai K, Yamada M, Muramatsu T, Sakai N, Okada S (2001) Mutation analysis of MECP2 in Japanese patients with atypical Rett syndrome. *Brain Dev* 23:212-215
 - Jacobson M (1991) *Developmental Neurology*, 3rd edn. Plenum Press, New York
 - Jan MM, Dooley JM, Gordon KE (1999) Male Rett syndrome variant: application of diagnostic criteria *Pediatr Neurol* 20:238-40

- Jellinger K & Seitelberg F (1986) Neuropathology of Rett syndrome. *Am J Med Genet Suppl* 1:259-288
- Jellinger KA (2003) Rett Syndrome - an update. *J Neural Transm.* 110:681-701
- Johnson WF (1980) Metabolic interference and the +/- heterozygote: a hypothetical form of simple inheritance which is neither dominant nor recessive. *Am J Hum Genet* 32:374-386
- Jones PL, Veenstra GJ, Wade PA, Vermaak D, Kass SU, Landsberger N, Strtoubouli J, Wolffe AP (1998) Methylated DNA and MeCP2 recruit histone deacetylase to repress transcription. *Nat Genet* 19:187-191
- Journel H, Melki J, Turleau C, Munnich A, de Grouchy J (1990) Rett phenotype with X/autosome translocation: possible mapping to the short arm of chromosome X. *Am J Med Genet* 35:142-147
- Jung BP, Jugloff FGM, Zhang G, Logan R, Brown S, Eubanks JH (2003) The expression of methyl-CpG binding factor MeCP2 correlates with cellular differentiation in the developing rat brain and in cultured cells. *J Neurobiol* 55:86-96
- Kaludov NF & Wolffe AP (2000) MeCP2 driven transcriptional repression in vitro: selectivity for methylated DNA, action at a distance and contacts with the basal transcription machinery. *Nucleic Acids Res* 28:1921-19-28
- Kaufmann WE, Naidu S, Budden S (1995) Abnormal expression of microtubule-associated protein 2 (MAP-2) in neocortex in Rett syndrome. *Neuropediatrics* 26:109-113
- Kaufmann WE, Worley PF, Pegg J, Mremer M, Isakson P (1996) COX-2, a synaptically induced enzyme, is expressed by excitatory neurons at postsynaptic sites in rat cerebral cortex. *Proc Natl Acad Sci USA* 93:2317-2321
- Kaufmann WE, Worley PF, Taylor CV, Bremer M, Isakson PC (1997) Cyclooxygenase-2 expression during rat neocortical development and in Rett syndrome. *Brain Dev* 19:25-34
- Kerr AM & Stephenson JB (1985) Rett's syndrome in the west of Scotland. *Br Med J (Clin Res Ed)* 291:579-582
- Kim SJ & Cook EH Jr (2000) Novel de novo nonsense mutation of MECP2 in a patient with Rett syndrome. *Hum Mut* 15:382-383
- Klauck SM, Lindsay S, Beyer KS, Splitt M, Burn J, Poustka A (2002) A mutation hot spot for nonspecific X-linked mental retardation in the MECP2 gene causes the PPM-X syndrome. *Am J Hum Genet* 70:1034-7
- Kokura K, Kaul SC; Wadhwa R, Nomura T, Khan MM, Shinagawa T, Yasukawa T, Colmenares C, Ishii S (2001) The Ski protein family is required for MeCP2-mediated transcriptional repression. *J Biol Chem* 276:34115-34121

-
- Kozinetz CA, Skender ML, MacNaughton N, Almes MJ, Schultz RJ, Percy AK, Glaze DG (1993) Epidemiology of Rett syndrome: a population-based registry. *Pediatrics* 91:445-450
 - Krawczak M, Ball EV, Cooper DN (1998) Neighboring-nucleotide effects on the rates of germ-line single-base-pair substitution in human genes. *Am J Hum Genet* 63:474-488
 - Kresppischi AC, Kok F, Otto PG (1998) X chromosome-inactivation patterns in patients with Rett syndrome. *Hum Genet* 102:319-321
 - Kudo S, Fukuda M (1993) Transcriptional activation of human leukosialin (CD43) gene by Sp1 through binding to a GGGTGG motif. *Eur J Biochem* 223:319-327
 - Kudo S (1998) Methyl-CpG-binding protein MeCP2 represses Sp1-activated transcription of the human leukosialin gene when the promoter is methylated. *Mol Cell Biol* 18:5492-5499
 - Kudo S, Nomura Y, Segawa M, Fujita N, Nakao M, Dragich J, Schanen C, Tamura M (2001) Functional analyses of MeCP2 mutations associated with Rett syndrome using transient expression systems. *Brain Dev* 23 Suppl 1:S165-S173
 - Kudo S, Nomura Y, Segawa M, Fujita N, Nakao M, Hammer S, Schanen C, Terai I, Tamura M (2002) Functional characterization of MeCP2 mutations found in male patients with X linked mental retardation. *J Med Genet* 39:132-136
 - Laan AEM & Vein AA (2002) A Rett patient with a typical Angelman EGG. *Epilepsia* 43:1590-1592
 - Laccone F, Zoll B, Huppke P, Hanefeld F, Pepinski W, Trappe R (2002) MECP2 gene nucleotide changes and their pathogenicity in males: proceed with caution. *J Med Genet* 39:586-588
 - Lam CW, Yeung WL, Ko CH, Poon PM, Tong SF, Chan KY, Lo IF, Chan LY, Hui J, Wong V, Pang CP, Lo YM, Fok TF (2000) Spectrum of mutations in the MECP2 gene in patients with infantile autism and Rett syndrome. *J Med Genet* 37:E41
 - LaSalle JM, Goldstine J, Balmer D, Greco CK (2001) Quantitative localization of heterogeneous methyl-CpG-binding protein 2 (MeCP2) expression phenotypes in normal and Rett syndrome brain by laser scanning cytometry. *Hum Mol Genet* 10:1729-1740
 - LeCaptain DJ, Michel MA, Van Orden A (2001) Characterization of DNA-protein complexes by capillary electrophoresis-single molecule fluorescence correlation spectroscopy. *Analyst* 126:1279-1284.
 - Leonard H, Silberstein J, Falk R, Houwink-Manville I, Ellaway C, Raffaele LS, Engerstrom IW, Schanen C (2001) Occurrence of Rett syndrome in boys. *J Child Neurol* 16:333-338
 - Leonard H, Colvin L, Christodoulou J, Schiavello T, Williamson S, Davis M, Ravine D, Fyfe S, de Klerk N, Matsuishi T, Kondo I, Clarke A, Hackwell, Yamashita Y (2003) Patients with the R133C mutation: is their phenotype different from patients with Rett syndrome with other mutations? *J Med Genet* 40:e52
-

- Lewis JD, Meehan RR, Henzel WJ, Maurer-Fogy I, Jeppersen P, Klein F, Bird A (1992) Purification, sequence and cellular localization of a novel chromosomal protein that binds to methylated DNA. *Cell* 69:905-914
- Lobo-Menendez F, Sossey-Alaoui K, Bell JM, Copeland-Yates SA, Plank SM, Sanford SO, Skinner C, Simensen RJ, Schroer RJ, Michaelis RC (2003) Absence of MeCP2 mutations in patients from the South Carolina autism project. *Am J Med Genet* 15:97-101
- Lynch SA, Whatley SD, Ramesh V, Sinha, Ravine D (2003) Sporadic case of fatal encephalopathy with neonatal onset associated with a T158M missense mutation in MECP2. *Arch Dis Child Fetal Neonatal Ed* 88:F250-F252
- Lyon MF (1961) Gene action in the X-chromosome of the mouse (*Mus musculus* L.) *Nature* 190:372
- Lyon MF (2002) X-chromosome inactivation and human genetic disease. *Acta Paediatr Suppl* 91:107-112
- Maiwald R, Bönnte A, Jung H, Bitter P, Storm Z, Laccone F, Herkenrath P (2002) De novo MECP2 mutation in a 46,XX male patient with Rett syndrome. *Neurogenetics* 4:107-108
- Mak SC, Chi CS, Chen CH, Shian WJ (1993) Abnormal mitochondria in Rett syndrome: one case report. *Zhonghua Yi Xue Za Zhi (Taipei)* 52:116-119
- Maquat J (2001) The power of point mutations. *Nat Genet* 27:5-6
- Marin M, Karis A, Visser P, Grosveld F, Philipsen S (1997) Transcription factor Sp1 is essential for early embryonic development but dispensable for cell growth and differentiation. *Cell* 89:619-628
- Matsuishi T, Nagamitsu S, Yamashita Y, Murakami Y, Kimura A, Sakai T, Shoji H, Kato H, Percy AK (1997) Decreased cerebrospinal fluid levels of substance P in patients with Rett syndrome. *Ann Neurol* 42:978-981
- Meloni I, Bruttini M, Longo I, Mari F, Rizzolio F, D'Adamo P, Denvriendt K, Fryns JP, Toniolo D, Renieri A (2000) A mutation in the Rett syndrome gene, MECP2, causes X-linked mental retardation and progressive spasticity in males. *Am J Hum Genet* 67:982-985
- Miggeon BR, Dunn MA, Thomas G, Schmeckpeper BJ, Naidu S (1995) Studies of X inactivation and isodisomy in twins provide further evidence that the X chromosome is not involved in Rett syndrome. *Am J Hum Genet* 56:647-653
- Moncla A, Kpebe A, Missirian C, Mancini J, Villard L (2002) Polymorphisms in the C-terminal domain of MECP2 in mentally handicapped boys: implications for genetic counselling. *Eur J Hum Genet* 10:86-89
- Monrós E, Armstrong J, Aibar E, Poo P, Canós I, Pineda M (2001) Rett syndrome in Spain: mutation analysis, and clinical correlations. *Brain Dev* 23 Suppl 1:S251-S253

-
- Monrós E, Moltó MD, Martínez F, Cañizares J, Blanca J, Vilchez JJ, Prieto F, de Frutos R, Palau F (1997) Phenotype correlation and Intergenerational dynamics of the Friedreich ataxia GAA trinucleotide repeat. *Am J Hum Genet* 61:101-110
 - Moog U, Smeets EE, Van Roosendaal KE, Schoenmakers S, Herbergs J, Shoonbrood-Lenssen AM, Schnader-Stumpel CT (2003) Neurodevelopmental disorders in males related to the gene causing Rett syndrome in females (MECP2). *Eur J Paedr Neurol* 7:5-12
 - Moses LM*, Armstrong J*. Hoffbuhr K, Pineda M, Bibat G, González E, Galán F, Naidu S, Hoffman E, Monrós E. In-frame deletions in the MECP2 gene inherited as polymorphisms in two girls with Rett syndrome. *Both authors contributed equally to this work (*submitted*).
 - Motil KJ, Schultz RJ, Wong WW, Glaze DG (1998) Increased energy expenditure associated with repetitive involuntary movement does not contribute to growth failure in girls with Rett syndrome. *J Pediatr* 132:228-233
 - Nagy L, Kao HY, Chakravarti D, Lin RJ, Hassig CA, Ayer DE, Schreiber SL, Evans RM (1997) Nuclear receptor repression mediated by a complex containing SMRT, mSin2A, and histone deacetylase. *Cell* 89:373-380.
 - Naidu S, Murphy M, Moser HW, Rett A (1986) Rett syndrome-natural history in 70 cases. *Am J Med Genet Suppl* 1:61-72
 - Naidu SB (1997) Rett syndrome. *Indian J Pediatr* 64:651-9
 - Naidu S, Kaufmann WE, Abrams MT, Pearlson GD, Lanham DC, Fredericksen KA, Barker PB, Horska A, Golay X, Mori S, Wong DF, Yablonski M, Moser HW, Johnston MV (2001) Neuroimaging studies in Rett Syndrome. *Brain Dev* 23 Suppl 1S62-S71
 - Nan XS, Meehan RR, Bird A (1993) Dissection of the methyl-CpG-binding domain from the chromosomal protein MeCP2. *Nucleic Acids Res* 21:4886-4892
 - Nan X, Tate P, Li E, Bird A (1996) DNA methylation specifies chromosomal localization of MeCP2. *Mol Cell Biol* 16:414-421
 - Nan XS, Campoy FJ, Bird A (1997) MeCP2 is a transcriptional repressor with abundant binding sites in genomic chromatin. *Cell* 88:471-481
 - Nan XS, Ng HH, Johnson CA, Laherty CD, Turner BM, Eisenman RN, Bird A (1998) Transcriptional repression by the methyl-CpG-binding protein MeCP2 involves a histone deacetylase complex. *Nature* 393:386-389
 - Narayanan V, Olinsky S, Dahle E, Naidu S, Zoghbi HY (1998) Mutation analysis of the M6b gene in patients with Rett syndrome. *Am J Med Genet* 78:165-168
 - Ng HH & Bird (1999) A DNA methylation and chromatin modification *Curr Opin Genet Dev* 9:158-163
-

- Ng HH, Jeppesen P, Bird A (2000) Active repression of methylated genes by the chromosomal protein MBD1. *Mol Cell Biol* 20:1394-1406
- Nielsen JB, Henriksen KF, Hansen C, Silahatoglu A, Schwartz M, Tommerup N (2001) MECP2 mutations in Danish patients with Rett syndrome: high frequency of mutations but no consistent correlations with clinical severity or with the X chromosome inactivation pattern. *Eur J Hum Genet* 9:178-184
- Nomura T, Khan MM, Kaul SC, Dong HD, Wadhwa R, Colmenares C, Kohno I, Ishii S (1999) Ski is a component of the histone deacetylases complex required for transcriptional repression by Mad and thyroid hormone receptor. *Genes Dev* 13:412-423
- Noumova AK, Plengue RM, Bird LM, Leppert M, Morgan k, Willard HF, Sapienza C (1996) Heritability of X chromosome-inactivation phenotype in a large family. *Am J Hum Genet* 58:1111-1119
- Oberlé I, Rousseau F, Heitz D, Kretz C, Devys D, Hanauer A, Boue J, Bertheas MF, Mandel JL (1991) Instability of a 550-base pair DNA segment and abnormal methylation in fragile X syndrome. *Science* 252:1097-1102
- Ohki I, Shimatoku N, Fujita N, Nakao M, Shirakawa M (1999) Solution structure of the methyl-CpG-binding domain of the methylation-dependent transcriptional repressor MBD1. *EMBO J* 18:6653-6661
- Orrico A, Lam C, Galli L, Dotti MT, Hayek G, Tong SF, Poon PM, Zappella M, Federico A, Sorrentino V (2000) MECP2 mutation in male patients with non-specific X-linked mental retardation. *FEBS Lett* 481:285-288
- Paratcha G, Stein ML, Szapiro G, Lopez M, Bevilacqua L, Cammarota M, Iraldi AP, Izquierdo I, Medina JH (2000) Experience-dependent decrease in synaptically localized Fra-1. *Mol Brain Res* 70:120-130
- Percy AK, Schanen C, Dure LS (1998) The genetic basis of Rett syndrome: Candidate gene considerations. *Mol Genet Metab* 64:1-6
- Prokhortchouk A, Hendrich B, Jorgensen H, Ruzov A, Wilm M, Georgiev G, Bird A, Prokhortchouk E (2001) The p120 catenin partner Kaiso is a DNA methylation-dependent transcriptional repressor. *Genes Dev* 15:1613-1618
- Quaderi NA, Meehan RR, Tate PH, Cross Sh, Bird AP, Chatterjee A, Herman GE, Brown SD (1994) Genetic and physical mapping of a gene encoding a methyl CpG binding protein, *Mecp2*, to the mouse X chromosome. *Genomics* 22:648-651
- Reichwald K, Thiesen J, Wiehe T, Weitzel J, Poustka WA, Rosenthal A, Platzer M, Stratling WH, Kioschis P (2000) Comparative sequence analysis of the MECP2-locus in human and mouse reveals new transcribed regions. *Mamm Genome* 11:182-90
- Rett A (1966) Über ein eigenartiges hirnatrophisches syndrom bei hyperammonemie im kindesalter. *Wien Med Wochenschr* 116:723-728

-
- Roloff TC, Ropers HH, Nuber UA (2003) Comparative study of methyl-CpG-binding domain proteins. *BCM Genomics* 4:1
 - Schanen NC, Dahle EJ, Capozzoli F, Holm VA, Zoghbi HY, Francke U (1997) A new Rett syndrome family consistent with X-linked inheritance expands the X chromosome exclusion map. *Am J Hum Genet* 61:634-641
 - Schanen C & Francke U (1998) A severity affected male born into a Rett syndrome kindred supports X-linked inheritance and allows extension of the exclusion map. *Am J Hum Genet* 63:267-269
 - Schanen C (2001) Rethinking the fate of males with mutations in the gene that causes Rett syndrome. *Brain Dev* 23:S144-S146
 - Schwartzman JS, Bernardino A, Nishimura A, Gomes RR, Zats M (2001) Rett syndrome in a boy with a 47,XXY karyotype confirmed by a rare mutation in the MECP2 gene. *Neuropediatrics* 32:162-164
 - Shahbazian MD & Zoghbi HY (2001) Molecular genetics of Rett syndrome and clinical spectrum of MECP2 mutations. *Curr Opin Neurol* 14:171-176
 - Shahbazian MD & Zoghi H (2002) Rett syndrome and MeCP2: linking epigenetics and neuronal function. *Am J Hum Genet* 71:1259-1272
 - Shahbazian MD, Antalffy BA, Armstrong D, Zoghi H (2002a) Insight into Rett syndrome: MeCP2 levels display tissue- and cell-specific differences and correlate with neuronal maturation. *Hum Mol Genet* 11:115-124
 - Shahbazian MD, Sun Y, Zoghi H (2002b) Balanced X chromosome inactivation patterns in the Rett syndrome brain. *Am J Med Genet* 111:164-168
 - Shahbazian MD, Young JI, Yuva-Paylor LA, Spencer CM, Antalffy BA, Noebels JL, Armstrong D, Paylor R, Zoghi H (2002c) Mice with truncated MeCP2 recapitulate many Rett syndrome features and display hyperacetylation of histone H3. *Neuron* 35:243-254
 - Sirianni N, Naidu S, Pereira J, Pillotto RF, Hoffman EP (1998) Rett syndrome: confirmation of X-linked dominant inheritance, and localization of the gene to Xq28. *Am J Hum Genet* 63:1552-1558
 - Song HR & Schanen NC (1998) Search for mutation in the GdX in Rett syndrome. In 48th Annual Meeting of the American Society of Human Genetics, Denver, CO
 - Song HR, Gu A, Schanen NC (1999) Identification of a second polymorphism in the human serotonin receptor ν 2C gene (5-HT ν 2C). *Mol Genet Metab* 66:224-227
 - Summer C, Shinohara T, Durham L, Traub R, Major EO, Amemiya K (1996) Expression of multiple classes of the Nuclear Factor-1 family in the developing human brain: differential expression of two classes of NF-1 gene. *J Neurovirol* 2:87-100.
-

- Takahashi Y, Fukushige S, Murotsu T, Matsubara K (1986) Structure of human cholecystokinin gene and its chromosomal location. *Gene* 50:353-60.
- Tang J, Qi Y, Bao XH, Wu XR (1997) Mutational analysis of mitochondrial DNA of children with Rett syndrome. *Pediatric Neurol* 17:327-330
- Tate P, Skarnes W, Bird A (1996) The methyl-CpG binding protein MeCP2 is essential for embryonic development in the mouse. *Nat Genet* 12:205-208
- Tatematsu K, Yamazaki T, Ishikawa F (2000) MBD2-MBD3 complex binds to hemimethylated DNA and forms a complex containing DNMT1 at the replication foci in late S phase. *Genes to Cells* 5:677-688.
- The Huntington Disease Collaborative Research Group (1993) A novel gene containing a trinucleotide repeat that is expanded and unstable on Huntington's disease chromosomes. *Cell* 72:971-983
- Thomas GH (1996) High male:female ratio of germ-line mutations: an alternative explanation for postulated gestational lethality in males in X-linked dominant disorders. *Am J Hum Genet* 58:1364-1368
- Topçu M, Akyerli C, Sayi A, Toruner GA, Kocoglu SR, Cimbis M, Ozcelik T (2002) Somatic mosaicism for a MECP2 mutation associated with classic Rett syndrome in a boy. *Eur J Hum Genet* 10:77-81
- Trappe R, Laccone F, Cobilanschi J, Meins M, Huppke P, Hanefeld F, Engel W (2001) MECP2 mutations in sporadic cases of Rett syndrome are almost exclusively of paternal origin. *Am J Hum Genet* 68:1093-2001
- Turner H, MacDonald F, Warburton S, Latif F, Webb T (2003) Developmental delay and the methyl binding genes. *J Med Genet* 40:E13
- Vacca M, Filippini F, Budillon A, Rossi V, Della Ragione F, De Bonis ML, Mercadante G, Manzati E, Gualandi F, Bigoni S, TrabANELLI C, Pini G, Calzolari E, Ferlini A, Meloni I, Hayek G, Zappella M, Renieri A, D'Urso M, D'Esposito M, Macdonald F, Kerr A, Dhanjal S, Hulten M (2001) MECP2 gene mutation analysis in the British and Italian Rett Syndrome patients: hot spot map of the most recurrent mutations and bioinformatic analysis of a new MECP2 conserved region. *Brain Dev* 23 Suppl 1:S246-S250
- Van der Veyver IB, Subramanian S, Zoghbi HY (1998) Genomic structure of a human holocytochrome c-type synthetase gene in Xp22.3 and mutation analysis in patients with Rett syndrome. *Am J Med Genet* 78:179-181
- Verma NP, Chheda RL, Nigro MA, Hart ZH (1986) Electroencephalographic findings in Rett syndrome. *Electroencephalogr Clin Neurophysiol* 64:394-401
- Vilain A, Apiou F, Vogt N, Dutrillaux B, Malfoy B (1996) Assignment of the gene for methyl-CpG-binding protein 2 (MECP2) to human chromosome band Xq28 by in situ hybridization. *Cytogenet Cell Genet* 74:293-294

-
- Villard L, Kpebe A, Cardoso AK, Chelly PJ, Tardieu M, Fontes M (2000) Two affected boys in a Rett syndrome family: clinical and molecular findings. *Neurology* 55:1188-1193
 - Villard L, Lévy N, Xiang F, Kpebe A, Labelle V, Chevillard C, Zhang Z, Schwartz CE, Tardieu M, Chelly J, Anvret M, Fontes M (2001) Segregation of a totally skewed pattern of X chromosome inactivation in four familial cases of Rett syndrome without MECP2 mutation: implications for the disease. *J Med Genet* 38:435-442
 - Vorsanova SG, Yurov YB, Ulas VY, Demidova IA, Sharonin VO, Kolotti AD, Gorbatchevskaia NL, Beresheva AK, Soloviev IV (2001) Cytogenetic and molecular-cytogenetic studies of Rett syndrome (RTT): a retrospective analysis of a Russian cohort of RTT patients (the investigation of 57 girls and three boys). *Brain Dev* 23 Suppl 1:S196-S201
 - Vourch P, Bienvenu T, Beldjord C, Chelly J, Barthelemy C, Muh JP, Andres C (2001) No mutations in the coding region of the Rett syndrome gene MECP2 in 59 autistic patients. *Eur J Hum Genet* 9:556-558
 - Wade PA (2001) Methyl CpG-binding proteins and transcriptional repression. *Bioessays* 23:1131-1137.
 - Wakefield IR, Smith BO, Nan X, Free A, Soteriu A, Uhrin D, Bird AP, Barlow PN (1999) The solution structure of the domain from MeCP2 that binds to methylated DNA. *J Mol Biol* 291:1055-1065
 - Wan M, Francke U (1998) Evaluation of two X chromosomal candidate genes for Rett syndrome: glutamate dehydrogenase-2 (GLUD2) and rab GDP-dissociation inhibitor (GDI1). *Am J Med Genet* 78:169-172
 - Wan M, Sung Jae Lee S, Zhang X, Houwink-Manville I, Song H-R, Amir R, Budden S, Naidu S, Pereira JL, Lo IF, Zoghbi HY, Schanen NC, Francke U (1999) Rett Syndrome and beyond: recurrent spontaneous and familial MECP2 mutations at CpG hotspots. *Am J Hum Genet* 65:1520-1529
 - Wan M, Zhao K, Sung Jae Lee S, Francke U (2001) MECP2 truncating mutations cause histone H4 hyperacetylation in Rett syndrome. *Hum Mol Genet* 10:1085-1092
 - Watson P, Black G, Ramsden S, Barrow M, Super M, Kerr B, Clayton-Smith J (2001) Angelman syndrome phenotype associated with mutations in MECP2, a gene encoding a methyl CpG binding protein. *J Med Genet* 38:224-228
 - Weaving LS, Williamson SL, Bennetts B, Davis M, Ellaway CJ, Leonard H, Thong MK, Delatycki M, Thompson EM, Laing N, Christodoulou J (2003) Effects of MECP2 mutation type, location and X-inactivation in modulating Rett syndrome phenotype. *Am J Med Genet* 118A:103-114
 - Webb T, Watkiss E, Woods CG (1993) Neither uniparental disomy nor skewed X-inactivation explains Rett syndrome. *Clin Genet* 44:236-240

- Webb T & Watkiss E (1996) A comparative study of X-inactivation in Rett syndrome probands and control subjects. *Clin Genet* 49:198-195
- Webb T, Clarke A, Hanefeld F, Pereira JL, Rosenbloom L, Woods CG (1998) Linkage analysis in Rett syndrome families suggests that there may be a critical region at Xq28. *J Med Genet* 35:997-1003
- Webb T & Latif F (2001) Rett syndrome and the MECP2 gene. *J Med Genet* 38:217-223
- Weir BS (1990) Genetic Data Analysis. Sinauer Associates, Inc. Publishers. Sunderland, Massachusetts.
- Whitaker-Azmitia, PM (2001) Serotonin and brain development: Role in human developmental diseases. *Brain Research Bulletin* 56:479-485
- Whitty CJ, Kapatos G, Bannon MJ (1993) Neurotrophic effects of substance P on hippocampal neurons in vivo. *Neurosci Lett* 164:141-144
- Winnepeninckx B, Errijgers V, Hayez-Delatte F, Reyniers E, Frank-Kooy R (2002) Identification of a family with nonspecific mental retardation (MRX79) with the A140V mutation in the MECP2 gene: is there a need for routine screening? *Hum Mutat* 20:249-52
- Xiang F, Zhang Z, Clarke A, Joseluiz P, Sakkubai N, Saroniji B, Delozier-Blanchet CD, Hansmann I, Edstrom L, Anvret M (1998) Chromosome mapping of Rett syndrome: a likely candidate region on the telomere of Xq. *J Med Genet* 35:297-300
- Xiang F, Buervenich S, Prieto N, Bailey MES, Zhang Z, Anvret M (2000) Mutation screening in Rett syndrome patients. *J Med Genet* 37:250-255
- Xiang F, Stenbom Y, Anvret M (2002) MECP2 mutations in Swedish Rett syndrome clusters. *Clin Genet* 61:384-385
- Yamashita Y, Kondo I, Fukuda T, Morishima R, Kusaga A, Iwanaga R, Matsuishi T (2001) Mutation analysis of the methyl-CpG-binding protein 2 gene (MECP2) in Rett syndrome with preserved speech. *Brain Dev* 23 Suppl 1:S157-S160
- Yntema HG, Oudakker AR, Kleefstra T, Hamel BCJ, Bokhoven H, Chelly J, Kalscheuer VM, Fryns JP, Raynaud M, Moizard MP, Moraine C (2002a) In-frame deletion in MECP2 causes mild nonspecific mental retardation. *Am J Med Genet* 107:81-83
- Yntema HG, Kleefstra T, Oudakker AR, Romein T, de Vries BBA, Nillesen W, Sistermans EA, Bruner HG, Hamel BCJ, Bokhoven H (2002b) Low frequency of MECP2 mutations in mentally retarded males. *Eur J Hum Genet* 10:487-490
- Yu F, Thiesen J, Stratling WH (2000) Histone deacetylase-independent transcriptional repression by methyl-CpG-binding protein 2. *Nucleic Acids Res* 28:2201-2206
- Yusufzai TM & Wolffe AP (2000) Functional consequences of Rett syndrome mutations on human MeCP2. *Nucleic Acids Res* 28:4172-4179

- Zappella M (1992) The Rett girls with preserved speech. *Brain Dev* 14:98-101
- Zappella M, Meloni I, Longo I, Hayek G, Renieri A (2001) Preserved speech variants of the Rett syndrome: molecular and clinical analysis. *Am J Med Genet* 104:14-22
- Zappella M, Meloni I, Longo I, Canitano R, Hayek G, Rosaia L, Mari F, Renieri A (2003) Study of MECP2 gene in Rett syndrome variants and autistic girls. *Am J Med Genet* 15;119B:102-7
- Zeev BB, Yaron Y, Schanen NC, Wolf H, Brandt N, Ginot N, Shomrat R, Orr-Urtreger A (2002) Rett syndrome: clinical manifestations in males with MECP2 mutations. *J Child Neurol* 17:20-24
- Zoghbi HY, Percy AK, Schultz RJ, Fill C (1990) Patterns of X chromosome inactivation in the Rett syndrome. *Brain Dev* 12:131-135

