



UNIVERSITAT DE BARCELONA

Facultat de Farmàcia

Departament de Microbiologia i Parasitologia Sanitàries

Estudi dels polihidroxialcanoats acumulats per *Pseudomonas aeruginosa* 42A2: producció i caracterització

Mònica Bassas i Galià 2007



UNIVERSITAT DE BARCELONA

FACULTAT DE FARMÀCIA

Departament de Microbiologia i Parasitologia Sanitàries

Programa de doctorat: Química Orgànica (Facultat de Químiques)

BIENNI 2002-2004

Estudi dels polihidroxialcanoats acumulats per *Pseudomonas aeruginosa* 42A2: producció i caracterització

Memòria presentada per **Mònica Bassas i Galià** per optar al títol de doctor per la Universitat de Barcelona

Director/a:

Dra. Àngels Manresa i Presas

Dr. Joan Llorens i Llacuna

Doctorand:

Mònica Bassas i Galià

Mònica Bassas i Galià, 2007

6. BIBLIOGRAFIA

- Agbenyega, J. k., M. Claybourn and G. Ellis (1991). "A study of the autoxidation of some unsaturated fatty acid methyl esters using Fourier Transform Raman Spectroscopy." Spectrochimica Acta 47A(9/10): 1375-1388.
- Ahimou, F., F. A. Denis, A. Touhami and Y. F. Dufrene (2002). "Probing microbial cell surface charges by atomic force microscopy." Langmuir 18: 9937-9941.
- Akita, S., Y. Einaga, Y. Miyaki and H. Fujita (1976). "Solution properties of poly(D- β -hydroxybutyrate). 1. Biosynthesis and characterization." Macromolecules 9(5): 774-780.
- Álvarez, H. M., F. Mayer, D. Fabritius and A. Steinbüchel (1996). "Formation of intracytoplasmic lipid inclusions by *Rhodococcus opacus* strain PD630." Archives of Microbiology 165: 377-386.
- Álvarez, H. M., O. H. Pucci and A. Steinbüchel (1997). "Lipid storage compounds in marine bacteria." Applied Microbiology and Biotechnology 47: 132-139.
- Anderson, A. J. and E. A. Dawes (1990). "Ocurrence, metabolism, metabolic role and industrial uses of bacterial polyhydroxyalkaonates." Microbiological Reviews 54(4): 450-472.
- Andrade, A. P., P. Neuenschwander, R. Hany, T. Egli and B. Witholt (2002). "Synthesis and Characterization of novel copoly(ester-urethane) containing blocks of Poly-[*(R*)-3-hydroxyoctanoate] and Poly-[*(R*)-3-hydroxybutyrate]." Macromolecules 35: 4946-4950.
- Ashby, R., D. Solaiman and T. Foglia (2002). "The synthesis of short- and medium-chain-length poly(hydroxyalkanoate) mixtures from glucose -or alkanoic acid- grown *Pseudomonas oleovorans*." Journal of Industrial Microbiology & Biotechnology 28: 147-153.
- Ashby, R. D., A. Cromwick and T. A. Foglia (1998). "Radiation crosslinking of a bacterial medium-chain-length poly(hydroxyalkanoate) elastomer from tallow." International Journal of Biological Macromolecules 23: 61-72.
- Ashby, R. D., T. A. Foglia, C.-K. Liu and J. W. Hampson (1998). "Improved film properties of radiation treated medium chain length poly(hydroxyalkanoates)." Biotechnology Letters 20(11): 1047-1052.
- Ashby, R. D., T. A. Foglia, D. K. Y. Solaiman, C.-K. Liu, A. Núñez and G. Eggink (2000). "Viscoelastic properties of linseed oil-based medium-chain-length poly(hydroxyalkanoate) films: Effects of epoxidation and curing." International Journal of Biological Macromolecules 27: 355-368.
- Ballistreri, A., G. Montaudo, M. Giuffrida and R. W. Lenz (1992). "Determination of sequence distributions in bacterial copolymers containing higher alkyl and alkenyl pendant groups." Macromolecules 25: 1845-1851.

- Barbuzzi, T., M. Giuffrida, S. Carnazza, A. Ferreri, S. P. P. Guglielmino and A. Ballistreri (2004). "Microbial Synthesis of Poly(3-hydroxyalkanoates) by *Pseudomonas aeruginosa* from fatty acids: Identification of higher monomer units and structural characterization." *Biomacromolecules* 5: 2469-2478.
- Bassas, M., E. Rodríguez, J. Llorens and À. Manresa (2006). "Poly(3-hydroxyalkanoate) produced from *Pseudomonas aeruginosa* 42A2 (NCBIM 40045): Effect of fatty acid nature as nutrient." *Journal of Non-Crystalline Solids* 352: 2243-2247.
- Bear, M. M., Leboucher-Durand, M.A., Langlois, V., Lenz, R.W., Goodwin,S. and Guerin, P. (1997). "Bacterial poly-3-hydroxyalkenoates with epoxy groups in the side chains." *Reactive and Functional polymers* 34: 65-67.
- Binning, G., C. F. Quate and C. Gerber (1986). "Atomic force microscopy." *Phys. Rev. Lett.* 56: 930-933.
- Bolshakova, A. V., O. I. Kiselyova and I. V. Yaminsky (2004). "Microbial surfaces investigated using atomic force microscopy." *Biotechnology Progress* 20(6): 1615-1622.
- Boopathy, R. (2000). "Factor limiting bioremediation technologies." *Bioresource Technology* 74: 63-67.
- Brandl, H., R. A. Gross, R. W. Lenz and R. C. Fuller (1988). "Pseudomonas oleovorans as a source of poly(β -hydroxyalkanoates) for potential application as biodegradable polyester." *Applied and Environmental Microbiology* 54: 1977-1982.
- Cabellos, A. (2003). "Obtención y Caracterización del biopolímero producido por *Pseudomonas aeruginosa* 42A2 a partir de ácido undecanoico." *Proyecto Final de Carrera. Ingenieria Química. Universitat de Barcelona.*
- Casini, E., C. de Rijk, P. de Waard and G. Eggink (1997). "Synthesis of Poly(hydroxyalkanoate) from Hydrolyzed Linseed Oil." *Journal of Environmental Polymer Degradation*. 5(3): 153-158.
- Cromwick, A., T. Foglia and R. W. Lenz (1996). "The microbial production of poly(hydroxyalkanoates) from tallow." *Applied Microbiology and Biotechnology* 46: 464-469.
- Culleré, J. (2002). "Utilització de l'àcid oleic per *Pseudomonas* sp.42A2." *Tesis Doctoral. Universitat de Barcelona.*
- Culleré, J., O. Durany, M. Busquets and À. Manresa (2001). "Biotransformation of oleic acid into (E)-10-hydroxy-8-octadecenoic acid and (E)-7,10-dihydroxy-8-octadecenoic acid by *Pseudomonas* sp.42A2 in an immobilized system." *Biotechnology Letters* 23: 215-219.
- Curley, J. M., R. W. Lenz and R. C. Fuller (1996). "Sequential production of two different polyesters in the inclusion bodies of *Pseudomonas oleovorans*." *International Journal of Biological Macromolecules* 19: 29-34.

- Choi, J.-I. and S. Y. Lee (1999). "High-level production of poly-(3-hydroxybutyrate-co-3-hydroxyvalerate) by fed-batch culture of recombinant *Escherichia coli*." Applied and Environmental Microbiology 65(10): 4363-4368.
- Dawes, E. A. (1973). "The role and regulation of energy reserve polymers in microorganisms." Adv. Microb. Physiol. 10: 135-266.
- De Koning, G. (1995). "Physical properties of bacterial poly((R)-3-hydroxyalkanoates)." Can. J. Microbiol. 41(suppl.1): 303-309.
- De Koning, G. J. M., M. Kellerhals, C. van Meurs and Witholt, B. (1997a). "A process for the recovery of poly(hydroxyalkanoates) from *Pseudomonads* Part 2: Process development and economic evaluation." Bioprocess Engineering 17: 15-21.
- De Koning, G. J. M., Witholt, B. (1997b). "A process for the recovery of poly(hydroxyalkanoates) from *Pseudomonads*. Part 1: Solubilization." Bioprocess and Biosystems Engineering 17: 7-13.
- De Smet, M. J., G. Eggink, B. Witholt, J. Kingma and H. Wynberg (1983). "Characterization of intracellular inclusions formed by *Pseudomonas oleovorans* during growth on octane." Journal of Bacteriology 154(2): 870-878.
- De Waard, P., H. van der Wal, G. M. N. Huijborts and G. Eggink (1993). "Heteronuclear NMR analysis of unsaturated fatty acids in poly(3-hydroxyalkanoates)." The Journal of Biological Chemistry 268(1): 315-319.
- Dennis, D., Liebig, C., Holley, t. and al. (2003). "Preliminary analysis of polyhydroxyalkanoate inclusions using atomic force microscopy." FEMS Microbiology Letters 226: 113-119.
- Doi, Y. and C. Abe (1990). "Biosynthesis and characterization of a new bacterial copolyester of 3-hydroxy- ω -chloroalkenoates." Macromolecules 23: 3703-3707.
- Du, G. and J. Yu (2002). "Green Technology for Conversion of Food Scraps to Biodegradable Thermoplastic Polyhydroxyalkanoates." Environ. Sci. Technol. 36: 5511-5516.
- Dufresne, A., L. Reche, R. H. Marchessault and M. Lacroix (2001). "Gamma-ray crosslinking of poly(3-hydroxyoctanoate-co-undecenoate)." International Journal of Biological Macromolecules 29: 73-82.
- Eggink, G., P. de Waard and G. N. M. Huijborts (1995). "Formation of novel poly(hydroxyalkanoates) from long-chain fatty acids." Canadian Journal of Microbiology 41 (suppl 1): 14-21.
- Eggink, G., P. d. Waard and G. N. M. Huijborts (1992). "The role of fatty acid biosynthesis and degradation in the supply of substrates for poly(3-hydroxyalkanoate) formation in *Pseudomonas putida*." FEMS Microbiological reviews 103: 159-164.
- Eglinton, G., H. Hunneman and A. McCormick (1968). "Gas chromatographic-mass spectrometric studies of long chain hydroxy acid. III. The mass spectra of the methyl

esters trimethylsilyl ethers of aliphatic hydroxy acids. A facile method of double bond location." Org. Mass Spec. 1: 593-611.

El Kirat, K., I. Burton, V. Dupres and D. Y.F (2005). "Sample preparation procedures for biologiacal atomic force microscopy." Journal of Microscopy 219(3): 199-207.

Fernández, D. (2000). "Producción y caracterización de los polihidroxialcanoatos producidos por *Pseudomonas sp.* 42A2." Máster Experimental en Biología. Universitat de Barcelona.

Fernández, D., E. Rodríguez, M. Bassas, M. Viñas, A. M. Solanas, J. Llorens, M. A.M. and A. Manresa (2005). "Agro-industrial oily wastes as substrates for PHA production by the new strain *Pseudomonas aeruginosa* NCIB 40045: Effect of culture conditions." Biochemical Engineering Journal 26(2-3): 159-167.

Fernandez del Castillo, R., F. Rodriguez Valera, J. Gonzalez Ramos and F. Ruiz Berraquero (1986). "Accumulation of Poly(β -Hydroxybutyrate) by Halobacteria." Applied and Environmental Microbiology 51(1): 214-216.

Fiedler, S., A. Steinbüchel and B. Rehm (2002). "The role of the fatty acid b-oxidation multienzyme complex from *Pseudomonas oleovorans* in polyhydroxyalkanoate biosynthesis: molecular characterization of the *fadBA* operon from *P.oleovorans* and of the enoyl-CoA hydratase genes *phaJ* from *P.oleovorans* and *Pseudomonas putida*." Arch Microbiol 178: 149-160.

Findlay, R. H. and D. C. White (1983). "Polymeric β -hydroxyalkanoates from environmental samples and *Bacillus megaterium*." Applied and Environmental Microbiology 45(1): 71.

Flechter, A. (1993). "Plastics from Bacteria and for Bacteria: PHA as Natural, Biodegradable Polyesters." Springer Verlag, New York: 77-93.

Fritzsche, K., R. W. Lenz and R. C. Fuller (1990). "Bacterial polyesters containing branched poly (β -hydroxyalkanoate) units." J. Biol. Macromol Chem. 191: 92-101.

Fritzsche, K., R. W. Lenz and R. C. Fuller (1990). "An unusual bacterial polyester with a phenyl pendant group." Macromol. Chem. 191: 1957-1965.

Fukui, T., M. Kato and H. Matsusaki (1998). "Morphological and ^{13}C -nuclear magnetic resonance studies for polyhydroxyalkanoates biosynthesis in *Pseudomonas sp.61-3*." FEMS Microbiology Letters 164: 219-225.

Furrer, P., R. Hany, D. Rentsch, A. Grubelnik, K. Ruth, S. Pake and M. Zinn (2007). "Quantitative analysis of bacterial medium-chain-length pol([R]-3-hydroxyalkanoates) by gas chromatography." Journal of Chromatography A 1143: 199-206.

Garcia-Ribera, R., M. Monteoliva-Sanchez and A. Ramos-Comerzana (2001). "Production of polyhydroxyalkanoates by *Pseudomonas putida* KT2442 harboring pSK2665 in wastewater from olive oil mills (alpechin)." Electronic Journal of Biotechnology 4(2): 1-4.

- Gerngross, T. U., P. Reilly, P. Stubbe, A. J. Sinskey and O. P. Peoples (1993). "Immunocytochemical analysis of poly- β -hydroxybutyrate (PHB) synthase in *Alcaligenes eutrophus* H16: localization of the synthase enzyme at the surface of PHB-granules." *Journal of Bacteriology* 175: 5289-5293.
- Gogolewski, S., M. Jovanovic, S. M. Perren, J. G. Dillon and M. K. Hughes (1993). "Tissue response and in vivo degradation of selected polyhydroxyacids: polylactides (PLA), poly(3-hydroxybutyrate) (PHB), and poly(3-hydroxybutyrate-co-3-hydroxyvalerate) (PHB/VA)." *Journal of biomedical materials research* 27(9): 1135-1148.
- Gorenflo, V., A. Steinbüchel, S. Marose, M. Reiseberg and T. Scheper (1999). "Quantification of bacterial polyhydroxyalkanoic acids by Nile red staining." *Applied Microbiology and Biotechnology* 51: 765-772.
- Griebel, R., Z. Smith and J. M. Merrick (1968). "Metabolism of poly- β -hydroxybutyrate granules from *Bacillus megaterium*." *Biochemistry* 7: 3676-3681.
- Gross, R. A., C. DeMello and R. W. Lenz (1989). "Biosynthesis and characterization of poly(β -hydroxyalkanoates) produced by *Pseudomonas oleovorans*." *Macromolecules* 22: 1106-1115.
- Guerrero, A., I. Casals, M. Busquets, Y. León and A. Manresa (1997). "Oxidation of oleic acid to (E)-10-hydroperoxy-8-octadecenoic acid and (E)-10-hydroxy-8-octadecenoic acids by *Pseudomonas sp.* 42A2." *Biochimica et Biophysica Acta* 1347: 75-81.
- Haba, E., J. Vidal-Mas, M. Bassas, M. J. Espuny, J. Llorens and A. Manresa (2007). "Poly 3-(hydroxyalkanoates) produced from oily substrates by *Pseudomonas aeruginosa* 47T2 (NCBIM 40044): Effect of nutrients and incubation temperature on polymer composition." *Biochem. Eng. J.* 35(2): 99-106.
- Handrick, R., S. Reinhardt and D. Jendrossek (2000). "Mobilization of poly(3-hydroxybutyrate) in *Ralstonia eutropha*." *Journal of Bacteriology* 182(20).
- Hanley, S., D. J. C. Pappin, D. Rahman, A. j. White, K. M. Elborough and A. R. Slabs (1999). "Re-evaluation of the primary structure of *Ralstonia eutropha* phasin and implications for polyhydroxyalkanoic acid granule binding." *FEBS Lett.* 447: 99-105.
- Hartmann, R. (2005). "Tailored Medium-Chain-Length Poly[(R)-3-hydroxyalkanoates]: Biosynthetic and Chemical Approaches." *Tesis Doctoral. Swiss Federal Institute of Technology of Zurich.*
- Hartmann, R., R. Hany, T. geiger, T. Egli, B. Witholt and M. Zinn (2004). "Tailored biosynthesis of olefinic medium-chain-length poly[(R)-3-hydroxyalkanoates] in *Pseudomonas putida* GPoI with improved thermal properties." *Macromolecules* 37: 3780-6785.
- Hazer, B., S. I. Demirel, M. Borcakli, M. S. Eroglu, M. Cakmak and B. Erman (2001). "Free radical crosslinking of unsaturated bacterial polyesters obtained from soybean oily acids." *Polymer Bulletin* 46: 389-394.

Hernández, R. (2003). "Utilización de ácidos alcanoicos (C_5-C_9) para la obtención de polihidroxialcanoatos mediante *Pseudomonas aeruginosa* 42A2." Proyecto Final de Carrera. Ingenieria Química. Universitat de Barcelona.

T. Hiraishi, Y. Kikkawa, M. Fujita, N. Mohd, M. Kanesato, K. Tsuge, M. Maeda and Y. Doi (2005). "Atomic force microscopic observation of *in vitro* polymerized poly[(R)-hydroxybutyrate]: insight into possible mechanisms of granule formation." Biomacromolecules 6: 2671-2677.

Hirt, T. D., P. Neuenschwander and U. W. Suter (1996a). "Synthesis of degradable, biocompatible, and tough block-copolyesterurethanes." Macromol. Chem. Phys. 197: 4253-4268.

Hirt, T. D., P. Neuenschwander and U. W. Suter (1996b). "Telechelic diols from poly[(R)-3-hydroxybutyric acid] and poly[(R)-3-hydroxybutyric acid]-co-poly[(R)-3-hydroxyvaleric acid]." Macromol. Chem. Phys. 197: 1609-1614.

Hori, K., K. Soga and Y. Doi (1994). "Production of poly(3-hydroxyalkanoates-co-3-hydroxy- ω -fluoroalkanoates) by *Pseudomonas oleovorans* from 1-fluorononane and glucomate." Biotechnology Letters 16(5): 501.

Huijberts, G. N., G. Eggink, P. Waard, G. W. Huisman and B. Witholt (1992). "*Pseudomonas putida* KT2442 cultivated on glucose accumulates poly(3-hydroxyalkanoates) consisting of saturated and unsaturated monomers." Applied and Environmental Microbiology 58(2): 536-544.

Huijberts, G. N. M. and G. Eggink (1996). "Production of poly(3-hydroxyalkanoates) by *Pseudomonas putida* KT2442 in continuous cultures." Applied Microbiology and Biotechnology 46: 233-239.

Huisman, G. W., O. Leeuw, G. Eggink and B. Witholt (1989). "Synthesis of poly-3-hydroxyalkanoates is a common feature of fluorescent *Pseudomonads*." Applied and Environmental Microbiology 55(8): 1949-1954.

Johnstone, B. (1990). "A throw away answer." Far Eastern Econ. Rev. 147(6): 62-63.

Jung, Y. M., J. S. Park and Y. H. Lee (2000). "Metabolic engineering of *Alcaligenes eutrophus* through the transformation of cloned phvCAB genes for the investigation of the regulatory mechanism of polyhydroxyalkanoate biosynthesis." Enzyme and Microbial Technology 26: 201-208.

Jurasek, L., Marchessault (2004). "Polyhydroxyalkanoate (PHA) granule formation in *Ralstonia eutropha* cells: a computer simulation." Applied Microbiology and Biotechnology 64: 611-617.

Kadouri, D., E. Jurkevitch and Y. Okon (2005). "Ecological and Agricultural Significance of Bacterial Polyhydroxyalkanoates." Critical Reviews in Microbiology 31: 55-67.

Kalia, V. C., N. Raizada and V. Sonakya (2000). "Bioplastics." J. Sci. Ind. Res. 59: 433-445.

- Kelley, A. S. and F. Srienc (1999). "Production of two phase polyhydroxyalkanoic acid granules in *Ralstonia eutropha*." International Journal of Biological Macromolecules 25: 61-67.
- Kidwell, P. A. and K. Bleman (1982). "Determination of double bond position and geometry of olefins by mass spectrometry of their Diels-Alder adducts." Anal. Chem. 54: 2462-2465.
- Kim, B. S. (2000). "Production of poly(3-hydroxybutyrate) from inexpensive substrates." Enzyme and Microbial Technology 27: 774-777.
- Kim, B. S. (2002). "Production of medium chain length polyhydroxyalkanoates by fed-batch culture of *Pseudomonas oleovorans*." Biotechnol Lett. 24: 125-130.
- Kim, B. S. and H. N. Chang (1998). "Production of poly(3-hydroxybutyrate) from starch by *Azotobacter chroococcum*." Biotechnology Letters 20(2): 109-112.
- Kim, D. Y., S. B. Jung, G. G. Choi, Y. B. Kim and Y. H. Rhee (2001). "Biosynthesis of polyhydroxyalkanoate copolyester containing cyclohexyl groups by *Pseudomonas oleovorans*." International Journal of Biological Macromolecules 29: 145-150.
- Kim, D. Y., Y. B. Kim and Y. H. Rhee (1998). "Bacterial poly(3-hydroxyalkanoates) bearing carbon-carbon triple bonds." Macromolecules 31: 4760-4763.
- Kim, D. Y., Y. B. Kim and Y. H. Rhee (2000). "Evaluation of various carbon substrates for the biosynthesis of polyhydroxyalkanoates bearing functional groups by *Pseudomonas putida*." International Journal of Biological Macromolecules 28: 23-29.
- Kim, H. W., G. H.W. and C. T. Hou (2000). Production of isomeric 9,10,13 (9,12,13)-trihidroxy-11-E-(10E)-octadecenoic acid from linoleic acid by *Pseudomonas aerugiosa* PR3. 25: 109-115.
- Kim, O., R. A. Gross and D. R. Rutherford (1995). "Bioengineering of poly(β -hydroxyalkanoates) for advanced material applications: incorporation of cyano and nitrophenoxy side chain substituents." Canadian J. Microbiol 41(suppl.1): 32.
- Kim, Y. B. and R. W. Lenz (2001). "Polyesters from microorganism." Advances in biochemical Engineering Biotechnology 71: 52.
- Kim, Y. B., R. W. Lenz and R. C. Fuller (1992). "Poly(β -hydroxyalkanoate) copolymers containing brominated repeating units produced by *Pseudomonas oleovorans*." Macromolecules 25: 2852-2857.
- Kim, Y. B., Y.-H. Rhee, R. W. Lenz and R. C. Fuller (1997). "Poly(3-hydroxyalkanoate)s Produced by *Pseudomonas oleovorans* Grown by Feeding Nonanoic and 10-Undecenoic Acids in Sequence." Polymer Journal 29(11): 894-898.
- Kinkle, S., M. Dauner, G. Scott, B. Kessler and B. Witholt (2000). "Inactivation of isocitrate lyase leads to increased production of medium-chain-length poly(3-hydroxyalkanoates) in *Pseudomonas putida*." Applied and Environmental Microbiology. 66(3): 909-913.

- Koritala, S. and M. O. Bagby (1992). "Microbial Conversions of Linoleic and Linolenic Acids to Unsaturated Hydroxy Fatty Acids." JAACS 69(6): 575-578.
- Kumar, N. and M. N. V. Ravikumar (2001). "Biodegradable block copolymer." Adv. Drug Deliv. Rev. 53: 23-44.
- Kunioka, M., Y. Kawaguchi and Y. Doi (1989). "Production of biodegradable copolyesters of 3-hydroxybutyrate and 4-hydroxybutyrate by *Alcaligenes eutrophus*." Applied Microbiology and Biotechnology 30: 569-573.
- Kunststoffe (2004). "Oktober Ausgabe. Carl Hanser Verlag, München."
- Kuo, T. M., L. K. Manthey and C. T. Hou (1998). "Fatty acid bioconversions by *Pseudomonas aeruginosa* PR3." JAACS 75(7): 875-879.
- Kurth, N., Renard, E., Brachet, F., Robic, D., Guerin, PH., Bourbouze, R. (2002). "Poly(hydroxyoctanoate) containing pendant carboxylic groups for the preparation of nanoparticles aimed at drug transport and release." Polymer 43: 1095-1101.
- Lageveen, R. G. B., G. W. Huisman, H. Preusting, P. Ketelar, G. Eggink and Witholt (1988). "Formation of polyesters by *Pseudomonas oleovorans*: Effect of the substrates on formation and composition of poly-(R)-3-hydroxyalkanoates and poly-(R)-3-hydroxyalkenoates." Applied and Environmental Microbiology 54: 2924-2932.
- Lazzari, M. a. C., O. (1999). "Drying and oxidative degradation of linseed oil." Polymer Degradation and Stability 65: 303-313.
- Lee, E.-Y. and C.-Y. Choi (1995). "Gas Chromatography-Mass Spectrometric Analysis and Its Application to a Screening Procedure for Novel Bacterial Polyhydroxyalkanoic Acids Containing Long Chain Saturated and Unsaturated Monomers." Journal of Fermentation and Bioengineering 80(4): 408-414.
- Lee, M. Y., W. H. Park and R. W. Lenz (2000). "Hydrophilic bacterial polyesters modified with pendant hydroxyl groups." Polymer 41: 1703-1709.
- Lee, S. Y. (1996). "Bacterial polyhydroxyalkanoates." Biotechnology and Bioengineering 49: 1-14.
- Lee, S. Y. (1996). "Plastic bacteria? Progress and prospects for polyhydroxyalkanoate production in bacteria." TIBTECH 14: 431-438.
- Lee, S. Y. (1999). "Chiral compounds from bacterial polyesters: Sugars to plastics to fine chemicals." Biotechnology and Bioengineering 65: 363.
- Lee, S. Y., J. I. Choi and H. H. Wong (1999). "Recent advances in polyhydroxyalkanoate production by bacterial fermentation: mini-review." International Journal of Biological Macromolecules 25: 31-36.
- Lenz, R. W., Y. B. Kim and R. C. Fuller (1992). "Production of unusual bacterial polyesters by *Pseudomonas oleovorans* through cometabolism." FEMS Microbiology Reviews 103: 207-214.

- Madison, L. L. and G. W. Huisman (1999). "Metabolic engineering of poly(3-hydroxyalkanoates): from DNA to plastic." Microbiology and Molecular Biology Reviews 63(1): 21-53.
- Maehara, A., Y. Doi, Z. Nishiyama, Y. Takagi, S. Ueda, S. Nakano and T. Yamane (2001). "PhaR: a protein of unknown function conserved among short-chain-length polyhydroxyalkanoic acids producing bacteria, is a DNA-binding protein and represses *Paracoccus denitrificans* phaP expression in vitro." FEMS Microbiology Letters 200: 9-15.
- Mallégol, J., L. Gonon, J. Lemaire and J.-L. Gardette (2001). "Long-term behaviour of oil-based varnishes and paints. 4. Influence of film thickness on the photooxidation." Polymer Degradation and Stability 72: 191-197.
- Matsumoto, K., Matsusaki, H., Taguchi, K., Seki, M. and Doi, Y. (2002). "Isolation and characterization of polyhydroxyalkanoates inclusions and their associated proteins in *Pseudomonas* sp.61-3." Biomacromolecules 3: 787-792.
- Marqués, A. M. (1982). "Aislamiento y caracterización de *Pseudomonas aeruginosa* procedentes de distintos hábitats: Aspectos ecológicos y sanitarios." Tesis Doctoral. Departament de Microbiologia i Parasitologia Sanitàries. Facultat de Farmàcia. Universitat de Barcelona.
- McCloskey, J. A. (1969). "Mass spectrometry of lipids and steroids." Methods of Enzymology 14: 382-450.
- Moore, C. J., S. L. Moore, M. K. Leecaster and S. B. Weisberg (2001). "A comparision of plastic and plankton in the North Pacific Central Gyre." Marine Pollution Bulletin 49: 1298-1300.
- Numata, K., Y. Kikkawa, T. Tsuge, T. Iwata, Y. Doi and H. Abe (2005). "Enzymatic degradation processes of poly[(R)-3-hydroxybutyric acid] and poly[(R)-3-hydroxybutyric acid-co-(R)-3-hydroxyvaleric acid] single crystals revealed by atomic force microscopy: effect of molecular weight and the second monomer composition on erosion rates." Biomacromolecules 6: 2008-2016.
- Nuñez, M. J., M. O. Martin, P. H. Chan, L.-K. Duonog and A. R. Sindhuurakar (2005). "Atomic force microscopy of bacterial communities in Environmental Microbiology." AC. Press.
- Page, W. J. (1992). "Suitability of commercial beet molasses fractions as substrates for polyhydroxyalkanoate production by *Azotobacter vinelandii* UWD." Biotechnology Letters 14(5): 385-390.
- Park, J.S., H. C. Park, T. L. Huh and Y. H. Lee (1995). "Production of poly- β -hydroxybutyrate by *Alcaligenes eutrophus* transformants harbouring cloned *phbCAB* genes." Biotechnology Letters 17(7): 735-740.
- Park, H. G., Lenz, R.W. and Goodwin, S. (1998). "Epoxidation of bacterial polyesters with unsaturated side chains.I. Production and epoxidation of polyesters from 10-undecenoic acid." Macromolecules 31: 1480-1486.

Peters, V. and B. H. A. Rehm (2005). "In vivo monitoring of PHA granule formation using GFP-labeled PHA synthases." FEMS Microbiology Letters 248: 93-100.

Philip, S., T. Keshavarz and I. Roy (2007). "Review. Polyhydroxyalkanoates: biodegradable polymers with a range of applications." Journal of Chemical Technology and Biotechnology 82: 233-247.

Pieper-Fürst, U., M. H. Madkour, F. Mayer and A. Steinbüchel (1994). "Purification and characterization of a 14-kilodalton protein that is bound to the surface of polyhydroxyalkanoic acid granules in *Rhodococcus ruber*." Journal of Bacteriology 176: 4328-4337.

Pötter, M., M. H. Madkour, F. Mayer and A. Steinbüchel (2002). "Regulation of phasin expression and polyhydroxyalkanoate (PHA) granule formation in *Ralstonia eutropha* H16." Microbiol. 148: 2413-2426.

Pretsch, E., J. Seibl and W. Simon (1998). "Tablas para la determinación structural por métodos espectroscópicos." Springer -Verlag Ibérica.

Preusting, H., W. Hazenberg and B. Witholt (1993). "Continuous production of poly(3-hydroxyalkanoates by *Pseudomonas oleovorans* in a high-cell-density two-liquid-phase chemostat." Enzyme microbial Technology 15(April): 311-316.

Preusting, H., Nijenhuis,A., and Witholt, B. (1990). "Physical Characteristics of Poly(3-hydroxyalkanoates) and Poly(3-hydroxyalkanoates) produced by *Pseudomonas oleovorans* grown on aliphatic hydrocarbons." Macromolecules 23: 4220-4224.

Prieto, M. A., B. Bühler, K. Jung, B. Witholt and B. Kessler (1999). "PhaF, a polyhydroxyalkanoate-granule-associated protein od *Pseudomonas oleovorans* Gpo1 involved in the regulatory expression system for pha Genes." Journal of Bacteriology 181(3): 858-868.

Reddy, C. S. K., R. Ghai and K. V. C. Rashmi (2003). "Polyhydroxyalkanoates: an overview." Bioresource Technology 87: 137-146.

Rehm, B., N. Krüger and A. Steinbüchel (1998). "A new metabolic link between fatty acid *de novo* synthesis and polyhydroxyalkanoic acid synthesis." The Journal of Biological Chemistry 273, No.37(11): 24044-24051.

Rehm, B. H. A. (2006). "Genetics and biochemistry of polyhydroxyalkanoate granule self-assembly: The key role of polyester synthases." Biotechnology Letters 28: 207-213.

Rehm, B. H. A. and A. Steinbüchel (1999). "Biochemical and genetic analysis of PHA synthases and other proteins required for PHA synthesis." International Journal of Biological Macromolecules 25: 3-19.

Ren, Q., A. Grubelnik, M. Hoerler, K. Ruth, R. Hartmann, H. Felber and M. Zinn (2005). "Bacterial Poly(hydroxyalkanoates) as a Source of Chiral Hydroxyalkanoic Acids." Biomacromolecules 6(4): 2290-2298.

- Renard, E., Walls, M., Guerin, P and Langlois V. (2004). "Hydrolytic degradation of blends of polyhydroxyalkanoates and functionalized polyhydroxyalkanoates." Polymer Degradation and Stability 85: 779-787.
- Reynolds, E. S. (1963). "The use of lead citrate at high pH as an electron opaque strain in electron microscopy." J. Cel. Bio. 17: 208-212.
- Robert, M. (1989). "Aislamiento y selección de microorganismos productores de tensioactivos. Producción y caracterización de biotensioactivos por *Pseudomonas 44T1*." Departamento de Microbiología Facultad de Farmacia. Barcelona, Barcelona.
- Rodríguez, E. (2006). "Aplicación de la Metodología de Superficies de Respuesta den matraces y estrategias de producción en bioreactor para la obtención de biomasa y polihidroxialcanoatos por *Pseudomonas aeruginosa 42A2*." Tesis Doctoral. Universitat de Barcelona.
- Rossini, C., J. Arceo, E. McCarney, B. Augustine, D. Dennis, M. D. Flythe and S. F. baron (2001). "Use of in situ atomic force microscopy to monitor the biodegradation of polyhydroxyalkanoates (PHAs)." Macromol Symp 167: 139-151.
- Ruth, K., A. Grubelnik, R. Hartmann, T. Egli, M. Zinn and Q. Ren (2007). "Efficient production of (R)-3-hydroxycarboxylic acids by biotechnological conversion of Polyhydroxyalkanoates and their purification." Biomacromolecules 8(1): 279-286.
- Saad, B., P. Neuenschwander, G. K. Uhlsmitd and U. W. Suter (1999). "New versatile, elastomeric, degradable polymeric materials for medicine." International Journal of Biological Macromolecules 25: 293-301.
- Saegusa, H., K. M. Shiraki, C. Kanai and T. Saito (2001). "Cloning of an intracellular poly[D8-)3-hydroxybutyrate] depolymerase gene from *Ralstonia eutropha* H16 and characterization of the gene product." Journal of Bacteriology 183: 94-100.
- Schmid, M., A. Ritter, A. Grubelnik and M. Zinn (2006). "Autoxidation of Medium Chain Length Polyhydroxyalkanoate." Biomacromolecules 8(2): 579-584.
- Solaiman, D. K. Y., R. D. Ashby and T. A. Foglia (1999). "Medium-Chain-Length poly(β -hydroxyalkanoate) synthesis from triacylglycerols by *Pseudomonas saccharophila*." Current Microbiology 38: 151-154.
- Song, J. J., S. C. Yoon, S. M. Yu and R. W. Lenz (1998). "Differential scanning calorimetric study of poly(3-hydroxyoctanoate) inclusions in bacterial cells." International Journal of Biological Macromolecules 23: 165-173.
- Spurr, A. R. (1969). "A low-viscosity epoxy resin embedding medium for electron microscopy." J. Ultrastruct. Res. 26: 31.
- Steinbüchel, A. (2001). "Perspectives for Biotechnological Production and Utilization of Biopolymer: Metabolic Engineering of Polyhydroxyalkanoates Biosynthesis Pathways as a successful example." Macromol Biosci. 1: 1-24.
- Steinbüchel, A., K. Aerts, W. Babek, C. Föller, M. Liebergesell, M. H. Madkour, A. Pries, F. E. Valentin and R. Wieczorek (1995). "Considerations on the structure of

bacterial polyhydroxyalkanoic acid inclusions." Canadian J. Microbiol (Suppl-1) 41(1): 94-105.

Steinbüchel, A. and B. Füchtenbusch (1998). "Bacterial and other biological systems for polyester production." Trends in Biotechnology (TIBTECH) 16: 419-427.

Steinbüchel, A. and P. Schubert (1989). "Expression of the *Alcaligenes eutrophus* poly(β -hydroxybutyric acid)- synthetic pathway in *Pseudomonas sp.*" Archives of Microbiology 153: 101-104.

Steinbüchel, A. and H. E. Valentin (1995). "Diversity of bacterial polyhydroxyalkanoic acids." FEMS Microbiology Letters 128: 219-228.

Steinbüchel, A. and S. Wiese (1992). "A *Pseudomonas* strain accumulating Polyesters of 3-hydroxybutyric acid and Medium-Chain-Length 3-hydroxyalkanoic acids." Applied Microbiology and Biotechnology 37: 691-697.

Stuart, E. S., R. W. Lenz and R. C. Fuller (1995). "The ordered macromolecular surface of polyester inclusion bodies in *Pseudomonas oleovorans*." Canadian J. Microbiol 41(suppl.1): 84-93.

Stuart, E. S., A. Tehrani, H. E. Valentin, D. Dennis, R. W. Lenz and R. C. Fuller (1998). "Protein organization on the PHA inclusion cytoplasmic boundary." J. Biotechnol (2-3): 137-144.

Sudesh, K., H. Abe and Y. Doi (2000). "Synthesis, structure and properties of polyhydroxyalkanoates: biological polyesters." Progress in Polymer Science 25: 1503-1555.

Sudesh, K. and Y. Doi (2000). "Molecular Design and Biosynthesis of Biodegradable polyesters." Polymers for advanced technologies 11: 865-872.

Sudesh, K., Gan, Z., Maehara, A., Doi, Y. (2002). "Surface structure, morphology and stability of polyhydroxyalkanoate inclusions characterised by atomic force microscopy." Polymer Degradation and Stability 77: 77-85.

Takagi, Y., H. Hashii, A. Maehara and T. Yamane (1999). "Biosynthesis of Polyhydroxyalkanoate with a thiophenoxy Side Group Obtained from *Pseudomonas putida*." Macromolecules 32: 8315-8318.

Tanaka, T., M. Fujita, et al. (2005). "Structure investigation of narrow banded spherulites in polyhydroxyalkanoates by microbeam X-ray diffraction with synchrotron radiation." Polymer 46: 5673-5679.

Tian, W., K. Hong, G. Q. Chen, Q. Wu, R. Zhang and W. Huang (2000). "Production of polyesters consisting of medium chain length 3-hydroxyalkanoic acids by *Pseudomonas mendocina* 0806 from various carbon sources." Antonie van Leeuwenhoek 77: 31-36.

- Timm, A., S. Wiese and A. Steinbüchel (1994). "A general method for identification of polyhydroxyalkanoic acid synthase genes from *pseudomonads* belonging to the rRNA homolgy group I." Applied Microbiology and Biotechnology 40: 669-675.
- Tokiwa, Y. and T. Suzuki (1977). "Hydrolisis of polyesters by lipases." Nature 270: 76-78.
- Ubbink, J. and P. Schär-Zammaretti (2005). "Probing bacterial interactions: integrated approches combining atomic force microscopy, electron microscopy and biophysical techniques," Micron 36: 293-320.
- Valentin, H. E., Y. E. Lee, C. Y. Choi and A. Steinbüchel (1994). "Identification of 4-hydroxyhexanoic acid as a new constituent of biosynthetic polyhydroxyalkanoic acids from bacteria." Applied Microbiology and Biotechnology 40: 710-716.
- Valentin, H. E., S. Reiser and K. J. Gruys (2000). "Poly(3-hydroxybutyrate-co-4-hydroxybutyrate) formation from γ -Aminobutyrate and glutamate." Biotechnology and Bioengineering 67(3): 291-299.
- Valentin, H. E., A. Schönebaum and A. Steinbüchel (1996). "Identificaction of 5-hydroxyhexanoic acid, 4-hydroxyheptanoic acid and 4-hydroxyoctanoic acid as new constituents of bacterial polyhydroxyalkaonic acids." Applied Microbiology and Biotechnology 46: 261-267.
- van der Walle, G. A., G. J. de Koning, G. Weusthuis and G. Eggink (2001). "Properties, modifications and applications of biopolymers." Adv. Biochem. Eng. Biotechnol. 71: 263-291.
- Van der Walle, G. A. M., G. J. H. Buisman, R. A. Weusthuis and G. Eggink (1999). "Development of environmentally friendly coatings and paints using medium-chain-length poly(3-hydroxyalkanoates) as the polymer binder." International Journal of Biological Macromolecules 25: 123-128.
- Wallen, L. L. and W. K. Rohwedder (1974). "Poly- β -hydroxyalkanoates from activated sludge." Environ. Sci. Technol. 8: 576-579.
- Wang, J. G. and L. R. Bakken (1998). "Screening of soil bacteia for polybeta-hydroxybutyric acid production and its role in the survival of starvation." Microb. Ecol. 35: 94-101.
- Williams, S. F., D. P. Martin, D. M. Horowitz and O. P. Peoples (1999). "PHA applications: addressing the price performance issue I. Tissue engineering." International Journal of Biological Macromolecules 25: 111-121.
- Witholt, B. and B. Kessler (1999). "Perspectives of medium chain length poly(hydroxyalkanoates), a versatile set of bacterial bioplastics." Current Opinion in Biotechnology 10: 279-285.
- Xi, J., Q. Wu, Y. Yan and G.-Q. Chen (2000). "Hyperproduction of polyesters consisting of medium-chain-lenght hydroxyalkanoate monomers by strain *Pseudomonas stutzeri* 1317." Antonie van Leeuwenhoek 78: 43-49.

Xu, J., B. H. Guo, et al. (2003). "Terraces on banded spherulites of polyhydroxyalkanoates." Journal of Polymer Science: Part B: Polymer Physics 41: 2128-2134.

York, G. M., B. H. Junker, J. Stubbe and A. J. Sinskey (2001). "Accumulation of the PhaP phasin on *Ralstonia eutropha* is dependent on production of polyhydroxybutyrate in cells." Journal of Bacteriology 183(14): 4217-4226.

York, G. M., J. Stubbe and A. J. Sinskey (2002). "The *Ralstonia eutropha* PhaR protein couples synthesis of the PhaP phasin to the presence of polyhydroxybutyrate in cells and promotes polyhydroxybutyrate production." Journal of Bacteriology 184: 59-66.

Yu, P. H., H. Chua, A. L. Huang, W. Lo and G. Q. Chen (1998). "Conversion of food industrial wastes into bioplastics." Applied Biochemistry and Biotechnology 70-72: 603-614.

Zinn, M., B. Wiltholt and T. Egli (2001). "Occurrence, synthesis and medical application of bacterial polyhydroxyalkanoates." Adv. Drug Deliv. Rev. 53: 5-21.

Zinn, M., M., H.-U. Weilenmann, R. Hany, M. Schmid and T. Egli (2003). "Tailored synthesis of poly([R]-3-hydroxybutyrate-co-3-hydroxyvalerate)(PHB/HV) in *Ralstonia eutropha* DSM 428." Acta Biotechnologica 23(2-3): 309-316.