



Caracterización funcional de la interacción de la proteína p53 con la ubiquitina ligasa HERC2

Mónica Cubillos Rojas

ADVERTIMENT. La consulta d'aquesta tesi queda condicionada a l'acceptació de les següents condicions d'ús: La difusió d'aquesta tesi per mitjà del servei TDX (www.tdx.cat) i a través del Dipòsit Digital de la UB (diposit.ub.edu) ha estat autoritzada pels titulars dels drets de propietat intel·lectual únicament per a usos privats emmarcats en activitats d'investigació i docència. No s'autoritza la seva reproducció amb finalitats de lucre ni la seva difusió i posada a disposició des d'un lloc aliè al servei TDX ni al Dipòsit Digital de la UB. No s'autoritza la presentació del seu contingut en una finestra o marc aliè a TDX o al Dipòsit Digital de la UB (framing). Aquesta reserva de drets afecta tant al resum de presentació de la tesi com als seus continguts. En la utilització o cita de parts de la tesi és obligat indicar el nom de la persona autora.

ADVERTENCIA. La consulta de esta tesis queda condicionada a la aceptación de las siguientes condiciones de uso: La difusión de esta tesis por medio del servicio TDR (www.tdx.cat) y a través del Repositorio Digital de la UB (diposit.ub.edu) ha sido autorizada por los titulares de los derechos de propiedad intelectual únicamente para usos privados enmarcados en actividades de investigación y docencia. No se autoriza su reproducción con finalidades de lucro ni su difusión y puesta a disposición desde un sitio ajeno al servicio TDR o al Repositorio Digital de la UB. No se autoriza la presentación de su contenido en una ventana o marco ajeno a TDR o al Repositorio Digital de la UB (framing). Esta reserva de derechos afecta tanto al resumen de presentación de la tesis como a sus contenidos. En la utilización o cita de partes de la tesis es obligado indicar el nombre de la persona autora.

WARNING. On having consulted this thesis you're accepting the following use conditions: Spreading this thesis by the TDX (www.tdx.cat) service and by the UB Digital Repository (diposit.ub.edu) has been authorized by the titular of the intellectual property rights only for private uses placed in investigation and teaching activities. Reproduction with lucrative aims is not authorized nor its spreading and availability from a site foreign to the TDX service or to the UB Digital Repository. Introducing its content in a window or frame foreign to the TDX service or to the UB Digital Repository is not authorized (framing). Those rights affect to the presentation summary of the thesis as well as to its contents. In the using or citation of parts of the thesis it's obliged to indicate the name of the author.



**Caracterización funcional
de la interacción de la proteína p53
con la ubiquitina ligasa HERC2**

Mónica Cubillos Rojas

Tesis Doctoral

2014

Bibliografía

- Abida, W. M., Nikolaev, A., Zhao, W., Zhang, W. and Gu, W. (2007). FBXO11 promotes the Neddylation of p53 and inhibits its transcriptional activity. *J. Biol. Chem.* 282, 1797–804.
- Al-Hakim, A. K., Bashkurov, M., Gingras, A.-C., Durocher, D. and Pelletier, L. (2012). Interaction proteomics identify NEURL4 and the HECT E3 ligase HERC2 as novel modulators of centrosome architecture. *Mol. Cell. Proteomics* 11, M111.014233.
- Ali, M. H. and Imperiali, B. (2005). Protein oligomerization: how and why. *Bioorg. Med. Chem.* 13, 5013–20.
- Amit, I., Citri, A., Shay, T., Lu, Y., Katz, M., Zhang, F., Tarcic, G., Siwak, D., Lahad, J., Jacob-Hirsch, J., et al. (2007). A module of negative feedback regulators defines growth factor signaling. *Nat. Genet.* 39, 503–12.
- Andrews, P., He, Y. J. and Xiong, Y. (2006). Cytoplasmic localized ubiquitin ligase cullin 7 binds to p53 and promotes cell growth by antagonizing p53 function. *Oncogene* 25, 4534–48.
- Attardi, L. D., Reczek, E. E., Cosmas, C., Demicco, E. G., McCurrach, M. E., Lowe, S. W. and Jacks, T. (2000). PERP, an apoptosis-associated target of p53, is a novel member of the PMP-22/gas3 family. *Genes Dev.* 14, 704–18.
- Badie, C., Bourhis, J., Sobczak-Thépot, J., Haddada, H., Chiron, M., Janicot, M., Janot, F., Tursz, T. and Vassal, G. (2000). p53-dependent G2 arrest associated with a decrease in cyclins A2 and B1 levels in a human carcinoma cell line. *Br. J. Cancer* 82, 642–50.
- Baeriswyl, V. and Christofori, G. (2009). The angiogenic switch in carcinogenesis. *Semin. Cancer Biol.* 19, 329–37.
- Barboza, J. A., Liu, G., Ju, Z., El-Naggar, A. K. and Lozano, G. (2006). p21 delays tumor onset by preservation of chromosomal stability. *Proc. Natl. Acad. Sci. U. S. A.* 103, 19842–7.
- Bassing, C. H., Suh, H., Ferguson, D. O., Chua, K. F., Manis, J., Eckersdorff, M., Gleason, M., Bronson, R., Lee, C. and Alt, F. W. (2003). Histone H2AX: a dosage-dependent suppressor of oncogenic translocations and tumors. *Cell* 114, 359–70.
- Bech-Otschir, D., Kraft, R., Huang, X., Henklein, P., Kapelari, B., Pollmann, C. and Dubiel, W. (2001). COP9 signalosome-specific phosphorylation targets p53 to degradation by the ubiquitin system. *EMBO J.* 20, 1630–9.
- Bekker-Jensen, S. and Mailand, N. (2010). Assembly and function of DNA double-strand break repair foci in mammalian cells. *DNA Repair (Amst)*. 9, 1219–28.
- Bekker-Jensen, S., Rendtlew Danielsen, J., Fugger, K., Gromova, I., Nerstedt, A., Lukas, C., Bartek, J., Lukas, J. and Mailand, N. (2010). HERC2 coordinates ubiquitin-dependent assembly of DNA repair factors on damaged chromosomes. *Nat. Cell Biol.* 12, 80–6; sup pp 1–12.
- Ben-David, Y., Lavigne, A., Cheong, G. Y. and Bernstein, A. (1990). Insertional inactivation of the p53 gene during friend leukemia: a new strategy for identifying tumor suppressor genes. *New Biol.* 2, 1015–23.
- Bennett, M., Macdonald, K., Chan, S. W., Luzio, J. P., Simari, R. and Weissberg, P. (1998). Cell surface trafficking of Fas: a rapid mechanism of p53-mediated apoptosis. *Science* 282, 290–3.
- Bernassola, F., Karin, M., Ciechanover, A. and Melino, G. (2008). The HECT family of E3 ubiquitin ligases: multiple players in cancer development. *Cancer Cell* 14, 10–21.
- Bibeau-Poirier, A., Gravel, S.-P., Clément, J.-F., Rolland, S., Rodier, G., Coulombe, P., Hiscott, J., Grandvaux, N., Meloche, S. and Servant, M. J. (2006). Involvement of the I κ B kinase (IKK)-related kinases tank-binding kinase 1/IKKi and cullin-based ubiquitin ligases in IFN regulatory factor-3 degradation. *J. Immunol.* 177, 5059–67.

Bibliografia

- Bischof, O. and Dejean, A. (2007). SUMO is growing senescent. *Cell Cycle* 6, 677–81.
- Blasco, M. A. (2005). Telomeres and human disease: ageing, cancer and beyond. *Nat. Rev. Genet.* 6, 611–22.
- Bogunovic, D., Boisson-Dupuis, S. and Casanova, J.-L. (2013). ISG15: leading a double life as a secreted molecule. *Exp. Mol. Med.* 45, e18.
- Bonner, W. M., Redon, C. E., Dickey, J. S., Nakamura, A. J., Sedelnikova, O. A., Solier, S. and Pommier, Y. (2008). GammaH2AX and cancer. *Nat. Rev. Cancer* 8, 957–67.
- Bourdon, J.-C. (2007). p53 and its isoforms in cancer. *Br. J. Cancer* 97, 277–82.
- Bouvard, V., Zaitchouk, T., Vacher, M., Duthu, A., Canivet, M., Choisy-Rossi, C., Nieruchalski, M. and May, E. (2000). Tissue and cell-specific expression of the p53-target genes: bax, fas, mdm2 and waf1/p21, before and following ionising irradiation in mice. *Oncogene* 19, 649–60.
- Bramall, A. N., Wright, A. F., Jacobson, S. G. and McInnes, R. R. (2010). The genomic, biochemical, and cellular responses of the retina in inherited photoreceptor degenerations and prospects for the treatment of these disorders. *Annu. Rev. Neurosci.* 33, 441–72.
- Branicki, W., Brudnik, U. and Wojas-Pelc, A. (2009). Interactions between HERC2, OCA2 and MC1R may influence human pigmentation phenotype. *Ann. Hum. Genet.* 73, 160–70.
- Brehm, A., Miska, E. A., McCance, D. J., Reid, J. L., Bannister, A. J. and Kouzarides, T. (1998). Retinoblastoma protein recruits histone deacetylase to repress transcription. *Nature* 391, 597–601.
- Brooks, C. L. and Gu, W. (2003). Ubiquitination, phosphorylation and acetylation: the molecular basis for p53 regulation. *Curr. Opin. Cell Biol.* 15, 164–71.
- Brooks, C. L., Li, M., Hu, M., Shi, Y. and Gu, W. (2007). The p53--Mdm2--HAUSP complex is involved in p53 stabilization by HAUSP. *Oncogene* 26, 7262–6.
- Brugarolas, J., Bronson, R. T. and Jacks, T. (1998). p21 is a critical CDK2 regulator essential for proliferation control in Rb-deficient cells. *J. Cell Biol.* 141, 503–14.
- Budanov, A. V. (2011). Stress-responsive sestrins link p53 with redox regulation and mammalian target of rapamycin signaling. *Antioxid. Redox Signal.* 15, 1679–90.
- Budhram-Mahadeo, V., Morris, P. J., Smith, M. D., Midgley, C. A., Boxer, L. M. and Latchman, D. S. (1999). p53 suppresses the activation of the Bcl-2 promoter by the Brn-3a POU family transcription factor. *J. Biol. Chem.* 274, 15237–44.
- Burkhardt, D. L. and Sage, J. (2008). Cellular mechanisms of tumour suppression by the retinoblastoma gene. *Nat. Rev. Cancer* 8, 671–82.
- Cai, H., Shim, H., Lai, C., Xie, C., Lin, X., Yang, W. J. and Chandran, J. (2008). ALS2/alsin knockout mice and motor neuron diseases. *Neurodegener. Dis.* 5, 359–66.
- Campana, A. L. M., Rondi-Reig, L., Tobin, C., Lohof, A. M., Picquet, F., Falempin, M., Weitzman, J. B. and Mariani, J. (2003). p53 inactivation leads to impaired motor synchronization in mice. *Eur. J. Neurosci.* 17, 2135–46.

- Carazo-Salas, R. E., Guarguaglini, G., Gruss, O. J., Segref, A., Karsenti, E. and Mattaj, I. W. (1999). Generation of GTP-bound Ran by RCC1 is required for chromatin-induced mitotic spindle formation. *Nature* 400, 178–81.
- Carter, S., Bischof, O., Dejean, A. and Vousden, K. H. (2007). C-terminal modifications regulate MDM2 dissociation and nuclear export of p53. *Nat. Cell Biol.* 9, 428–35.
- Casas-Terradellas, E., Garcia-Gonzalo, F. R., Hadjebi, O., Bartrons, R., Ventura, F. and Rosa, J. L. (2006). Simultaneous electrophoretic analysis of proteins of very high and low molecular weights using low-percentage acrylamide gel and a gradient SDS-PAGE gel. *Electrophoresis* 27, 3935–8.
- Celeste, A., Petersen, S., Romanienko, P. J., Fernandez-Capetillo, O., Chen, H. T., Sedelnikova, O. A., Reina-San-Martin, B., Coppola, V., Meffre, E., Difilippantonio, M. J., et al. (2002). Genomic instability in mice lacking histone H2AX. *Science* 296, 922–7.
- Chen, J. and Sadowski, I. (2005). Identification of the mismatch repair genes PMS2 and MLH1 as p53 target genes by using serial analysis of binding elements. *Proc. Natl. Acad. Sci. U. S. A.* 102, 4813–8.
- Chen, J., Saha, P., Kornbluth, S., Dynlacht, B. D. and Dutta, A. (1996). Cyclin-binding motifs are essential for the function of p21CIP1. *Mol. Cell. Biol.* 16, 4673–82.
- Chen, D., Kon, N., Li, M., Zhang, W., Qin, J. and Gu, W. (2005). ARF-BP1/Mule is a critical mediator of the ARF tumor suppressor. *Cell* 121, 1071–83.
- Chène, P. (2001). The role of tetramerization in p53 function. *Oncogene* 20, 2611–7.
- Cheng, Q. and Chen, J. (2010). Mechanism of p53 stabilization by ATM after DNA damage. *Cell Cycle* 9, 472–8.
- Cheng, N., Chytil, A., Shyr, Y., Joly, A. and Moses, H. L. (2008). Transforming growth factor-beta signaling-deficient fibroblasts enhance hepatocyte growth factor signaling in mammary carcinoma cells to promote scattering and invasion. *Mol. Cancer Res.* 6, 1521–33.
- Chipuk, J. E., Bouchier-Hayes, L., Kuwana, T., Newmeyer, D. D. and Green, D. R. (2005). PUMA couples the nuclear and cytoplasmic proapoptotic function of p53. *Science* 309, 1732–5.
- Chong-Kopera, H., Inoki, K., Li, Y., Zhu, T., Garcia-Gonzalo, F. R., Rosa, J. L. and Guan, K.-L. (2006). TSC1 stabilizes TSC2 by inhibiting the interaction between TSC2 and the HERC1 ubiquitin ligase. *J. Biol. Chem.* 281, 8313–6.
- Choudhury, A. R., Ju, Z., Djojotubroto, M. W., Schienke, A., Lechel, A., Schaetzlein, S., Jiang, H., Stepczynska, A., Wang, C., Buer, J., et al. (2007). Cdkn1a deletion improves stem cell function and lifespan of mice with dysfunctional telomeres without accelerating cancer formation. *Nat. Genet.* 39, 99–105.
- Chuikov, S., Kurash, J. K., Wilson, J. R., Xiao, B., Justin, N., Ivanov, G. S., McKinney, K., Tempst, P., Prives, C., Gamblin, S. J., et al. (2004). Regulation of p53 activity through lysine methylation. *Nature* 432, 353–60.
- Ciechanover, A. (1998). The ubiquitin-proteasome pathway: on protein death and cell life. *EMBO J.* 17, 7151–60.
- Ciechanover, A., Elias, S., Heller, H., Ferber, S. and Hershko, A. (1980). Characterization of the heat-stable polypeptide of the ATP-dependent proteolytic system from reticulocytes. *J. Biol. Chem.* 255, 7525–8.
- Clarke, P. R. and Zhang, C. (2001). Ran GTPase: a master regulator of nuclear structure and function during the eukaryotic cell division cycle? *Trends Cell Biol.* 11, 366–71.

Bibliografia

- Clarke, P. R. and Zhang, C. (2008). Spatial and temporal coordination of mitosis by Ran GTPase. *Nat. Rev. Mol. Cell Biol.* 9, 464–77.
- Clore, G. M., Omichinski, J. G., Sakaguchi, K., Zambrano, N., Sakamoto, H., Appella, E. and Gronenborn, A. M. (1994). High-resolution structure of the oligomerization domain of p53 by multidimensional NMR. *Science* 265, 386–91.
- Colman, M. S., Afshari, C. A. and Barrett, J. C. (2000). Regulation of p53 stability and activity in response to genotoxic stress. *Mutat. Res.* 462, 179–88.
- Contente, A., Dittmer, A., Koch, M. C., Roth, J. and Dobbelstein, M. (2002). A polymorphic microsatellite that mediates induction of PIG3 by p53. *Nat. Genet.* 30, 315–20.
- Cory, S., Huang, D. C. S. and Adams, J. M. (2003). The Bcl-2 family: roles in cell survival and oncogenesis. *Oncogene* 22, 8590–607.
- Cruz, C., Nadal, M., Ventura, F., Bartrons, R., Estivill, X. and Rosa, J. L. (1999). The human HERC3 gene maps to chromosome 4q21 by fluorescence in situ hybridization. *Cytogenet. Cell Genet.* 87, 263–4.
- Cruz, C., Ventura, F., Bartrons, R. and Rosa, J. L. (2001). HERC3 binding to and regulation by ubiquitin. *FEBS Lett.* 488, 74–80.
- Cubillos-Rojas, M., Amair-Pinedo, F., Tato, I., Bartrons, R., Ventura, F. and Rosa, J. L. (2010). Simultaneous electrophoretic analysis of proteins of very high and low molecular mass using Tris-acetate polyacrylamide gels. *Electrophoresis* 31, 1318–21.
- Cubillos-Rojas, M., Amair-Pinedo, F., Tato, I., Bartrons, R., Ventura, F. and Rosa, J. L. (2012). Tris-acetate polyacrylamide gradient gels for the simultaneous electrophoretic analysis of proteins of very high and low molecular mass. *Methods Mol. Biol.* 869, 205–13.
- Cubillos-Rojas, M., Amair-Pinedo, F., Peiro-Jordan, R., Bartrons, R., Ventura, F. and Rosa, J. L. (2014). The E3 Ubiquitin Protein Ligase HERC2 Modulates Activity of Tumor Protein p53 by Regulating its Oligomerization. *J. Biol. Chem.*
- Cukierman, E., Huber, I., Rotman, M. and Cassel, D. (1995). The ARF1 GTPase-activating protein: zinc finger motif and Golgi complex localization. *Science* 270, 1999–2002.
- D'Orazi, G., Cecchinelli, B., Bruno, T., Manni, I., Higashimoto, Y., Saito, S., Gostissa, M., Coen, S., Marchetti, A., Del Sal, G., et al. (2002). Homeodomain-interacting protein kinase-2 phosphorylates p53 at Ser 46 and mediates apoptosis. *Nat. Cell Biol.* 4, 11–9.
- Danielsen, J. R., Povlsen, L. K., Villumsen, B. H., Streicher, W., Nilsson, J., Wikström, M., Bekker-Jensen, S. and Mailand, N. (2012). DNA damage-inducible SUMOylation of HERC2 promotes RNF8 binding via a novel SUMO-binding Zinc finger. *J. Cell Biol.* 197, 179–87.
- Darchen, F., Senyshyn, J., Brondyk, W. H., Taatjes, D. J., Holz, R. W., Henry, J. P., Denizot, J. P. and Macara, I. G. (1995). The GTPase Rab3a is associated with large dense core vesicles in bovine chromaffin cells and rat PC12 cells. *J. Cell Sci.* 108 (Pt 4, 1639–49.
- Dastur, A., Beaudenon, S., Kelley, M., Krug, R. M. and Huibregtse, J. M. (2006). Herc5, an interferon-induced HECT E3 enzyme, is required for conjugation of ISG15 in human cells. *J. Biol. Chem.* 281, 4334–8.
- Davies, W., Smith, R. J., Kelsey, G. and Wilkinson, L. S. (2004). Expression patterns of the novel imprinted genes Nap115 and Peg13 and their non-imprinted host genes in the adult mouse brain. *Gene Expr. Patterns* 4, 741–7.

- Davison, T. S., Yin, P., Nie, E., Kay, C. and Arrowsmith, C. H. (1998). Characterization of the oligomerization defects of two p53 mutants found in families with Li-Fraumeni and Li-Fraumeni-like syndrome. *Oncogene* 17, 651–6.
- Diouf, B., Cheng, Q., Krynetskaia, N. F., Yang, W., Cheok, M., Pei, D., Fan, Y., Cheng, C., Krynetskiy, E. Y., Geng, H., et al. (2011). Somatic deletions of genes regulating MSH2 protein stability cause DNA mismatch repair deficiency and drug resistance in human leukemia cells. *Nat. Med.* 17, 1298–303.
- Donehower, L. A. and Lozano, G. (2009). 20 years studying p53 functions in genetically engineered mice. *Nat. Rev. Cancer* 9, 831–41.
- Donehower, L. A., Harvey, M., Slagle, B. L., McArthur, M. J., Montgomery, C. A., Butel, J. S. and Bradley, A. (1992). Mice deficient for p53 are developmentally normal but susceptible to spontaneous tumours. *Nature* 356, 215–21.
- Dornan, D., Wertz, I., Shimizu, H., Arnott, D., Frantz, G. D., Dowd, P., O'Rourke, K., Koeppen, H. and Dixit, V. M. (2004). The ubiquitin ligase COP1 is a critical negative regulator of p53. *Nature* 429, 86–92.
- Du, H., Wolf, J., Schafer, B., Moldoveanu, T., Chipuk, J. E. and Kuwana, T. (2011). BH3 domains other than Bim and Bid can directly activate Bax/Bak. *J. Biol. Chem.* 286, 491–501.
- Durfee, L. A., Lyon, N., Seo, K. and Huijbregtse, J. M. (2010). The ISG15 conjugation system broadly targets newly synthesized proteins: implications for the antiviral function of ISG15. *Mol. Cell* 38, 722–32.
- Eiberg, H., Troelsen, J., Nielsen, M., Mikkelsen, A., Mengel-From, J., Kjaer, K. W. and Hansen, L. (2008). Blue eye color in humans may be caused by a perfectly associated founder mutation in a regulatory element located within the HERC2 gene inhibiting OCA2 expression. *Hum. Genet.* 123, 177–87.
- el-Deiry, W. S. (1998). Regulation of p53 downstream genes. *Semin. Cancer Biol.* 8, 345–57.
- el-Deiry, W. S., Tokino, T., Velculescu, V. E., Levy, D. B., Parsons, R., Trent, J. M., Lin, D., Mercer, W. E., Kinzler, K. W. and Vogelstein, B. (1993). WAF1, a potential mediator of p53 tumor suppression. *Cell* 75, 817–25.
- Enari, M., Ohmori, K., Kitabayashi, I. and Taya, Y. (2006). Requirement of clathrin heavy chain for p53-mediated transcription. *Genes Dev.* 20, 1087–99.
- Engert, A., Franklin, J., Eich, H. T., Brillant, C., Sehlen, S., Cartoni, C., Herrmann, R., Pfreundschuh, M., Sieber, M., Tesch, H., et al. (2007). Two cycles of doxorubicin, bleomycin, vinblastine, and dacarbazine plus extended-field radiotherapy is superior to radiotherapy alone in early favorable Hodgkin's lymphoma: final results of the GHSG HD7 trial. *J. Clin. Oncol.* 25, 3495–502.
- Faraonio, R., Vergara, P., Di Marzo, D., Pierantoni, M. G., Napolitano, M., Russo, T. and Cimino, F. (2006). p53 suppresses the Nrf2-dependent transcription of antioxidant response genes. *J. Biol. Chem.* 281, 39776–84.
- Fernandez-Fernandez, M. R., Veprintsev, D. B. and Fersht, A. R. (2005). Proteins of the S100 family regulate the oligomerization of p53 tumor suppressor. *Proc. Natl. Acad. Sci. U. S. A.* 102, 4735–40.
- Fidler, I. J., Kim, S.-J. and Langley, R. R. (2007). The role of the organ microenvironment in the biology and therapy of cancer metastasis. *J. Cell. Biochem.* 101, 927–36.
- Fogal, V., Hsieh, J.-K., Royer, C., Zhong, S. and Lu, X. (2005). Cell cycle-dependent nuclear retention of p53 by E2F1 requires phosphorylation of p53 at Ser315. *EMBO J.* 24, 2768–82.

Bibliografía

- Foo, R. S.-Y., Nam, Y.-J., Ostreicher, M. J., Metz, M. D., Whelan, R. S., Peng, C.-F., Ashton, A. W., Fu, W., Mani, K., Chin, S.-F., et al. (2007). Regulation of p53 tetramerization and nuclear export by ARC. *Proc. Natl. Acad. Sci. U. S. A.* 104, 20826–31.
- Franco, S., Gostissa, M., Zha, S., Lombard, D. B., Murphy, M. M., Zarrin, A. A., Yan, C., Tepsuporn, S., Morales, J. C., Adams, M. M., et al. (2006). H2AX prevents DNA breaks from progressing to chromosome breaks and translocations. *Mol. Cell* 21, 201–14.
- Franken, N. A. P., Rodermond, H. M., Stap, J., Haveman, J. and van Bree, C. (2006). Clonogenic assay of cells in vitro. *Nat. Protoc.* 1, 2315–9.
- Franklin, D. S., Godfrey, V. L., O'Brien, D. A., Deng, C. and Xiong, Y. (2000). Functional collaboration between different cyclin-dependent kinase inhibitors suppresses tumor growth with distinct tissue specificity. *Mol. Cell. Biol.* 20, 6147–58.
- Frudakis, T., Thomas, M., Gaskin, Z., Venkateswarlu, K., Chandra, K. S., Ginjupalli, S., Gunturi, S., Natrajan, S., Ponnuswamy, V. K. and Ponnuswamy, K. N. (2003). Sequences associated with human iris pigmentation. *Genetics* 165, 2071–83.
- Gaglia, G., Guan, Y., Shah, J. V and Lahav, G. (2013). Activation and control of p53 tetramerization in individual living cells. *Proc. Natl. Acad. Sci. U. S. A.* 110, 15497–501.
- Gakovic, M., Shu, X., Kasioulis, I., Carpanini, S., Moraga, I. and Wright, A. F. (2011). The role of RPGR in cilia formation and actin stability. *Hum. Mol. Genet.* 20, 4840–50.
- Galluzzi, L. and Kroemer, G. (2008). Necroptosis: a specialized pathway of programmed necrosis. *Cell* 135, 1161–3.
- Garami, A., Zwartkruis, F. J. T., Nobukuni, T., Joaquin, M., Rocco, M., Stocker, H., Kozma, S. C., Hafen, E., Bos, J. L. and Thomas, G. (2003). Insulin activation of Rheb, a mediator of mTOR/S6K/4E-BP signaling, is inhibited by TSC1 and 2. *Mol. Cell* 11, 1457–66.
- Garbarini, N. and Delpire, E. (2008). The RCC1 domain of protein associated with Myc (PAM) interacts with and regulates KCC2. *Cell. Physiol. Biochem.* 22, 31–44.
- Garcia-Gonzalo, F. R. and Rosa, J. L. (2005). The HERC proteins: functional and evolutionary insights. *Cell. Mol. Life Sci.* 62, 1826–38.
- Garcia-Gonzalo, F. R., Cruz, C., Muñoz, P., Mazurek, S., Eigenbrodt, E., Ventura, F., Bartrons, R. and Rosa, J. L. (2003). Interaction between HERC1 and M2-type pyruvate kinase. *FEBS Lett.* 539, 78–84.
- Garcia-Gonzalo, F. R., Muñoz, P., González, E., Casaroli-Marano, R. P., Vilaró, S., Bartrons, R., Ventura, F. and Rosa, J. L. (2004). The giant protein HERC1 is recruited to aluminum fluoride-induced actin-rich surface protrusions in HeLa cells. *FEBS Lett.* 559, 77–83.
- Garcia-Gonzalo, F. R., Bartrons, R., Ventura, F. and Rosa, J. L. (2005). Requirement of phosphatidylinositol-4,5-bisphosphate for HERC1-mediated guanine nucleotide release from ARF proteins. *FEBS Lett.* 579, 343–8.
- Ghosh, M., Huang, K. and Berberich, S. J. (2003). Overexpression of Mdm2 and MdmX fusion proteins alters p53 mediated transactivation, ubiquitination, and degradation. *Biochemistry* 42, 2291–9.
- Giannakopoulos, N. V., Luo, J.-K., Papov, V., Zou, W., Lenschow, D. J., Jacobs, B. S., Borden, E. C., Li, J., Virgin, H. W. and Zhang, D.-E. (2005). Proteomic identification of proteins conjugated to ISG15 in mouse and human cells. *Biochem. Biophys. Res. Commun.* 336, 496–506.

- Green, D. R. and Kroemer, G. (2009). Cytoplasmic functions of the tumour suppressor p53. *Nature* 458, 1127–30.
- Grossberger, R., Gieffers, C., Zachariae, W., Podtelejnikov, A. V., Schleiffer, A., Nasmyth, K., Mann, M. and Peters, J. M. (1999). Characterization of the DOC1/APC10 subunit of the yeast and the human anaphase-promoting complex. *J. Biol. Chem.* 274, 14500–7.
- Gu, W. and Roeder, R. G. (1997). Activation of p53 sequence-specific DNA binding by acetylation of the p53 C-terminal domain. *Cell* 90, 595–606.
- Guo, Q., Xie, J., Dang, C. V., Liu, E. T. and Bishop, J. M. (1998). Identification of a large Myc-binding protein that contains RCC1-like repeats. *Proc. Natl. Acad. Sci. U. S. A.* 95, 9172–7.
- Hachmann, J. P. and Amshey, J. W. (2005). Models of protein modification in Tris-glycine and neutral pH Bis-Tris gels during electrophoresis: effect of gel pH. *Anal. Biochem.* 342, 237–45.
- Hadjebi, O., Casas-Terradellas, E., Garcia-Gonzalo, F. R. and Rosa, J. L. (2008). The RCC1 superfamily: from genes, to function, to disease. *Biochim. Biophys. Acta* 1783, 1467–79.
- Han, S., Kim, S., Bahl, S., Li, L., Burande, C. F., Smith, N., James, M., Beauchamp, R. L., Bhide, P., DiAntonio, A., et al. (2012). The E3 ubiquitin ligase protein associated with Myc (Pam) regulates mammalian/mechanistic target of rapamycin complex 1 (mTORC1) signaling in vivo through N- and C-terminal domains. *J. Biol. Chem.* 287, 30063–72.
- Hanahan, D. and Weinberg, R. A. (2011). Hallmarks of cancer: the next generation. *Cell* 144, 646–74.
- Harlalka, G. V., Baple, E. L., Cross, H., Kühnle, S., Cubillos-Rojas, M., Matentzoglou, K., Patton, M. A., Wagner, K., Coblentz, R., Ford, D. L., et al. (2013). Mutation of HERC2 causes developmental delay with Angelman-like features. *J. Med. Genet.* 50, 65–73.
- Harper, J. W., Adami, G. R., Wei, N., Keyomarsi, K. and Elledge, S. J. (1993). The p21 Cdk-interacting protein Cip1 is a potent inhibitor of G1 cyclin-dependent kinases. *Cell* 75, 805–16.
- Harper, J. W., Elledge, S. J., Keyomarsi, K., Dynlacht, B., Tsai, L. H., Zhang, P., Dobrowolski, S., Bai, C., Connell-Crowley, L. and Swindell, E. (1995). Inhibition of cyclin-dependent kinases by p21. *Mol. Biol. Cell* 6, 387–400.
- Hartlerode, A. J. and Scully, R. (2009). Mechanisms of double-strand break repair in somatic mammalian cells. *Biochem. J.* 423, 157–68.
- Haupt, Y., Maya, R., Kazaz, A. and Oren, M. (1997). Mdm2 promotes the rapid degradation of p53. *Nature* 387, 296–9.
- Haupt, S., Berger, M., Goldberg, Z. and Haupt, Y. (2003). Apoptosis - the p53 network. *J. Cell Sci.* 116, 4077–85.
- Helleday, T., Petermann, E., Lundin, C., Hodgson, B. and Sharma, R. A. (2008). DNA repair pathways as targets for cancer therapy. *Nat. Rev. Cancer* 8, 193–204.
- Hermeking, H. (2012). MicroRNAs in the p53 network: micromanagement of tumour suppression. *Nat. Rev. Cancer* 12, 613–26.
- Hermeking, H., Lengauer, C., Polyak, K., He, T. C., Zhang, L., Thiagalingam, S., Kinzler, K. W. and Vogelstein, B. (1997). 14-3-3 sigma is a p53-regulated inhibitor of G2/M progression. *Mol. Cell* 1, 3–11.

Bibliografía

- Hetzer, M., Bilbao-Cortés, D., Walther, T. C., Gruss, O. J. and Mattaj, I. W. (2000). GTP hydrolysis by Ran is required for nuclear envelope assembly. *Mol. Cell* 5, 1013–24.
- Hochrainer, K., Mayer, H., Baranyi, U., Binder, B., Lipp, J. and Kroismayr, R. (2005). The human HERC family of ubiquitin ligases: novel members, genomic organization, expression profiling, and evolutionary aspects. *Genomics* 85, 153–64.
- Hochrainer, K., Kroismayr, R., Baranyi, U., Binder, B. R. and Lipp, J. (2008). Highly homologous HERC proteins localize to endosomes and exhibit specific interactions with hPLIC and Nm23B. *Cell. Mol. Life Sci.* 65, 2105–17.
- Hochstrasser, M. (2000a). Biochemistry. All in the ubiquitin family. *Science* 289, 563–4.
- Hochstrasser, M. (2000b). Evolution and function of ubiquitin-like protein-conjugation systems. *Nat. Cell Biol.* 2, E153–7.
- Hock, A. and Vousden, K. H. (2010). Regulation of the p53 pathway by ubiquitin and related proteins. *Int. J. Biochem. Cell Biol.* 42, 1618–21.
- Hock, A. K. and Vousden, K. H. (2014). The role of ubiquitin modification in the regulation of p53. *Biochim. Biophys. Acta* 1843, 137–49.
- HOLLANDER, W. F., BRYAN, J. H. and GOWEN, J. W. A male sterile pink-eyed mutant type in the mouse. *Fertil. Steril.* 11, 316–24.
- Hollander, M. C., Sheikh, M. S., Bulavin, D. V., Lundgren, K., Augeri-Henmueller, L., Shehee, R., Molinaro, T. A., Kim, K. E., Tolosa, E., Ashwell, J. D., et al. (1999). Genomic instability in Gadd45a-deficient mice. *Nat. Genet.* 23, 176–84.
- Honda, R. and Yasuda, H. (1999). Association of p19(ARF) with Mdm2 inhibits ubiquitin ligase activity of Mdm2 for tumor suppressor p53. *EMBO J.* 18, 22–7.
- Honda, R., Tanaka, H. and Yasuda, H. (1997). Oncoprotein MDM2 is a ubiquitin ligase E3 for tumor suppressor p53. *FEBS Lett.* 420, 25–7.
- Hu, X., Kim, J. A., Castillo, A., Huang, M., Liu, J. and Wang, B. (2011). NBA1/MERIT40 and BRE interaction is required for the integrity of two distinct deubiquitinating enzyme BRCC36-containing complexes. *J. Biol. Chem.* 286, 11734–45.
- Huang, L., Kinnucan, E., Wang, G., Beaudenon, S., Howley, P. M., Huibregtse, J. M. and Pavletich, N. P. (1999). Structure of an E6AP-UbcH7 complex: insights into ubiquitination by the E2-E3 enzyme cascade. *Science* 286, 1321–6.
- Huang, J., Perez-Burgos, L., Placek, B. J., Sengupta, R., Richter, M., Dorsey, J. A., Kubicek, S., Opravil, S., Jenuwein, T. and Berger, S. L. (2006). Repression of p53 activity by Smyd2-mediated methylation. *Nature* 444, 629–32.
- Huang, J., Sengupta, R., Espejo, A. B., Lee, M. G., Dorsey, J. A., Richter, M., Opravil, S., Shiekhhattar, R., Bedford, M. T., Jenuwein, T., et al. (2007). p53 is regulated by the lysine demethylase LSD1. *Nature* 449, 105–8.
- Huibregtse, J. M., Scheffner, M., Beaudenon, S. and Howley, P. M. (1995). A family of proteins structurally and functionally related to the E6-AP ubiquitin-protein ligase. *Proc. Natl. Acad. Sci. U. S. A.* 92, 5249.
- Innocente, S. A., Abrahamson, J. L., Cogswell, J. P. and Lee, J. M. (1999). p53 regulates a G2 checkpoint through cyclin B1. *Proc. Natl. Acad. Sci. U. S. A.* 96, 2147–52.

- Inoki, K., Li, Y., Xu, T. and Guan, K.-L. (2003). Rheb GTPase is a direct target of TSC2 GAP activity and regulates mTOR signaling. *Genes Dev.* 17, 1829–34.
- Itahana, Y., Ke, H. and Zhang, Y. (2009). p53 Oligomerization is essential for its C-terminal lysine acetylation. *J. Biol. Chem.* 284, 5158–64.
- Ito, A., Lai, C. H., Zhao, X., Saito, S., Hamilton, M. H., Appella, E. and Yao, T. P. (2001). p300/CBP-mediated p53 acetylation is commonly induced by p53-activating agents and inhibited by MDM2. *EMBO J.* 20, 1331–40.
- Itoh, M., Kim, C.-H., Palardy, G., Oda, T., Jiang, Y.-J., Maust, D., Yeo, S.-Y., Lorick, K., Wright, G. J., Ariza-McNaughton, L., et al. (2003). Mind bomb is a ubiquitin ligase that is essential for efficient activation of Notch signaling by Delta. *Dev. Cell* 4, 67–82.
- Izawa, N., Wu, W., Sato, K., Nishikawa, H., Kato, A., Boku, N., Itoh, F. and Ohta, T. (2011). HERC2 Interacts with Claspin and regulates DNA origin firing and replication fork progression. *Cancer Res.* 71, 5621–5.
- Jackson, S. P. and Bartek, J. (2009). The DNA-damage response in human biology and disease. *Nature* 461, 1071–8.
- Jansson, M., Durant, S. T., Cho, E.-C., Sheahan, S., Edelman, M., Kessler, B. and La Thangue, N. B. (2008). Arginine methylation regulates the p53 response. *Nat. Cell Biol.* 10, 1431–9.
- Jaroudi, S. and SenGupta, S. DNA repair in mammalian embryos. *Mutat. Res.* 635, 53–77.
- Jeffrey, P. D., Gorina, S. and Pavletich, N. P. (1995). Crystal structure of the tetramerization domain of the p53 tumor suppressor at 1.7 angstroms. *Science* 267, 1498–502.
- Jeon, B.-N., Choi, W.-I., Yu, M.-Y., Yoon, A.-R., Kim, M.-H., Yun, C.-O. and Hur, M.-W. (2009). ZBTB2, a novel master regulator of the p53 pathway. *J. Biol. Chem.* 284, 17935–46.
- Jeon, Y. J., Yoo, H. M. and Chung, C. H. (2010). ISG15 and immune diseases. *Biochim. Biophys. Acta* 1802, 485–96.
- Ji, Y., Walkowicz, M. J., Buiting, K., Johnson, D. K., Tarvin, R. E., Rinchik, E. M., Horsthemke, B., Stubbs, L. and Nicholls, R. D. (1999). The ancestral gene for transcribed, low-copy repeats in the Prader-Willi/Angelman region encodes a large protein implicated in protein trafficking, which is deficient in mice with neuromuscular and spermiogenic abnormalities. *Hum. Mol. Genet.* 8, 533–42.
- Ji, Y., Rebert, N. A., Joslin, J. M., Higgins, M. J., Schultz, R. A. and Nicholls, R. D. (2000). Structure of the highly conserved HERC2 gene and of multiple partially duplicated paralogs in human. *Genome Res.* 10, 319–29.
- Jiang, B.-H. and Liu, L.-Z. (2009). PI3K/PTEN signaling in angiogenesis and tumorigenesis. *Adv. Cancer Res.* 102, 19–65.
- Joerger, A. C. and Fersht, A. R. (2007). Structure-function-rescue: the diverse nature of common p53 cancer mutants. *Oncogene* 26, 2226–42.
- Jones, R. G. and Thompson, C. B. (2009). Tumor suppressors and cell metabolism: a recipe for cancer growth. *Genes Dev.* 23, 537–48.
- Jones, S. N., Roe, A. E., Donehower, L. A. and Bradley, A. (1995). Rescue of embryonic lethality in Mdm2-deficient mice by absence of p53. *Nature* 378, 206–8.

Bibliografia

- Jung, Y. and Lippard, S. J. (2007). Direct cellular responses to platinum-induced DNA damage. *Chem. Rev.* 107, 1387–407.
- Kalab, P., Pu, R. T. and Dasso, M. (1999). The ran GTPase regulates mitotic spindle assembly. *Curr. Biol.* 9, 481–4.
- Kamadurai, H. B., Souphron, J., Scott, D. C., Duda, D. M., Miller, D. J., Stringer, D., Piper, R. C. and Schulman, B. A. (2009). Insights into ubiquitin transfer cascades from a structure of a UbcH5B approximately ubiquitin-HECT(NEDD4L) complex. *Mol. Cell* 36, 1095–102.
- Kang, T.-H., Lindsey-Boltz, L. A., Reardon, J. T. and Sancar, A. (2010). Circadian control of XPA and excision repair of cisplatin-DNA damage by cryptochrome and HERC2 ubiquitin ligase. *Proc. Natl. Acad. Sci. U. S. A.* 107, 4890–5.
- Kang, T.-H., Reardon, J. T. and Sancar, A. (2011). Regulation of nucleotide excision repair activity by transcriptional and post-transcriptional control of the XPA protein. *Nucleic Acids Res.* 39, 3176–87.
- Kasper, J. S., Arai, T. and DeCaprio, J. A. (2006). A novel p53-binding domain in CUL7. *Biochem. Biophys. Res. Commun.* 348, 132–8.
- Kaustov, L., Lukin, J., Lemak, A., Duan, S., Ho, M., Doherty, R., Penn, L. Z. and Arrowsmith, C. H. (2007). The conserved CPH domains of Cul7 and PARC are protein-protein interaction modules that bind the tetramerization domain of p53. *J. Biol. Chem.* 282, 11300–7.
- Kawaguchi, T., Kato, S., Otsuka, K., Watanabe, G., Kumabe, T., Tominaga, T., Yoshimoto, T. and Ishioka, C. (2005). The relationship among p53 oligomer formation, structure and transcriptional activity using a comprehensive missense mutation library. *Oncogene* 24, 6976–81.
- Kawaguchi, Y., Ito, A., Appella, E. and Yao, T.-P. (2006). Charge modification at multiple C-terminal lysine residues regulates p53 oligomerization and its nucleus-cytoplasm trafficking. *J. Biol. Chem.* 281, 1394–400.
- Kawai, K. and Akaza, H. (2010). Current status of chemotherapy in risk-adapted management for metastatic testicular germ cell cancer. *Cancer Sci.* 101, 22–8.
- Kennedy, K. M. and Dewhirst, M. W. (2010). Tumor metabolism of lactate: the influence and therapeutic potential for MCT and CD147 regulation. *Future Oncol.* 6, 127–48.
- Kerscher, O., Felberbaum, R. and Hochstrasser, M. (2006). Modification of proteins by ubiquitin and ubiquitin-like proteins. *Annu. Rev. Cell Dev. Biol.* 22, 159–80.
- Ketscher, L., Basters, A., Prinz, M. and Knobloch, K.-P. (2012). mHERC6 is the essential ISG15 E3 ligase in the murine system. *Biochem. Biophys. Res. Commun.* 417, 135–40.
- Kim, K. II and Zhang, D.-E. (2003). ISG15, not just another ubiquitin-like protein. *Biochem. Biophys. Res. Commun.* 307, 431–4.
- Kishino, T., Lalonde, M. and Wagstaff, J. (1997). UBE3A/E6-AP mutations cause Angelman syndrome. *Nat. Genet.* 15, 70–3.
- Kitayner, M., Rozenberg, H., Kessler, N., Rabinovich, D., Shaulov, L., Haran, T. E. and Shaked, Z. (2006). Structural basis of DNA recognition by p53 tetramers. *Mol. Cell* 22, 741–53.
- Kleijnen, M. F., Shih, A. H., Zhou, P., Kumar, S., Soccio, R. E., Kedersha, N. L., Gill, G. and Howley, P. M. (2000). The hPLIC proteins may provide a link between the ubiquitination machinery and the proteasome. *Mol. Cell* 6, 409–19.

- Ko, L. J. and Prives, C. (1996). p53: puzzle and paradigm. *Genes Dev.* 10, 1054–72.
- Komander, D. and Rape, M. (2012). The ubiquitin code. *Annu. Rev. Biochem.* 81, 203–29.
- Kornitzer, D. and Ciechanover, A. (2000). Modes of regulation of ubiquitin-mediated protein degradation. *J. Cell. Physiol.* 182, 1–11.
- Koseki, T., Inohara, N., Chen, S. and Núñez, G. (1998). ARC, an inhibitor of apoptosis expressed in skeletal muscle and heart that interacts selectively with caspases. *Proc. Natl. Acad. Sci. U. S. A.* 95, 5156–60.
- Krause, K., Wasner, M., Reinhard, W., Haugwitz, U., Dohna, C. L., Mössner, J. and Engeland, K. (2000). The tumour suppressor protein p53 can repress transcription of cyclin B. *Nucleic Acids Res.* 28, 4410–8.
- Kravtsova-Ivantsiv, Y. and Ciechanover, A. (2012). Non-canonical ubiquitin-based signals for proteasomal degradation. *J. Cell Sci.* 125, 539–48.
- Kroismayr, R., Baranyi, U., Stehlik, C., Dorfleutner, A., Binder, B. R. and Lipp, J. (2004). HERC5, a HECT E3 ubiquitin ligase tightly regulated in LPS activated endothelial cells. *J. Cell Sci.* 117, 4749–56.
- Kruse, J.-P. and Gu, W. (2009). Modes of p53 regulation. *Cell* 137, 609–22.
- Kubbutat, M. H., Jones, S. N. and Vousden, K. H. (1997). Regulation of p53 stability by Mdm2. *Nature* 387, 299–303.
- Kubo, K. (1995). Effect of incubation of solutions of proteins containing dodecyl sulfate on the cleavage of peptide bonds by boiling. *Anal. Biochem.* 225, 351–3.
- Kubota, T., Matsuoka, M., Chang, T.-H., Taylor, P., Sasaki, T., Tashiro, M., Kato, A. and Ozato, K. (2008). Virus infection triggers SUMOylation of IRF3 and IRF7, leading to the negative regulation of type I interferon gene expression. *J. Biol. Chem.* 283, 25660–70.
- Kühnle, S., Kogel, U., Glockzin, S., Marquardt, A., Ciechanover, A., Matentzoglou, K. and Scheffner, M. (2011). Physical and functional interaction of the HECT ubiquitin-protein ligases E6AP and HERC2. *J. Biol. Chem.* 286, 19410–6.
- Kulathu, Y. and Komander, D. (2012). Atypical ubiquitylation - the unexplored world of polyubiquitin beyond Lys48 and Lys63 linkages. *Nat. Rev. Mol. Cell Biol.* 13, 508–23.
- Kumar, K. P., McBride, K. M., Weaver, B. K., Dingwall, C. and Reich, N. C. (2000). Regulated nuclear-cytoplasmic localization of interferon regulatory factor 3, a subunit of double-stranded RNA-activated factor 1. *Mol. Cell. Biol.* 20, 4159–68.
- Laan, L. A., v Haeringen, A. and Brouwer, O. F. (1999). Angelman syndrome: a review of clinical and genetic aspects. *Clin. Neurol. Neurosurg.* 101, 161–70.
- Laptenko, O. and Prives, C. (2006). Transcriptional regulation by p53: one protein, many possibilities. *Cell Death Differ.* 13, 951–61.
- Laronga, C., Yang, H. Y., Neal, C. and Lee, M. H. (2000). Association of the cyclin-dependent kinases and 14-3-3 sigma negatively regulates cell cycle progression. *J. Biol. Chem.* 275, 23106–12.
- Latil, A., Morant, P., Fournier, G., Mangin, P., Berthon, P. and Cussenot, O. (2002). CHC1-L, a candidate gene for prostate carcinogenesis at 13q14.2, is frequently affected by loss of heterozygosity and underexpressed in human prostate cancer. *Int. J. Cancer* 99, 689–96.

Bibliografia

- Lavigne, A., Maltby, V., Mock, D., Rossant, J., Pawson, T. and Bernstein, A. (1989). High incidence of lung, bone, and lymphoid tumors in transgenic mice overexpressing mutant alleles of the p53 oncogene. *Mol. Cell. Biol.* 9, 3982–91.
- Lebel, M., Cardiff, R. D. and Leder, P. (2001). Tumorigenic effect of nonfunctional p53 or p21 in mice mutant in the Werner syndrome helicase. *Cancer Res.* 61, 1816–9.
- Lee, J. T. and Gu, W. (2010). The multiple levels of regulation by p53 ubiquitination. *Cell Death Differ.* 17, 86–92.
- Lee, J.-H. and Paull, T. T. (2007). Activation and regulation of ATM kinase activity in response to DNA double-strand breaks. *Oncogene* 26, 7741–8.
- Lee, M. H., Reynisdóttir, I. and Massagué, J. (1995). Cloning of p57KIP2, a cyclin-dependent kinase inhibitor with unique domain structure and tissue distribution. *Genes Dev.* 9, 639–49.
- Lee, J., Gold, D. A., Shevchenko, A., Shevchenko, A. and Dunphy, W. G. (2005). Roles of replication fork-interacting and Chk1-activating domains from Claspin in a DNA replication checkpoint response. *Mol. Biol. Cell* 16, 5269–82.
- Lee, T.-H., Park, J.-M., Leem, S.-H. and Kang, T.-H. (2014). Coordinated regulation of XPA stability by ATR and HERC2 during nucleotide excision repair. *Oncogene* 33, 19–25.
- Lehman, A. L., Nakatsu, Y., Ching, A., Bronson, R. T., Oakey, R. J., Keiper-Hrynko, N., Finger, J. N., Durham-Pierre, D., Horton, D. B., Newton, J. M., et al. (1998). A very large protein with diverse functional motifs is deficient in rjs (runty, jerky, sterile) mice. *Proc. Natl. Acad. Sci. U. S. A.* 95, 9436–41.
- Leng, R. P., Lin, Y., Ma, W., Wu, H., Lemmers, B., Chung, S., Parant, J. M., Lozano, G., Hakem, R. and Benchimol, S. (2003). Pirh2, a p53-induced ubiquitin-protein ligase, promotes p53 degradation. *Cell* 112, 779–91.
- Leu, J. I.-J., Dumont, P., Hafey, M., Murphy, M. E. and George, D. L. (2004). Mitochondrial p53 activates Bak and causes disruption of a Bak-Mcl1 complex. *Nat. Cell Biol.* 6, 443–50.
- Levine, A. J. (1997). p53, the cellular gatekeeper for growth and division. *Cell* 88, 323–31.
- Levine, A. J. and Oren, M. (2009). The first 30 years of p53: growing ever more complex. *Nat. Rev. Cancer* 9, 749–58.
- Li, M., Brooks, C. L., Wu-Baer, F., Chen, D., Baer, R. and Gu, W. (2003). Mono- versus polyubiquitination: differential control of p53 fate by Mdm2. *Science* 302, 1972–5.
- Li, T., Kon, N., Jiang, L., Tan, M., Ludwig, T., Zhao, Y., Baer, R. and Gu, W. (2012). Tumor suppression in the absence of p53-mediated cell-cycle arrest, apoptosis, and senescence. *Cell* 149, 1269–83.
- Liang, S. H. and Clarke, M. F. (2001). Regulation of p53 localization. *Eur. J. Biochem.* 268, 2779–83.
- Lin, R., Heylbroeck, C., Pitha, P. M. and Hiscott, J. (1998). Virus-dependent phosphorylation of the IRF-3 transcription factor regulates nuclear translocation, transactivation potential, and proteasome-mediated degradation. *Mol. Cell. Biol.* 18, 2986–96.
- Lindahl, T. and Barnes, D. E. (2000). Repair of endogenous DNA damage. *Cold Spring Harb. Symp. Quant. Biol.* 65, 127–33.
- Lindström, M. S., Deisenroth, C. and Zhang, Y. (2007). Putting a finger on growth surveillance: insight into MDM2 zinc finger-ribosomal protein interactions. *Cell Cycle* 6, 434–7.

- Linnert, M. and Gehl, J. (2009). Bleomycin treatment of brain tumors: an evaluation. *Anticancer. Drugs* 20, 157–64.
- Liu, L., Scolnick, D. M., Trievel, R. C., Zhang, H. B., Marmorstein, R., Halazonetis, T. D. and Berger, S. L. (1999). p53 sites acetylated in vitro by PCAF and p300 are acetylated in vivo in response to DNA damage. *Mol. Cell. Biol.* 19, 1202–9.
- Liu, W., Quinto, I., Chen, X., Palmieri, C., Rabin, R. L., Schwartz, O. M., Nelson, D. L. and Scala, G. (2001). Direct inhibition of Bruton's tyrosine kinase by IBtk, a Btk-binding protein. *Nat. Immunol.* 2, 939–46.
- Liu, F., van Duijn, K., Vingerling, J. R., Hofman, A., Uitterlinden, A. G., Janssens, A. C. J. W. and Kayser, M. (2009). Eye color and the prediction of complex phenotypes from genotypes. *Curr. Biol.* 19, R192–3.
- Liu, J., Hutchison, K., Perrone-Bizzozero, N., Morgan, M., Sui, J. and Calhoun, V. (2010). Identification of genetic and epigenetic marks involved in population structure. *PLoS One* 5, e13209.
- Long, X., Lin, Y., Ortiz-Vega, S., Yonezawa, K. and Avruch, J. (2005). Rheb binds and regulates the mTOR kinase. *Curr. Biol.* 15, 702–13.
- Lowe, S. W. and Sherr, C. J. (2003). Tumor suppression by Ink4a-Arf: progress and puzzles. *Curr. Opin. Genet. Dev.* 13, 77–83.
- Lu, G., Reinert, J. T., Pitha-Rowe, I., Okumura, A., Kellum, M., Knobloch, K. P., Hassel, B. and Pitha, P. M. (2006). ISG15 enhances the innate antiviral response by inhibition of IRF-3 degradation. *Cell. Mol. Biol. (Noisy-le-grand)*. 52, 29–41.
- Lu, J.-Y., Lin, Y.-Y., Qian, J., Tao, S.-C., Zhu, J., Pickart, C. and Zhu, H. (2008). Functional dissection of a HECT ubiquitin E3 ligase. *Mol. Cell. Proteomics* 7, 35–45.
- Lukas, C., Savic, V., Bekker-Jensen, S., Doil, C., Neumann, B., Pedersen, R. S., Grøfte, M., Chan, K. L., Hickson, I. D., Bartek, J., et al. (2011). 53BP1 nuclear bodies form around DNA lesions generated by mitotic transmission of chromosomes under replication stress. *Nat. Cell Biol.* 13, 243–53.
- Lyon, M. F., King, T. R., Gondo, Y., Gardner, J. M., Nakatsu, Y., Eicher, E. M. and Brilliant, M. H. (1992). Genetic and molecular analysis of recessive alleles at the pink-eyed dilution (p) locus of the mouse. *Proc. Natl. Acad. Sci. U. S. A.* 89, 6968–72.
- Makde, R. D., England, J. R., Yennawar, H. P. and Tan, S. (2010). Structure of RCC1 chromatin factor bound to the nucleosome core particle. *Nature* 467, 562–6.
- Maki, C. G. (1999). Oligomerization is required for p53 to be efficiently ubiquitinated by MDM2. *J. Biol. Chem.* 274, 16531–5.
- Malkin, D. (2011). Li-fraumeni syndrome. *Genes Cancer* 2, 475–84.
- Mashimo, T., Hadjebi, O., Amair-Pinedo, F., Tsurumi, T., Langa, F., Serikawa, T., Sotelo, C., Guénet, J.-L. and Rosa, J. L. (2009). Progressive Purkinje cell degeneration in tambaleante mutant mice is a consequence of a missense mutation in HERC1 E3 ubiquitin ligase. *PLoS Genet.* 5, e1000784.
- Massagué, J. (2008). TGFbeta in Cancer. *Cell* 134, 215–30.
- McBride, K. A., Ballinger, M. L., Killick, E., Kirk, J., Tattersall, M. H. N., Eeles, R. A., Thomas, D. M. and Mitchell, G. (2014). Li-Fraumeni syndrome: cancer risk assessment and clinical management. *Nat. Rev. Clin. Oncol.*

- Merla, G., Ucla, C., Guipponi, M. and Reymond, A. (2002). Identification of additional transcripts in the Williams-Beuren syndrome critical region. *Hum. Genet.* 110, 429–38.
- Metzger, M. B., Hristova, V. A. and Weissman, A. M. (2012). HECT and RING finger families of E3 ubiquitin ligases at a glance. *J. Cell Sci.* 125, 531–7.
- Michalak, E. M., Villunger, A., Adams, J. M. and Strasser, A. (2008). In several cell types tumour suppressor p53 induces apoptosis largely via Puma but Noxa can contribute. *Cell Death Differ.* 15, 1019–29.
- Mitsui, K., Nakanishi, M., Ohtsuka, S., Norwood, T. H., Okabayashi, K., Miyamoto, C., Tanaka, K., Yoshimura, A. and Ohtsubo, M. (1999). A novel human gene encoding HECT domain and RCC1-like repeats interacts with cyclins and is potentially regulated by the tumor suppressor proteins. *Biochem. Biophys. Res. Commun.* 266, 115–22.
- Miyashita, T. and Reed, J. C. (1995). Tumor suppressor p53 is a direct transcriptional activator of the human bax gene. *Cell* 80, 293–9.
- Moll, U. M., Wolff, S., Speidel, D. and Deppert, W. (2005). Transcription-independent pro-apoptotic functions of p53. *Curr. Opin. Cell Biol.* 17, 631–6.
- Mollinari, C., Reynaud, C., Martineau-Thuillier, S., Monier, S., Kieffer, S., Garin, J., Andreassen, P. R., Boulet, A., Goud, B., Kleman, J.-P., et al. (2003). The mammalian passenger protein TD-60 is an RCC1 family member with an essential role in prometaphase to metaphase progression. *Dev. Cell* 5, 295–307.
- Montes de Oca Luna, R., Wagner, D. S. and Lozano, G. (1995). Rescue of early embryonic lethality in mdm2-deficient mice by deletion of p53. *Nature* 378, 203–6.
- Morgan, D. O. (1995). Principles of CDK regulation. *Nature* 374, 131–4.
- Moroishi, T., Nishiyama, M., Takeda, Y., Iwai, K. and Nakayama, K. I. (2011). The FBXL5-IRP2 axis is integral to control of iron metabolism in vivo. *Cell Metab.* 14, 339–51.
- Moroishi, T., Yamauchi, T., Nishiyama, M. and Nakayama, K. I. (2014). HERC2 Targets the Iron Regulator FBXL5 for Degradation and Modulates Iron Metabolism. *J. Biol. Chem.* 289, 16430–16441.
- Muzio, M. (1998). Signalling by proteolysis: death receptors induce apoptosis. *Int. J. Clin. Lab. Res.* 28, 141–7.
- Nagaich, A. K., Zhurkin, V. B., Sakamoto, H., Gorin, A. A., Clore, G. M., Gronenborn, A. M., Appella, E. and Harrington, R. E. (1997). Architectural accommodation in the complex of four p53 DNA binding domain peptides with the p21/waf1/cip1 DNA response element. *J. Biol. Chem.* 272, 14830–41.
- Nagase, T., Kikuno, R., Hattori, A., Kondo, Y., Okumura, K. and Ohara, O. (2000). Prediction of the coding sequences of unidentified human genes. XIX. The complete sequences of 100 new cDNA clones from brain which code for large proteins in vitro. *DNA Res.* 7, 347–55.
- Nagata, S. and Golstein, P. (1995). The Fas death factor. *Science* 267, 1449–56.
- Nakano, K. and Vousden, K. H. (2001). PUMA, a novel proapoptotic gene, is induced by p53. *Mol. Cell* 7, 683–94.
- Nemergut, M. E., Mizzen, C. A., Stukenberg, T., Allis, C. D. and Macara, I. G. (2001). Chromatin docking and exchange activity enhancement of RCC1 by histones H2A and H2B. *Science* 292, 1540–3.

- Nicholls, R. D. and Knepper, J. L. (2001). Genome organization, function, and imprinting in Prader-Willi and Angelman syndromes. *Annu. Rev. Genomics Hum. Genet.* 2, 153–75.
- Nicholls, C. D., McLure, K. G., Shields, M. A. and Lee, P. W. K. (2002). Biogenesis of p53 involves cotranslational dimerization of monomers and posttranslational dimerization of dimers. Implications on the dominant negative effect. *J. Biol. Chem.* 277, 12937–45.
- Nikolaev, A. Y., Li, M., Puskas, N., Qin, J. and Gu, W. (2003). Parc: a cytoplasmic anchor for p53. *Cell* 112, 29–40.
- Nomura, N., Miyajima, N., Sazuka, T., Tanaka, A., Kawarabayasi, Y., Sato, S., Nagase, T., Seki, N., Ishikawa, K. and Tabata, S. (1994). Prediction of the coding sequences of unidentified human genes. I. The coding sequences of 40 new genes (KIAA0001-KIAA0040) deduced by analysis of randomly sampled cDNA clones from human immature myeloid cell line KG-1. *DNA Res.* 1, 27–35.
- O'Keefe, K., Li, H. and Zhang, Y. (2003). Nucleocytoplasmic shuttling of p53 is essential for MDM2-mediated cytoplasmic degradation but not ubiquitination. *Mol. Cell. Biol.* 23, 6396–405.
- O'Reilly, K. E., Rojo, F., She, Q.-B., Solit, D., Mills, G. B., Smith, D., Lane, H., Hofmann, F., Hicklin, D. J., Ludwig, D. L., et al. (2006). mTOR inhibition induces upstream receptor tyrosine kinase signaling and activates Akt. *Cancer Res.* 66, 1500–8.
- Oda, E., Ohki, R., Murasawa, H., Nemoto, J., Shibue, T., Yamashita, T., Tokino, T., Taniguchi, T. and Tanaka, N. (2000a). Noxa, a BH3-only member of the Bcl-2 family and candidate mediator of p53-induced apoptosis. *Science* 288, 1053–8.
- Oda, K., Arakawa, H., Tanaka, T., Matsuda, K., Tanikawa, C., Mori, T., Nishimori, H., Tamai, K., Tokino, T., Nakamura, Y., et al. (2000b). p53AIP1, a potential mediator of p53-dependent apoptosis, and its regulation by Ser-46-phosphorylated p53. *Cell* 102, 849–62.
- Oestergaard, V. H., Pentzold, C., Pedersen, R. T., losif, S., Alpi, A., Bekker-Jensen, S., Mailand, N. and Lisby, M. (2012). RNF8 and RNF168 but not HERC2 are required for DNA damage-induced ubiquitylation in chicken DT40 cells. *DNA Repair (Amst).* 11, 892–905.
- Ogunjimi, A. A., Briant, D. J., Pece-Barbara, N., Le Roy, C., Di Guglielmo, G. M., Kavsak, P., Rasmussen, R. K., Seet, B. T., Sicheri, F. and Wrana, J. L. (2005). Regulation of Smurf2 ubiquitin ligase activity by anchoring the E2 to the HECT domain. *Mol. Cell* 19, 297–308.
- Ohtsubo, M., Kai, R., Furuno, N., Sekiguchi, T., Sekiguchi, M., Hayashida, H., Kuma, K., Miyata, T., Fukushige, S. and Murotsu, T. (1987). Isolation and characterization of the active cDNA of the human cell cycle gene (RCC1) involved in the regulation of onset of chromosome condensation. *Genes Dev.* 1, 585–93.
- Okada, T., Lopez-Lago, M. and Giancotti, F. G. (2005). Merlin/NF-2 mediates contact inhibition of growth by suppressing recruitment of Rac to the plasma membrane. *J. Cell Biol.* 171, 361–71.
- Okoshi, R., Ozaki, T., Yamamoto, H., Ando, K., Koida, N., Ono, S., Koda, T., Kamijo, T., Nakagawara, A. and Kizaki, H. (2008). Activation of AMP-activated protein kinase induces p53-dependent apoptotic cell death in response to energetic stress. *J. Biol. Chem.* 283, 3979–87.
- Ono, W., Hayashi, Y., Yokoyama, W., Kuroda, T., Kishimoto, H., Ito, I., Kimura, K., Akaogi, K., Waku, T. and Yanagisawa, J. (2014). The Nucleolar Protein Myb-binding Protein 1A (MYBBP1A) Enhances p53 Tetramerization and Acetylation in Response to Nucleolar Disruption. *J. Biol. Chem.* 289, 4928–40.

Bibliografia

- Ou, Y.-H., Chung, P.-H., Sun, T.-P. and Shieh, S.-Y. (2005). p53 C-terminal phosphorylation by CHK1 and CHK2 participates in the regulation of DNA-damage-induced C-terminal acetylation. *Mol. Biol. Cell* 16, 1684–95.
- Oudshoorn, D., van Boheemen, S., Sánchez-Aparicio, M. T., Rajsbaum, R., García-Sastre, A. and Versteeg, G. A. (2012). HERC6 is the main E3 ligase for global ISG15 conjugation in mouse cells. *PLoS One* 7, e29870.
- Papaioannou, V. E. and Behringer, R. R. (2012). Early embryonic lethality in genetically engineered mice: diagnosis and phenotypic analysis. *Vet. Pathol.* 49, 64–70.
- Parant, J., Chavez-Reyes, A., Little, N. A., Yan, W., Reinke, V., Jochemsen, A. G. and Lozano, G. (2001). Rescue of embryonic lethality in Mdm4-null mice by loss of Trp53 suggests a nonoverlapping pathway with MDM2 to regulate p53. *Nat. Genet.* 29, 92–5.
- Park, J.-I., Venteicher, A. S., Hong, J. Y., Choi, J., Jun, S., Shkreli, M., Chang, W., Meng, Z., Cheung, P., Ji, H., et al. (2009). Telomerase modulates Wnt signalling by association with target gene chromatin. *Nature* 460, 66–72.
- Pavletich, N. P., Chambers, K. A. and Pabo, C. O. (1993). The DNA-binding domain of p53 contains the four conserved regions and the major mutation hot spots. *Genes Dev.* 7, 2556–64.
- Perfettini, J.-L., Castedo, M., Nardacci, R., Ciccocanti, F., Boya, P., Roumier, T., Larochette, N., Piacentini, M. and Kroemer, G. (2005). Essential role of p53 phosphorylation by p38 MAPK in apoptosis induction by the HIV-1 envelope. *J. Exp. Med.* 201, 279–89.
- Perona, R. (2006). Cell signalling: growth factors and tyrosine kinase receptors. *Clin. Transl. Oncol.* 8, 77–82.
- Petermann, E., Helleday, T. and Caldecott, K. W. (2008). Claspin promotes normal replication fork rates in human cells. *Mol. Biol. Cell* 19, 2373–8.
- Petersen, S., Casellas, R., Reina-San-Martin, B., Chen, H. T., Difilippantonio, M. J., Wilson, P. C., Hanitsch, L., Celeste, A., Muramatsu, M., Pilch, D. R., et al. (2001). AID is required to initiate Nbs1/gamma-H2AX focus formation and mutations at sites of class switching. *Nature* 414, 660–5.
- Pośpiech, E., Draus-Barini, J., Kupiec, T., Wojas-Pelc, A. and Branicki, W. (2011). Gene-gene interactions contribute to eye colour variation in humans. *J. Hum. Genet.* 56, 447–55.
- Poyurovsky, M. V., Priest, C., Kentsis, A., Borden, K. L. B., Pan, Z.-Q., Pavletich, N. and Prives, C. (2007). The Mdm2 RING domain C-terminus is required for supramolecular assembly and ubiquitin ligase activity. *EMBO J.* 26, 90–101.
- Puffenberger, E. G., Jinks, R. N., Wang, H., Xin, B., Fiorentini, C., Sherman, E. A., Degrazio, D., Shaw, C., Sougnez, C., Cibulskis, K., et al. (2012). A homozygous missense mutation in HERC2 associated with global developmental delay and autism spectrum disorder. *Hum. Mutat.* 33, 1639–46.
- Raza, A., Franklin, M. J. and Dudek, A. Z. (2010). Pericytes and vessel maturation during tumor angiogenesis and metastasis. *Am. J. Hematol.* 85, 593–8.
- Reardon, J. T. and Sancar, A. (2005). Nucleotide excision repair. *Prog. Nucleic Acid Res. Mol. Biol.* 79, 183–235.
- Rees, J. L. (2003). Genetics of hair and skin color. *Annu. Rev. Genet.* 37, 67–90.
- Renault, L., Nassar, N., Vetter, I., Becker, J., Klebe, C., Roth, M. and Wittinghofer, A. (1998). The 1.7 Å crystal structure of the regulator of chromosome condensation (RCC1) reveals a seven-bladed propeller. *Nature* 392, 97–101.

- Renault, L., Nassar, N., Wittinghofer, A., Roth, M. and Vetter, I. R. (1999). Crystallization and preliminary X-ray analysis of human RCC1, the regulator of chromosome condensation. *Acta Crystallogr. D. Biol. Crystallogr.* 55, 272–5.
- Renault, L., Kuhlmann, J., Henkel, A. and Wittinghofer, A. (2001). Structural basis for guanine nucleotide exchange on Ran by the regulator of chromosome condensation (RCC1). *Cell* 105, 245–55.
- Ribbeck, K., Lipowsky, G., Kent, H. M., Stewart, M. and Görlich, D. (1998). NTF2 mediates nuclear import of Ran. *EMBO J.* 17, 6587–98.
- Rinaldo, C., Prodosmo, A., Mancini, F., Iacovelli, S., Sacchi, A., Moretti, F. and Soddu, S. (2007). MDM2-regulated degradation of HIPK2 prevents p53Ser46 phosphorylation and DNA damage-induced apoptosis. *Mol. Cell* 25, 739–50.
- Ringshausen, I., O'Shea, C. C., Finch, A. J., Swigart, L. B. and Evan, G. I. (2006). Mdm2 is critically and continuously required to suppress lethal p53 activity in vivo. *Cancer Cell* 10, 501–14.
- Rivera, A. and Maxwell, S. A. (2005). The p53-induced gene-6 (proline oxidase) mediates apoptosis through a calcineurin-dependent pathway. *J. Biol. Chem.* 280, 29346–54.
- Rodriguez, C. I. and Stewart, C. L. (2007). Disruption of the ubiquitin ligase HERC4 causes defects in spermatozoon maturation and impaired fertility. *Dev. Biol.* 312, 501–8.
- Rodriguez, M. S., Desterro, J. M., Lain, S., Lane, D. P. and Hay, R. T. (2000). Multiple C-terminal lysine residues target p53 for ubiquitin-proteasome-mediated degradation. *Mol. Cell. Biol.* 20, 8458–67.
- Rosa, J. L. and Barbacid, M. (1997). A giant protein that stimulates guanine nucleotide exchange on ARF1 and Rab proteins forms a cytosolic ternary complex with clathrin and Hsp70. *Oncogene* 15, 1–6.
- Rosa, J. L., Casaroli-Marano, R. P., Buckler, A. J., Vilaró, S. and Barbacid, M. (1996). p619, a giant protein related to the chromosome condensation regulator RCC1, stimulates guanine nucleotide exchange on ARF1 and Rab proteins. *EMBO J.* 15, 4262–73.
- Rotin, D. and Kumar, S. (2009). Physiological functions of the HECT family of ubiquitin ligases. *Nat. Rev. Mol. Cell Biol.* 10, 398–409.
- Rougeulle, C., Glatt, H. and Lalande, M. (1997). The Angelman syndrome candidate gene, UBE3A/E6-AP, is imprinted in brain. *Nat. Genet.* 17, 14–5.
- Rubbi, C. P. and Milner, J. (2003). Disruption of the nucleolus mediates stabilization of p53 in response to DNA damage and other stresses. *EMBO J.* 22, 6068–77.
- Russell, R. C., Fang, C. and Guan, K.-L. (2011). An emerging role for TOR signaling in mammalian tissue and stem cell physiology. *Development* 138, 3343–56.
- Saitoh, T., Tun-Kyi, A., Ryo, A., Yamamoto, M., Finn, G., Fujita, T., Akira, S., Yamamoto, N., Lu, K. P. and Yamaoka, S. (2006). Negative regulation of interferon-regulatory factor 3-dependent innate antiviral response by the prolyl isomerase Pin1. *Nat. Immunol.* 7, 598–605.
- Sandri, M. I., Isaacs, R. J., Ongkeko, W. M., Harris, A. L., Hickson, I. D., Broggin, M. and Vikhanskaya, F. (1996). p53 regulates the minimal promoter of the human topoisomerase IIalpha gene. *Nucleic Acids Res.* 24, 4464–70.

Bibliografia

- Sato, M., Suemori, H., Hata, N., Asagiri, M., Ogasawara, K., Nakao, K., Nakaya, T., Katsuki, M., Noguchi, S., Tanaka, N., et al. (2000). Distinct and essential roles of transcription factors IRF-3 and IRF-7 in response to viruses for IFN- α / β gene induction. *Immunity* 13, 539–48.
- Schägger, H. (2006). Tricine-SDS-PAGE. *Nat. Protoc.* 1, 16–22.
- Scheffner, M. and Kumar, S. (2014). Mammalian HECT ubiquitin-protein ligases: biological and pathophysiological aspects. *Biochim. Biophys. Acta* 1843, 61–74.
- Scheffner, M. and Staub, O. (2007). HECT E3s and human disease. *BMC Biochem.* 8 Suppl 1, S6.
- Scheffner, M., Huijbregtse, J. M., Vierstra, R. D. and Howley, P. M. (1993). The HPV-16 E6 and E6-AP complex functions as a ubiquitin-protein ligase in the ubiquitination of p53. *Cell* 75, 495–505.
- Schumacher, B., Mondry, J., Thiel, P., Weyand, M. and Ottmann, C. (2010). Structure of the p53 C-terminus bound to 14-3-3: implications for stabilization of the p53 tetramer. *FEBS Lett.* 584, 1443–8.
- Schwartz, A. L. and Ciechanover, A. (2009). Targeting proteins for destruction by the ubiquitin system: implications for human pathobiology. *Annu. Rev. Pharmacol. Toxicol.* 49, 73–96.
- Schwarz, S. E., Rosa, J. L. and Scheffner, M. (1998). Characterization of human hect domain family members and their interaction with UbcH5 and UbcH7. *J. Biol. Chem.* 273, 12148–54.
- Seki, T., Hayashi, N. and Nishimoto, T. (1996). RCC1 in the Ran pathway. *J. Biochem.* 120, 207–14.
- Shangary, S. and Wang, S. (2009). Small-molecule inhibitors of the MDM2-p53 protein-protein interaction to reactivate p53 function: a novel approach for cancer therapy. *Annu. Rev. Pharmacol. Toxicol.* 49, 223–41.
- Sharp, D. A., Kratowicz, S. A., Sank, M. J. and George, D. L. (1999). Stabilization of the MDM2 oncoprotein by interaction with the structurally related MDMX protein. *J. Biol. Chem.* 274, 38189–96.
- Sheng, Y., Laister, R. C., Lemak, A., Wu, B., Tai, E., Duan, S., Lukin, J., Sunnerhagen, M., Srisailam, S., Karra, M., et al. (2008). Molecular basis of Pirh2-mediated p53 ubiquitylation. *Nat. Struct. Mol. Biol.* 15, 1334–42.
- Sherr, C. J. (1994). Growth factor-regulated G1 cyclins. *Stem Cells* 12 Suppl 1, 47–55; discussion 55–7.
- Sherr, C. J. (2006). Divorcing ARF and p53: an unsettled case. *Nat. Rev. Cancer* 6, 663–73.
- Sherr, C. J. and Roberts, J. M. (1995). Inhibitors of mammalian G1 cyclin-dependent kinases. *Genes Dev.* 9, 1149–63.
- Shi, X., Kachirskaja, I., Yamaguchi, H., West, L. E., Wen, H., Wang, E. W., Dutta, S., Appella, E. and Gozani, O. (2007). Modulation of p53 function by SET8-mediated methylation at lysine 382. *Mol. Cell* 27, 636–46.
- Shi, H.-X., Yang, K., Liu, X., Liu, X.-Y., Wei, B., Shan, Y.-F., Zhu, L.-H. and Wang, C. (2010). Positive regulation of interferon regulatory factor 3 activation by Herc5 via ISG15 modification. *Mol. Cell. Biol.* 30, 2424–36.
- Sjölander, M., Uhlmann, J. and Ponstingl, H. (2002). DelGEF, a homologue of the Ran guanine nucleotide exchange factor RanGEF, binds to the exocyst component Sec5 and modulates secretion. *FEBS Lett.* 532, 211–5.
- Sjölander, M., Uhlmann, J. and Ponstingl, H. (2004). Characterisation of an evolutionary conserved protein interacting with the putative guanine nucleotide exchange factor DelGEF and modulating secretion. *Exp. Cell Res.* 294, 68–76.

- Skaug, B. and Chen, Z. J. (2010). Emerging role of ISG15 in antiviral immunity. *Cell* 143, 187–90.
- Smith, M. L., Chen, I. T., Zhan, Q., Bae, I., Chen, C. Y., Gilmer, T. M., Kastan, M. B., O'Connor, P. M. and Fornace, A. J. (1994). Interaction of the p53-regulated protein Gadd45 with proliferating cell nuclear antigen. *Science* 266, 1376–80.
- Spatuzza, C., Schiavone, M., Di Salle, E., Janda, E., Sardiello, M., Fiume, G., Fierro, O., Simonetta, M., Argiriou, N., Faraonio, R., et al. (2008). Physical and functional characterization of the genetic locus of IBtk, an inhibitor of Bruton's tyrosine kinase: evidence for three protein isoforms of IBtk. *Nucleic Acids Res.* 36, 4402–16.
- Stehmeier, P. and Muller, S. (2009). Regulation of p53 family members by the ubiquitin-like SUMO system. *DNA Repair (Amst)*. 8, 491–8.
- Stommel, J. M., Marchenko, N. D., Jimenez, G. S., Moll, U. M., Hope, T. J. and Wahl, G. M. (1999). A leucine-rich nuclear export signal in the p53 tetramerization domain: regulation of subcellular localization and p53 activity by NES masking. *EMBO J.* 18, 1660–72.
- Sturm, R. A. and Larsson, M. (2009). Genetics of human iris colour and patterns. *Pigment Cell Melanoma Res.* 22, 544–62.
- Sulem, P., Gudbjartsson, D. F., Stacey, S. N., Helgason, A., Rafnar, T., Magnusson, K. P., Manolescu, A., Karason, A., Palsson, A., Thorleifsson, G., et al. (2007). Genetic determinants of hair, eye and skin pigmentation in Europeans. *Nat. Genet.* 39, 1443–52.
- Sykes, S. M., Mellert, H. S., Holbert, M. A., Li, K., Marmorstein, R., Lane, W. S. and McMahon, S. B. (2006). Acetylation of the p53 DNA-binding domain regulates apoptosis induction. *Mol. Cell* 24, 841–51.
- Taira, N., Nihira, K., Yamaguchi, T., Miki, Y. and Yoshida, K. (2007). DYRK2 is targeted to the nucleus and controls p53 via Ser46 phosphorylation in the apoptotic response to DNA damage. *Mol. Cell* 25, 725–38.
- Takagi, M., Absalon, M. J., McLure, K. G. and Kastan, M. B. (2005). Regulation of p53 translation and induction after DNA damage by ribosomal protein L26 and nucleolin. *Cell* 123, 49–63.
- Tam, P. P., Parameswaran, M., Kinder, S. J. and Weinberger, R. P. (1997). The allocation of epiblast cells to the embryonic heart and other mesodermal lineages: the role of ingression and tissue movement during gastrulation. *Development* 124, 1631–42.
- Tanaka, H., Arakawa, H., Yamaguchi, T., Shiraishi, K., Fukuda, S., Matsui, K., Takei, Y. and Nakamura, Y. (2000). A ribonucleotide reductase gene involved in a p53-dependent cell-cycle checkpoint for DNA damage. *Nature* 404, 42–9.
- Tang, Y., Luo, J., Zhang, W. and Gu, W. (2006). Tip60-dependent acetylation of p53 modulates the decision between cell-cycle arrest and apoptosis. *Mol. Cell* 24, 827–39.
- Tang, Y., Zhong, G., Zhu, L., Liu, X., Shan, Y., Feng, H., Bu, Z., Chen, H. and Wang, C. (2010). Herc5 attenuates influenza A virus by catalyzing ISGylation of viral NS1 protein. *J. Immunol.* 184, 5777–90.
- Taniguchi, H., Jacinto, F. V., Villanueva, A., Fernandez, A. F., Yamamoto, H., Carmona, F. J., Puertas, S., Marquez, V. E., Shinomura, Y., Imai, K., et al. (2012). Silencing of Kruppel-like factor 2 by the histone methyltransferase EZH2 in human cancer. *Oncogene* 31, 1988–94.
- Tanimura, S., Ohtsuka, S., Mitsui, K., Shirouzu, K., Yoshimura, A. and Ohtsubo, M. (1999). MDM2 interacts with MDMX through their RING finger domains. *FEBS Lett.* 447, 5–9.

Bibliografia

- Teng, M. W. L., Swann, J. B., Koebel, C. M., Schreiber, R. D. and Smyth, M. J. (2008). Immune-mediated dormancy: an equilibrium with cancer. *J. Leukoc. Biol.* 84, 988–93.
- Thornberry, N. A. and Lazebnik, Y. (1998). Caspases: enemies within. *Science* 281, 1312–6.
- Uhlmann, J., Wiemann, S. and Ponstingl, H. (1999). DelGef, an RCC1-related protein encoded by a gene on chromosome 11p14 critical for two forms of hereditary deafness. *FEBS Lett.* 460, 153–60.
- Uldrijan, S., Pannekoek, W.-J. and Vousden, K. H. (2007). An essential function of the extreme C-terminus of MDM2 can be provided by MDMX. *EMBO J.* 26, 102–12.
- Valente, L. J., Gray, D. H. D., Michalak, E. M., Pinon-Hofbauer, J., Egle, A., Scott, C. L., Janic, A. and Strasser, A. (2013). p53 efficiently suppresses tumor development in the complete absence of its cell-cycle inhibitory and proapoptotic effectors p21, Puma, and Noxa. *Cell Rep.* 3, 1339–45.
- Van Attikum, H. and Gasser, S. M. (2005). The histone code at DNA breaks: a guide to repair? *Nat. Rev. Mol. Cell Biol.* 6, 757–65.
- Van Attikum, H. and Gasser, S. M. (2009). Crosstalk between histone modifications during the DNA damage response. *Trends Cell Biol.* 19, 207–17.
- Varshavsky, A. (2012). The ubiquitin system, an immense realm. *Annu. Rev. Biochem.* 81, 167–76.
- Vassilev, L. T. (2004). Small-molecule antagonists of p53-MDM2 binding: research tools and potential therapeutics. *Cell Cycle* 3, 419–21.
- Venkitaraman, A. R. (2002). Cancer susceptibility and the functions of BRCA1 and BRCA2. *Cell* 108, 171–82.
- Verdecia, M. A., Joazeiro, C. A. P., Wells, N. J., Ferrer, J.-L., Bowman, M. E., Hunter, T. and Noel, J. P. (2003). Conformational flexibility underlies ubiquitin ligation mediated by the WWP1 HECT domain E3 ligase. *Mol. Cell* 11, 249–59.
- Vinson, R. K. and Hales, B. F. (2002). DNA repair during organogenesis. *Mutat. Res.* 509, 79–91.
- Vousden, K. H. and Lane, D. P. (2007). p53 in health and disease. *Nat. Rev. Mol. Cell Biol.* 8, 275–83.
- Vousden, K. H. and Lu, X. (2002). Live or let die: the cell's response to p53. *Nat. Rev. Cancer* 2, 594–604.
- Vousden, K. H. and Prives, C. (2009). Blinded by the Light: The Growing Complexity of p53. *Cell* 137, 413–31.
- Wade, M., Wang, Y. V and Wahl, G. M. (2010). The p53 orchestra: Mdm2 and Mdmx set the tone. *Trends Cell Biol.* 20, 299–309.
- Wade, M., Li, Y.-C. and Wahl, G. M. (2013). MDM2, MDMX and p53 in oncogenesis and cancer therapy. *Nat. Rev. Cancer* 13, 83–96.
- Walkowicz, M., Ji, Y., Ren, X., Horsthemke, B., Russell, L. B., Johnson, D., Rinchik, E. M., Nicholls, R. D. and Stubbs, L. (1999). Molecular characterization of radiation- and chemically induced mutations associated with neuromuscular tremors, runting, juvenile lethality, and sperm defects in jdf2 mice. *Mamm. Genome* 10, 870–8.
- Wang, R. C. and Levine, B. (2010). Autophagy in cellular growth control. *FEBS Lett.* 584, 1417–26.

- Wang, X., Taplick, J., Geva, N. and Oren, M. (2004). Inhibition of p53 degradation by Mdm2 acetylation. *FEBS Lett.* 561, 195–201.
- Wang, J., Teves, M. E., Shen, X., Nagarkatti-Gude, D. R., Hess, R. A., Henderson, S. C., Strauss, J. F. and Zhang, Z. (2012). Mouse RC/BTB2, a member of the RCC1 superfamily, localizes to spermatid acrosomal vesicles. *PLoS One* 7, e39846.
- Weinberg, R. L., Veprintsev, D. B. and Fersht, A. R. (2004). Cooperative binding of tetrameric p53 to DNA. *J. Mol. Biol.* 341, 1145–59.
- White, E. (2013). Exploiting the bad eating habits of Ras-driven cancers. *Genes Dev.* 27, 2065–71.
- White, D. and Rabago-Smith, M. (2011). Genotype-phenotype associations and human eye color. *J. Hum. Genet.* 56, 5–7.
- Wilkinson, K. D., Urban, M. K. and Haas, A. L. (1980). Ubiquitin is the ATP-dependent proteolysis factor I of rabbit reticulocytes. *J. Biol. Chem.* 255, 7529–32.
- Wong, J. J. Y., Pung, Y. F., Sze, N. S.-K. and Chin, K.-C. (2006). HERC5 is an IFN-induced HECT-type E3 protein ligase that mediates type I IFN-induced ISGylation of protein targets. *Proc. Natl. Acad. Sci. U. S. A.* 103, 10735–40.
- Wu, G. S., Burns, T. F., McDonald, E. R., Jiang, W., Meng, R., Krantz, I. D., Kao, G., Gan, D. D., Zhou, J. Y., Muschel, R., et al. (1997). KILLER/DR5 is a DNA damage-inducible p53-regulated death receptor gene. *Nat. Genet.* 17, 141–3.
- Wu, W., Sato, K., Koike, A., Nishikawa, H., Koizumi, H., Venkitaraman, A. R. and Ohta, T. (2010). HERC2 is an E3 ligase that targets BRCA1 for degradation. *Cancer Res.* 70, 6384–92.
- Xirodimas, D. P., Saville, M. K., Bourdon, J.-C., Hay, R. T. and Lane, D. P. (2004). Mdm2-mediated NEDD8 conjugation of p53 inhibits its transcriptional activity. *Cell* 118, 83–97.
- Xu, J., Zhou, X., Wang, J., Li, Z., Kong, X., Qian, J., Hu, Y. and Fang, J.-Y. (2013). RhoGAPs attenuate cell proliferation by direct interaction with p53 tetramerization domain. *Cell Rep.* 3, 1526–38.
- Yenjerla, M., Panopoulos, A., Reynaud, C., Fotadar, R. and Margolis, R. L. (2013). TD-60 is required for interphase cell cycle progression. *Cell Cycle* 12, 837–41.
- Yoshida, K., Liu, H. and Miki, Y. (2006). Protein kinase C delta regulates Ser46 phosphorylation of p53 tumor suppressor in the apoptotic response to DNA damage. *J. Biol. Chem.* 281, 5734–40.
- You, H., Yamamoto, K. and Mak, T. W. (2006). Regulation of transactivation-independent proapoptotic activity of p53 by FOXO3a. *Proc. Natl. Acad. Sci. U. S. A.* 103, 9051–6.
- Yun, J., Chae, H. D., Choy, H. E., Chung, J., Yoo, H. S., Han, M. H. and Shin, D. Y. (1999). p53 negatively regulates cdc2 transcription via the CCAAT-binding NF-Y transcription factor. *J. Biol. Chem.* 274, 29677–82.
- Zhan, Q., Antinore, M. J., Wang, X. W., Carrier, F., Smith, M. L., Harris, C. C. and Fornace, A. J. (1999). Association with Cdc2 and inhibition of Cdc2/Cyclin B1 kinase activity by the p53-regulated protein Gadd45. *Oncogene* 18, 2892–900.
- Zhang, Y. and Lu, H. (2009). Signaling to p53: ribosomal proteins find their way. *Cancer Cell* 16, 369–77.

Bibliografia

- Zhang, Y. and Xiong, Y. (2001). A p53 amino-terminal nuclear export signal inhibited by DNA damage-induced phosphorylation. *Science* 292, 1910–5.
- Zhang, J., Zhao, D., Park, H. K., Wang, H., Dyer, R. B., Liu, W., Klee, G. G., McNiven, M. A., Tindall, D. J., Molina, J. R., et al. (2010). FAVL elevation in human tumors disrupts Fanconi anemia pathway signaling and promotes genomic instability and tumor growth. *J. Clin. Invest.* 120, 1524–34.
- Zhao, C., Denison, C., Huibregtse, J. M., Gygi, S. and Krug, R. M. (2005). Human ISG15 conjugation targets both IFN-induced and constitutively expressed proteins functioning in diverse cellular pathways. *Proc. Natl. Acad. Sci. U. S. A.* 102, 10200–5.
- Zhao, C., Hsiang, T.-Y., Kuo, R.-L. and Krug, R. M. (2010). ISG15 conjugation system targets the viral NS1 protein in influenza A virus-infected cells. *Proc. Natl. Acad. Sci. U. S. A.* 107, 2253–8.
- Zhao, Y., Brickner, J. R., Majid, M. C. and Mosammamarast, N. (2014). Crosstalk between ubiquitin and other post-translational modifications on chromatin during double-strand break repair. *Trends Cell Biol.*
- Zhou, X. and Münger, K. (2010). Cld7, a candidate tumor suppressor on chromosome 13q14, regulates pathways of DNA damage/repair and apoptosis. *Cancer Res.* 70, 9434–43.
- Zhou, H., Shi, R., Wei, M., Zheng, W.-L., Zhou, J.-Y. and Ma, W.-L. (2013). The expression and clinical significance of HERC4 in breast cancer. *Cancer Cell Int.* 13, 113.
- Zhu, Y., Poyurovsky, M. V., Li, Y., Biderman, L., Stahl, J., Jacq, X. and Prives, C. (2009). Ribosomal protein S7 is both a regulator and a substrate of MDM2. *Mol. Cell* 35, 316–26.