

**DEPARTAMENTO DE FISIOLOGÍA
FACULTAD DE BIOLOGÍA
UNIVERSIDAD DE BARCELONA**

Programa de doctorado de Fisiología
Bienio 2005-2007

**IDENTIFICACIÓN DE FEROMONAS Y PROTEÍNAS
IMPLICADAS EN LA PERCEPCIÓN FEROMONAL DE
LEPIDÓPTEROS PLAGA**

Memoria presentada por Patricia Acín Viu
para optar al título de Doctor
con mención europea

Tesis realizada en el Departamento de Química Biológica y Modelización Molecular del
Instituto de Química Avanzada de Cataluña (IQAC-CSIC) bajo la dirección de la Dra.
Carmen Quero López y la Dra. Glòria Rosell Pellisé

Directoras

Dra. Carmen Quero López
Dept. de Química Biológica y Modelización Molecular
ICAQ
CSIC

Tutora

Dra. Isabel Navarro Álvarez
Dept. de Fisiología
Facultad de Biología
Universidad de Barcelona

Doctoranda

Dra. Glòria Rosell Pellisé
Dept. de Farmacología y Química Terapéutica
Facultad de Farmacia
Universidad de Barcelona

Patricia Acín Viu

8. BIBLIOGRAFÍA

1. Panagiotakopulu, E., Buckland, P., Day, P. and Doumas, C. (1995) Natural insecticides and insect repellents in antiquity: a review of the evidence. *Journal of Archaeological Science*, 22: 705-710.
2. Wheeler, C.M. and Aus, C. (1946) Control of typhus in Italy 1943-1944 by use of DDT. *American Journal of Public Health*, 36: 119-129.
3. Carson, R. (1962) *Silent Spring*. Penguin, London.
4. Flodin, U., Frederiksson, M., Persson, B. and Axelson, O. (1988) Chronic lymphatic leukaemia and engine exhausts, fresh wood and DDT: a case-referent study. *British Journal of Industrial Medicine*, 45: 33-38.
5. Soliman, A.S., Smith, M. A., Cooper, S. P., Ismail, K., Khaled, H., Ismail, S., McPherson, R. S., Seifeldin, I. A. and Bondy, M.L. (1997) Serum organochlorine pesticide levels in patients with colorectal cancer in Egypt. *Archives of Environmental Health*, 52: 409-415.
6. Murray, A. (1985) Acute residual toxicity of a new pyrethroid insecticide, WL85871, to honey-bees. *Bulletin of Environmental Contamination and Toxicology*, 34: 560-564.
7. Hardensen, S. and Wratten, S.D. (2000) Sensitivity of aquatic life stages of *Xanthocnemis zealandica* (Odonata: Zygoptera) to azinphos-methyl and carbaryl. *New Zealand Journal of Marine and Freshwater Research*, 34: 117-123.
8. Mauck, W.L. and Olson, L.E. (1976) Toxicity of natural pyrethrins and five pyretroids to fish. *Archives of Environmental Contamination and Toxicology*, 4: 18-29.
9. Tumlison, J.H. and Teal, P.E.A. (1987) *Relationship of structure and function to biochemistry in insect pheromone systems*. In Prestwich, G.D and Blomquist, G.J.(eds), *Pheromone Biochemistry*. Academic Press. Orlando, Florida, pp 3-26.
10. Butenandt, A., Beckmann, R., Stamm, D. and Hecker, E. (1959) Über den sexual-lockstoff des seidenspinners *Bombyx mori*. Reindarstellung und konstitutionsermittlung. *Zeitschrift für Naturforschung*, 14b: 283-284.
11. Van de Veire, M. and Dirinck, P. (1986) Sex pheromone of the cabbage armyworm, *Mamestra brassicae*: Isolation, identification and field experiments. *Entomologia Experimentalis et Applicata*, 41: 153-155.
12. Bestmann, H.J., Attygale, A. B., Schwarz, J., Vostrowsky, O. and Knauf, W. (1988) Identification of sex pheromone components of *Spodoptera sunia* Gueneé (Lepidoptera:Noctuidae). *Journal of Chemical Ecology*, 14: 683-690.
13. Quero, C., Bau, J., Guerrero, A., Breuer, M., DeLoof, A., Kontzog, H. G. and Camps, F. (2003) Sex pheromone of the oak processionary moth *Thaumatomoea processionea*. Identification and biological activity. *Journal of Agricultural and Food Chemistry*, 51: 2987-2991.

14. Wall, C. (1989) *Monitoring and spray timin*. In Jutsum, A. R. and Gordon, R. F. S. (eds), *Insect Pheromones in Plant Protection*. Wiley, Chichester, UK, pp 39-66.
15. Burkholder, W.E. (1990) Practical use of pheromones and other attractants for stored products insects. In Ridgway, R. L., Silverstein, R. M. and Inscoe, M. N. (eds), *Behavior-Modifying Chemicals for Insect Management*. Marcel Dekker, New York, pp 497-516.
16. Borden, J.H. (1993) *Strategies and tactics for the use of semiochemicals against forest insect pests in North America*, In *Pest Management: Biologically Based Technologies*, American Chemical Society: Washington, D. C, pp 265-279.
17. Lucas, P. and Renou, M. (1989) Responses to pheromone compounds in *Mamestra suasa* (Lepidoptera:Noctuidae) olfactory neurons. *Journal of Insect Physiology*, 35: 837-845.
18. Steinbrecht, R.A. (1987) *Functional morphology of pheromone sensitive sensilla*, In: G. D. Prestwich and G. J. Blomquist (eds) *Pheromone Biochemistry*. Academic Press, Orlando, Florida, pp 353-383.
19. Flower, N.E. (1974) Variation in antennal sensilla of some noctuid moths, a scanning electron microscope study. *New Zealand Journal of Zoology*, 1: 59-66.
20. Grant, A.J., O'Connell, R. J. and Eisner, T. (1989) Pheromone mediated sexual selection in the moth *Utethesia ornatrix*: Olfactory receptor neurons responsive to a male-produced pheromone. *Journal of Insect Behavior*, 2: 371-383.
21. Steinbrecht, R.A., Laue, M., Maida, R. and Ziegelberger, G. (1996) Odorant-binding proteins and their role in the detection of plant odours. *Entomologia Experimentalis et Applicata*, 80: 15-18.
22. Pophof, B. (1997) Olfactory responses recorded from sensilla coeloconica of the silkworm *Bombyx mori*. *Physiological Entomology*, 22: 239-248.
23. Anderson, P., Hallberg, E. and Subchev, M. (2000) Morphology of antennal sensilla auricillica and their detection of plant volatiles in the Herald moth, *Scoliopteryx libatrix* L. (Lepidoptera: Noctuidae). *Arthropod Structure and Development*, 29: 33-41.
24. Chapman, R.F. (1982) Insect chemoreceptors. *Advances in Insect Physiology*, 16: 291-297.
25. Albert, P.J. and Seabrook, W.D. (1973) Morphology and histology of the antenna of the male eastern spruce budworm, *Choristoneura fumiferana* (Clem.) (Lepidoptera:Tortricidae). *Canadian Journal of Zoology*, 51: 443-448.
26. Van Der Pers, J.N.C. and Den Otter, C.J. (1978) Single cell responses from olfactory receptors of small ermine moths (Lepidoptera: Yponomeutidae) to sex attractants. *Journal of Insect Physiology*, 24: 337-343.
27. Shields, V.D.C. and Hildebrand, J.G. (1999) Fine structure of sensilla antennal of the female sphinx moth, *Manduca sexta* (Lepidoptera: Sphingidae). I. Trichoid and basiconic sensilla. *Canadian Journal of Zoology*, 77: 290-301.

28. Vogt, R.G. (2005) Molecular basis of pheromone detection in insects. In Gilbert, L. I., Latro, K., and S. Gill (eds), *Comprehensive Insect Physiology, Biochemistry, Pharmacology and Molecular Biology*. Elsevier, London, pp 753-804.
29. Ziegelberger, G. (1995) Redox-shift of the pheromone-binding protein in the silkworm *Antheraea polyphemus*. *European Journal of Biochemistry*, 232: 706-711.
30. Kaissling, K.E. (2001) Olfactory perireceptor and receptor events in moths: a kinetic model. *Chemical Senses*, 26: 125-150.
31. Damberger, F., Nikonova, L., Horst, R., Peng, G., Leal, W.S., and Würhrich, K. (2000) NMR characterization of a pH-dependent equilibrium between two-folded solution conformations of the pheromone-binding protein from *Bombyx mori*. *Protein Science*, 9: 1038-1041.
32. Horst, R., Damberger, F., Luginbuh , P., Guntert, P., Peng, G., Nikonova, L., Leal, W.S. and Wuthrich, K. (2001) NMR structure reveals intramolecular regulation mechanism for pheromone binding and release. *Proceedings of the National Academy of Sciences USA* 98: 14374-14379.
33. Leal, W.S., Chen, A. M., Ishida, Y., Chiang, V. P., Erickson, M. L., Morgan, T. I. and Tsuruda, J.M. (2005) Kinetics and molecular properties of pheromone binding and release. *Proceedings of the National Academy of Sciences of USA*, 102: 5386-5391.
34. Ishida, Y. and Leal, W.S. (2005) Rapid inactivation of a moth pheromone. *Proceedings of the National Academy of Sciences of USA*, 102: 14075-14079.
35. Leal, W.S. (2005) Pheromone reception. *Topics in Current Chemistry*, 240: 1-36.
36. Birch, M.C. and Haynes, K.F. (1990) *Insect Pheromones*. Barcelona, Oikos-Tau, S. A.
37. Koontz, M.A. and Schneider, D. (1987) Sexual dimorphism in neuronal projections from the antennae of silk moths (*Bombyx mori*, *Antheraea polyphemus*) and the gypsy moth (*Lymantria dispar*). *Cell Tissue Research*, 249: 39-50.
38. Homberg, U., Montague, R. A. and Hildebrand, J.G. (1988) Anatomy of antenno-cerebral pathways in the brain of the sphinx moth *Manduca sexta*. *Cell Tissue Research*, 254: 255-281.
39. Carlsson, M.A., Galizia, C. G. and Hansson, B.S. (2002) Spatial representation of odours in the antennal lobe of the moth *Spodoptera littoralis* (Lepidoptera: Noctuidae). *Chemical Senses*, 27: 231-244.
40. Hansson, B.S. (1995) Olfaction in Lepidoptera. *Experientia*, 51: 1003-1027.
41. Rospars, J.P. and Hildebrand, J.G. (1992) Anatomical identification of glomeruli in the antennal lobes of the male sphinx moth *Manduca sexta*. *Cell Tissue Research*, 270: 205-227.

42. Rospars, J.P. (1983) Invariance and sex-specific variations of the glomerular organization in the antennal lobes of a moth, *Mamestra brassicae*, and a butterfly, *Pieris brassicae*. *Journal of Comparative Neurology*, 220: 80-96.
43. Matsumoto, S.G. and Hildebrand, J.G. (1981) Olfactory mechanisms in the moth *Manduca sexta*: Response characteristics and morphology of central neurons in the antennal lobes. *Proceedings of the Royal Society of Sciences of London*, B213: 249-277.
44. Christensen, T.A. and Hildebrand, J.G. (1987) Male-specific, sex pheromone-selective projection neurons in the antennal lobes of the moth *Manduca sexta*. *Journal of Comparative Physiology*, 160: 553-569.
45. Hansson, B.S., Christensen, T. A. and Hildebrand, J.G. (1991) Functionally distinct subdivisions of the macroglomerular complex in the antennal lobe of the male sphinx moth *Manduca sexta*. *Journal of Comparative Neurology*, 312: 264-278.
46. Ochieng, S.A., Anderson, P. and Hansson, B.S. (1995) AL projection patterns of olfactory receptor neurons involved in sex pheromone detection in *Spodoptera littoralis* (Lepidoptera: Noctuidae). *Tissue Cell*, 27: 221-232.
47. Hansson, B.S. (1997) Antennal lobe projection patterns of pheromone-specific olfactory receptor neurons in moths. In Cardé, R. T. and Minks, A. K. (eds), *Insect Pheromone Research New Directions*. Chapman & Hall: New York, pp 164-183.
48. Hansson, B.S., Ljungberg, H., Hallberg, E. and Löfstedt, C. (1992) Functional specialization of olfactory glomeruli in a moth. *Science*, 256: 1313-1315.
49. Hildebrand, J.G. (1997) *Sensory processing of pheromone signals*. In Cardé, R. T. and Minks, A. K. (eds), *Insect Pheromone Research. New Directions*. Chapman & Hall, New York, pp 111-113.
50. Vogt, R.G., Riddiford, L. M. and Prestwich, G.D. (1985) Kinetic properties of a sex pheromone-degrading enzyme: The sensillar esterase of *Antheraea polyphemus*. *Proceedings of the National Academy of Sciences of USA*, 82: 8827-8831.
51. Hildebrand, J.G. (1996) Olfactory control of behavior in moths: Central processing of odor information and the functional significance of olfactory glomeruli. *Journal of comparative Physiology A*, 178: 5-19.
52. Zhu, J., Kozlov, M. V., Philipp, P., Francke, W. and Löfstedt, C. (1995) Identification of a novel moth sex pheromone in *Eriocrania cicatricella* (Lepidoptera: Eriocraniidae) and its phylogenetic implications. *Journal of Chemical Ecology*, 21: 29-43.
53. Löfstedt, C. and Kozlov, M. (1997) *A phylogenetic analysis of pheromone communication in primitive moths*. In Cardé, R. T. and Minks, A. K. (eds), *Insect Pheromone Research: New Directions*. Chapman & Hall, New York, pp 473-489.
54. Baker, T.C. (1988) *Pheromones and flight behavior*. In Goldsworthy, G.G. and Wheeler, C. (eds), *Insect Flight*. CRC Press, Boca Ratón, Florida.

55. Phelan, P.L. and Baker, T.C. (1987) Evolution of male pheromones in moths: Reproductive isolation through sexual selection? *Science*, 235: 205-207.
56. Phelan, P.L. and Baker, T.C. (1990) Comparative study of courtship in twelve phycitine moths (Lepidoptera:Pyralidae). *Journal of Insect Behavior*, 3: 303-326.
57. Ando, T., Inomata, S.I. and Yamamoto, M. (2004) Lepidopteran sex pheromones. *Topics in Current Chemistry*, 239: 51-96.
58. Scoble, M.J. (1992) *The Lepidoptera: Form, Function and Diversity*. Oxford University Press, Oxford
59. Rauser, C.L. and Rutowski, R.L. (2003) Male-specific structures on the wings of the Gulf Fritillary butterfly, *Agraulis vanillae* (Nymphalidae). *Journal of the Lepidopterists' Society*, 57: 279-283.
60. Pliske, T.E. and Salpenter, M.M. (1971) The structure and development of the hairpencil gland in males of the queen butterfly, *Danaus gilippus berenice*. *Journal of Morphology*, 134: 215-242.
61. Birch, M.C., Poppy, G.M. and Baker, T.C. (1990) Scents and eversible scent structures of male moths. *Annual review of Entomology*, 35: 25-58.
62. Schneider, R.A. (1957) Electrophysiologische untersuchungen von chemo-und mechanorezeptoren der antennen des seidenspinners *Bombyx mori* L. *Zeitschrift für Vergleichende Physiologie*, 40: 8-41.
63. Nagai, T. (1981) Electroantennogram response gradient on the antenna of the european corn borer, *Ostrinia nubilalis*. *Journal of Insect Physiology*, 27: 889-894.
64. Jang, E.B., Light, D. M., Flath, R. A., Nagata, J. T. and Mon, T.R. (1989) Electroantennogram responses of the mediterranean fruit fly, *Ceratitis capitata* to identified volatile constituents from calling males. *Entomolg Experimentalis et Applicata*, 50: 7-19.
65. Evans, K.A. and Bergeron, J. (1994) Behavioral and electrophysiological response of cabbage seed weevils (*Ceutorhynchus assimilis*) to conspecific odor. *Journal of Chemical Ecology*, 20: 979-989.
66. Roelofs, W.L. (1984) Electroantennogram assays: Rapid and convenient screening procedures for pheromones. In Hummel, H. E. and Miller, T. A. (eds), *Techniques in Pheromone Research*. Springer, New York, pp 131-160.
67. Moorhouse, J.E., Yeadon, R., Beevor, P. S. and Nesbitt, B.F. (1969) Method for use in studies of insect chemical communication. *Nature*, 223: 1174-1175.
68. Renou, M. (1991) Sex pheromone reception in the moth, *Mamestra thalassina*. Characterization and distribution of two types of olfactory hairs. *Journal of Insect Physiology*, 37: 617-626.

69. Renou, M., Gadenne, C. and Tauban, D. (1996) Electrophysiology investigations of pheromone-sensitive sensilla in the hybrids between two species *Journal of Insect Physiology*, 42: 267-277.
70. Cossé, A.A., Todd, J. L. and Baker, T.C. (1998) Neurons discovered in male *Helicoverpa zea* antennae that correlate with pheromone-mediated attraction and interspecific antagonism. *Journal of Comparative Physiology A*, 182: 585-594.
71. Baker, T.C. and Cardé, R.T. (1984) *Techniques for behavioral assays*. In Hummel, H. E. and Miller, T. A. (eds), *Techniques in Pheromone Research*. Springer-Verlag, New York, pp 45-73.
72. Klose, J. and Koboltz. (1995) Two-dimensional electrophoresis of proteins: An updated protocol and implications for a functional analysis of the genome. *Electrophoresis*, 16: 1034-1059.
73. Simpson, R.J. (2003) *Preparative two-dimensional gel electrophoresis with immobilized pH gradients*. In *Proteins and Proteomics, a Laboratory Manual*. Cold Spring Harbor Laboratory Press, New York.
74. Bradford, M.M. (1976) A rapid and sensitive method for the quantification of microgram quantities of protein utilizing the principle of protein-dye binding. *Analitical Biochemistry*, 72: 248-254.
75. Lowry, O.H., Rosebrough, N. J., Farr, A. L., and Randall, R. J. (1951) Protein measurement with the folin phenol reagent. *Journal of Biological Chemistry*, 193: 265-275.
76. Görg, A., Postel, W. and Günther, S. (1988) The current state of two-dimensional electrophoresis with immobilized pH gradients. *Electrophoresis*, 9: 531-546.
77. Shevchenko, A., Wilm, M. S., Vorm, O. and Mann, M. (1996) Mass spectrometric sequencing of proteins from silver-stained polyacrilamide gels. *Analytical Chemistry*, 68: 850-858.
78. Chapman, R.F. (1993) *Practical Organic Mass Spectrometry*. Wiley, Chichester, UK.
79. Carrascal, M., Carujo, S., Bachs, O. and Abián, J. (2002) Identification of p21Cip1 binding proteins by gel electrophoresis and capillary liquid chromatography microelectrospray tandem mass spectrometry. *Proteomics*, 2: 455-468.
80. Brady, U.E. and Ganyard, M.C. (1972) Identification of a sex pheromone of the female beet armyworm *Spodoptera exigua*. *Annals of the Entomology Society of America*, 65: 898-899.
81. Persoons, C.J., Van Der Kraan, C., Nooijen, W. J., Ritter, F. J., Voerman, S. and Baker, T.C. (1981) Sex pheromone of the beet armyworm, *Spodoptera exigua*: isolation, identification and preliminary field evaluation. *Entomology Experimentalis et Applicata*, 30:98-99.
82. Tumlinson, J.H., Mitchell, E. R. and Yu, H.S. (1990) Analysis and field evaluation of volatile blends emitted by calling virgin females of beet armyworm moth, *Spodoptera exigua* (HÜBNER). *Journal of Chemical Ecology*, 16: 3411-3423.

83. Mochizuki, F., Shibuya, T., Ihara, T. and Wakamura, S. (1993) Electrophysiological responses of the male antenna to compounds found in the female sex pheromone gland of *Spodoptera exigua* (HÜBNER) (Lepidoptera:Noctuidae). *Applied Entomology and Zoology*, 28: 489-496.
84. Dong, S.L. and Du, J. (2002) Chemical identification and field tests of sex pheromone of beet armyworm *Spodoptera exigua*. *Acta Phytophylacica Sinica*, 29: 19-24.
85. Rogers, C.E. and Underhill, E.W. (1981) A sex attractant for monitoring beet armyworm populations. *The SW Entomology*, 6: 211-214.
86. Mitchell, E.R., Sugie, H., and Tumlinson, J. H. (1983) *Spodoptera exigua*: Capture of feral males in traps baited with blends of pheromone components. *Journal of Chemical Ecology*, 9: 95-104.
87. Cheng, E.Y., Lin, D. F., Kao, C. H., Chen, S. H., Wang, S. S. and Lee, H.C. (1985) Studies on the synthetic pheromone of beet armyworm, *Spodoptera exigua* (Hübner). I. Evaluation of Mitchell's formula in Taiwan. *Journal of Agricultural Research in China* 34: 315-322.
88. Wakamura, S. (1987) Sex pheromone of the beet armyworm, *Spodoptera exigua* (Hübner) (Lepidoptera: Noctuidae): Field attraction of male moths in Japan to (Z,E)-9,12-tetradecadienyl acetate and (Z)-9-tetradecen-1-ol. *Applied Entomology and Zoology*, 22: 348-351.
89. Michell, E.R. and Tumlinson, J.H. (1994) Response of *Spodoptera exigua* and *S. eridania* (Lepidoptera: Noctuidae) males to synthetic pheromone and *S. exigua* females. *The Florida Entomologist*, 77: 237-247.
90. Mochizuki, F. and Shibuya, T. (1991) Antennal single sensillum responses to sex pheromone in male beet armyworm, *Spodoptera exigua* (HÜBNER) (Lepidoptera:Noctuidae). *Applied Entomology and Zoology*, 26: 409-411.
91. Meyers, J. and Miller, T. (1969) Starvation-induced activity in cockroach Malpighian tubules. *Annals of the Entomology Society of America*, 62: 725.
92. Martínez, T. and Camps, F. (1988) Stimulation of sex pheromone production by head extract in *Spodoptera littoralis* at different times of the photoperiod. *Archives of Insect Biochemistry and Physiology*, 9: 211-220.
93. Mas, E., Llòria, J., Quero, C., Camps, F. and Fabriás, G. (2000) Control of the biosynthetic pathway of *Sesamia nonagrioides* sex pheromone by the pheromone biosynthesis activating neuropeptide. *Insect Biochemistry and Molecular Biology*, 30: 455-459.
94. Bau, J. (2000) *Activitat electrofisiològica i comportamental de feromones sexuals de lepidòpters i dels seus inhibidors*. Universitat de Barcelona: Barcelona.
95. Coons, A.H., Creech, H. J. and Jones, R.N. (1941) Immunological properties of an antibody containing a fluorescent group. *Proceedings of the Society of the Experimental Biology and Medicine* 47: 200-202.

96. Dong, S.L. and Du, S.W. (2001) Diel rhythms of calling behavior and sex pheromone production of beet armyworm, *Spodoptera exigua* (Lepidoptera: Noctuidae). *Entomologia sinica*, 8: 89-96.
97. Mitchell, E.R. and Doolittle, R.E. (1976) Sex pheromones of *Spodoptera exigua*, *S. eridania*, and *S. frugiperda*: Bioessay for field activity. *Journal of Economic Entomology*, 69: 324-326.
98. Tumlinson, J.H., Mitchell, E. R. and Sonnet, P.E. (1981) Sex pheromone components of the beet armyworm, *Spodoptera exigua*. *Journal of Environmental Sciences of Health A*, 16: 189-200.
99. Tumlinson, J.H. (1988) Insect pheromone systems. *Comments of Agricultural Food Chemistry*, 1: 115-146.
100. Teal, P.E.A. and Tumlinson, J.H. (1986) Terminal steps in pheromone biosynthesis by *Heliothis virescens* and *H. zea*. *Journal of Chemical Ecology*, 12: 353-366.
101. Teal, P.E.A. and Tumlinson, J.H. (1987) The role of alcohols in pheromone biosynthesis by two noctuid moths that use acetate pheromone components. *Archives of Insect Biochemistry and Physiology*, 4: 261-269.
102. Teal, P.E.A., Heath, R. R., Tumlinson, J. H. and McLaughlin, J.R. (1981) Identification of a sex pheromone of *Heliothis subflexa* (Gn.) and field trapping studies using different blends of components. *Journal of Chemical Ecology*, 7: 1011-1022.
103. Mochizuki, F., takai, M., Shibuya, T., and Wakamura, S. (1994) Field traps responses of male *Spodoptera exigua* (HÜBNER) (Lepidoptera: Noctuidae) to (Z,E)-9,12-tetradecadien-1-ol and (Z)-9-tetradecenyl acetate. *Applied Entomology and Zoology*, 29: 436-438.
104. Baker, T.C. (1989) Sex pheromone in the Lepidoptera: New research progress. *Experientia*, 45: 248-262.
105. Linn, C.E.J., Campbell, M. G. and Roelofs, W.L. (1986) Male moth sensitivity to multicomponent pheromone: critical role of female-released blend in determining the functional role of components and active space of the pheromone. *Journal of Chemical Ecology*, 12: 659-668.
106. Linn, C.E.J., Campbell, M. G. and Roelofs, W.L. (1987) Pheromone components and active spaces: what do moths smell and where do they smell it? *Science*, 237: 650-652.
107. Tumlinson, J.H., Mitchell, E. R., Teal, P.E.A., Heath, R.R. and Mengelkoch, L.J. (1986) Sex pheromone of fall armyworm, *Spodoptera frugiperda* (J.E. Smith). Identification of components critical to attraction in the field. *Journal of Chemical Ecology*, 12: 1909-1926.
108. Dickens, J.C. and Payne, T.L. (1977) Bark beetle olfaction: Pheromone receptor system in *Dendroctonus frontalis*. *Journal of Insect Physiology*, 23: 481-489.
109. Visser, J.H. (1979) Electroantennogram responses of the Colorado beetle, *Leptinotarsa decemlineata*, to plant volatiles. *Entomologia Experimentalis et Applicata*, 25: 86-97.

110. Campion, D.G. (1975) Sex pheromones and their use for control of insects of the genus *Spodoptera*. *Mededelingen Faculteit landbouwwetenschappen Rijksuniversiteit Gent*, 40: 283-292.
111. Mitchell, E.R., Sugie, H. and Tumlinson, J.H. (1983) Rubber septa as dispensers for the fall armyworm sex attractant. *Journal of Environmental Sciences of Health A*, 18: 463-470.
112. Jung, C.R., Park, Y. J. and Boo, K.S. (2003) Optimal sex pheromone composition for monitoring *Spodoptera exigua* (Lepidoptera: Noctuidae) in Korea. *Journal of Asia-Pacific Entomology* 6: 175-182.
113. Baker, T.C., Meyer, W. and Roelofs, W.L. (1981) Sex pheromone dosage and blend specificity of responses by oriental fruit moth males. *Entomology Experimentalis et Applicata*, 30: 269-279.
114. Linn, C.E.J. and Roelofs, W.L. (1981) Modification of sex pheromone blend discrimination in male oriental fruit moths by pre-exposure to (E)-8-dodecetyl acetate. *Physiological Entomology*, 4: 421-429.
115. Dickens, J.C., Visser, J. H. and Van Der Pers, J.N.C. (1993) Detection and deactivation of pheromone and plant odor components by the beet armyworm, *Spodoptera exigua* (HÜBNER) (Lepidoptera: Noctuidae). *Journal of Insect Physiology*, 39: 503-516.
116. Mochizuki, F., Sugi, N., and Shibuya, T. (1992) Pheromone sensilla of the beet armyworm, *Spodoptera exigua* (HÜBNER) (Lepidoptera: Noctuidae). *Applied Entomology and Zoology*, 27: 547-556.
117. Vosshall, L.B. (2001) The molecular logic of olfaction in *Drosophila*. *Chemical Senses*, 26: 207-213.
118. Klein, U. (1987) Sensillum-lymph proteins from antennal olfactory hairs of the moth *Antheraea polyphemus* (Saturniidae). *Insect Biochemistry*, 17: 1193-1204.
119. Steinbrecht, R.A., Ozaki, M., and Ziegelberger, G. (1992) Immunocytochemical localization of pheromone binding protein in moth antennae. *Cell Tissue Research*, 270: 287-302.
120. Leal W. S., N.L., Peng G. (1999) Disulfide structure of the pheromone binding protein from the silkworm moth, *Bombyx mori*. *FEBS Letters*, 468: 85-90.
121. Briand, L., Nespolous, C., Huet, J. C. and Pernollet, J.C. (2001) Disulfide pairing and secondary structure of ASP1, an olfactory-binding protein from honeybee (*Apis mellifera* L.). *Journal of Peptide Research*, 58: 540-545.
122. Briand, L., Nespolous, C., Huet, J. C., Takahashi, M. and Pernollet, J.C. (2001) Ligand binding and physico-chemical properties of ASP2, a recombinant odorant-binding protein from honeybee (*Apis mellifera* L.). *European Journal of Biochemistry*, 268: 752-760.
123. Breer, H., Krieger, J. and Raming, K. (1990) A novel class of binding proteins in the antennae of the silk moth *Antheraea pernyi*. *Insect Biochemistry*, 20: 735-774.

124. Vogt, R.G., Prestwich, G. D. and Lerner, M.R. (1991) Odorant-binding-protein subfamilies associate with distinct classes of olfactory receptor neurons in insects. *Journal of Neurobiology*, 22: 74–84.
125. Vogt, R.G. and Riddiford, L.M. (1981) Pheromone binding and inactivation by moth antennae. *Nature*, 293: 161-163.
126. Sandler, B.H., Nikonova, L., Leal, W. S., and Clardy, J. (2000) Sexual attraction in the silkworm moth: Structure of the pheromone- binding protein- bombykol complex. *Chemical Biology*, 7: 143-151.
127. Vogt, R.G., Kohne, A. C., Dubnau, J. T. and Prestwich, G.D. (1989) Expression of pheromone-binding proteins during antennal development in the gypsy moth *Lymantria dispar*. *Journal of Neurosciences*, 9: 3332 – 3346.
128. Vogt, R.G. (1987) *The molecular basis of pheromone reception: its influence on behavior*. In Prestwich, G.D. and Blomquist, G.J. (eds), *Pheromone Biochemistry*. Academic Press, New York, pp 385-431.
129. Callahan, F.E., Vogt, R. G., Tuckerc, M. L., Dickens, J. C. and Mattoo, A.K. (2000) High level expression of "male specific" pheromone binding proteins (PBPs) in the antennae of female noctuid moths. *Insect Biochemistry and Molecular Biology*, 30: 507-514.
130. Maïbèche-Coisné, M., Jacquin-Joly, E, Francois, M.C., and Nagnan-Le Meillour, P. (1998) Molecular cloning of two pheromone binding proteins in the cabbage armyworm *Mamestra brassicae*. *Insect Biochem. Mol. Biol.*, 28: 815-818.
131. Krieger, J.a.B., H. and (1999) Olfactory reception in invertebrates. *Science*, 286: 720–723.
132. Cusson, M. and McNeil, J.N. (1989) Involvement of juvenile hormone in the regulation of pheromone release activities in a moth. *Science*, 243: 210-212.
133. Ljunberg, H., Anderson, P., and Hansson, B.S. (1993) Physiology and morphology of pheomone-specific sensilla on the antennae of male and female *Spodoptera littoralis* (Lepidoptera: Noctuidae). *Journal of Insect Physiology*, 39: 253-260.
134. Picimbon, J.F. and Gadenne, C. (2002) Evolution of noctuid pheromone binding proteins: identification of PBP in the black cutworm moth, *Agrotis ipsilon*. *Insect Biochemistry and Molecular Biology*, 32: 839-846.
135. Steinbrecht, R.A., Laue, M. and Ziegelberger, G. (1995) Immunolocalization of pheromone-binding protein and general odorant-binding protein in olfactory sensilla of the silk moths *Antheraea* and *Bombyx*. *Cell Tissue Research*, 282: 203-217.
136. Vogt, R.G., Rogers, M. E., Franco, M. D. and Sun, M. (2002) A comparative study of odorant binding protein genes: differential expression of the PBP1-GOBP2 gene cluster in *Manduca sexta* (Lepidoptera) and the organization of OBP genes in *Drosophila melanogaster* (Diptera). *Journal of Experimental Biology*, 205: 719-744.

137. Laue, M. and Steinbrecht, R.A. (1997) Topochemistry of moth olfactory sensilla. *International Journal of Insect Morphology and Embryology*, 26: 217–228.
138. Maida, R., Mameli, M., Krieger, J., Breer, H., Ziegelberger, G. and Steinbrecht, R.A. (1999) Complex expression pattern of odorant-binding proteins in *Bombyx mori*. In Elsner, N. and Eysel, U. (eds), *Gottingen Neurobiology Report*. Thieme, Stuttgart.
139. Pelosi, P. (1996) Perireceptor events in olfaction. *Journal of Neurobiology* 30: 3-19.
140. Krieger, J. and Breer, H. (1999) Olfactory reception in invertebrates. *Science*, 286: 720–723.
141. Steinbrecht, R.A. (1996) Are odorant-binding proteins involved in odorant discrimination? . *Chemical Senses* 21: 718–725.
142. Kaissling, E.K. (1986) Chemo-electrical transduction in insect olfactory receptors. *Annual review of Neurosciences*, 9: 121-145.
143. Kaissling, E.K. (1996) Peripheral mechanisms of pheromone reception in moths. *Chemical Senses*, 21: 257-268.
144. Du, G. and Prestwich, G.D. (1995) Protein structure encodes te ligand binding specificity in pheromone binding proteins. *Biochemistry*, 34: 8726-8732.
145. Kasang, G. (1971) Bombykol reception and metabolism on the antennae of the silkworm *Bombyx mori*. In G., Ohloff and A. F., Thomas (eds), *Gustation and Olfaction*. Academic Press, London, pp 245–250.
146. Ferkovich, S.M., Mayer, M. S. and Rutter, R.R. (1973) Conversion of the sex pheromone of the cabbage looper. *Nature*, 242: 53–55.
147. Ferkovich, S.M., Mayer, M. S. and Rutter, R.R. (1973) Sex pheromone of the cabbage looper: reactions with antennal proteins in vitro. *Journal of Insect Physiology*, 19: 2231–2243.
148. Quero, C., Camps, F. and Guerrero, A. (1995) Behavior of processionary males (*Thaumetopoea pityocampa*) induced by sex pheromone and analogs in a wind tunnel. *Journal of Chemical Ecology*, 21: 1957–1969.
149. Liotta, A., kohn, E. C. and Petricoin, E.F. (2001) Clinical Proteomics: Personalized Molecular Medicine. *Journal of American Medical Association*, 286: 2211-2214.
150. Kowcun A., H.N. and E., P. (2001) Olfaction in the gypsy moth, *Lymantria dispar*: effect of pH, ionic strength, and reductants on pheromone transport by pheromonebinding proteins. *Journal of Biological Chemistry*, 276: 44770–44776.
151. Olsson, I., Larsson, K., Palmgren, R. and Bjellqvist, B. (2002) Organic disulfides as a means to generate streak-free two-dimensional maps with narrow range basic immobilized pH gradient strips as first dimension. *Proteomics*, 2: 1630-1632.

152. Martinez, T. and Camps, F. (1988) Stimulation of sex pheromone production by head extract in *Spodoptera littoralis* at different times of the photoperiod. *Archives of Insect Biochemistry and Physiology*, 9: 211-220.
153. Ausubel, F.M., Brent, R., Kingston, R. E., Moore, D. D., Seidman, J. G., Smith, J. A. and Struhl, K. (1995) *Short Protocols in Molecular Biology*. John Wiley & Sons, Inc.
154. Quero, C., Colomé, N., Prieto, M. R., Carrascal, M., Posada, M., Gelpí, E. and Abián, J. (2004) Determination of protein markers in human serum: analysis of protein expression in toxic oil syndrome studies. *Proteomics*, 4: 303-315.
155. de Santis, F., François, M.-C., Merlin, C., Pelletier, J., Maibèche-Coisné, M., Conti, E. and Jacquin-Joly, E. (2006) Molecular cloning and in situ expression patterns of two new pheromone-binding proteins from the corn stemborer *Sesamia nonagrioides*. *Journal of Chemical Ecology*, 32: 1703-1717.
156. Mazomenos, B. (1989) Sex components of corn stalk borer *Sesamia nonagrioides* (LEF.): Isolation, Identification, and Field tests. *Journal of Chemical Ecology*, 15: 1241-1247.
157. Xiu, W.M. and Dong, S.L. (2007) Molecular characterization of two pheromone binding proteins and quantitative analysis of their expression in the beet armyworm, *Spodoptera exigua* Hübner. *Journal of Chemical Ecology*, 33: 947-961.
158. Ljungberg, H., Anderson, P. and Hansson, B.S. (1993) Physiology and morphology of pheromone-specific sensilla of male and female *Spodoptera littoralis* (Lepidoptera: Noctuidae). *Journal of Insect Physiology*, 39: 253-260.
159. Mazomenos, B. (1989) Sex pheromone components of corn stalk borer, *Sesamia nonagrioides* (Lef.). Partial chemical purification and its biological activity under laboratory conditions. *Journal of Chemical Ecology*, 6: 565-572.
160. Martínez, T., Fabriás, G., and Camps, F. (1990) Sex pheromone biosynthetic pathway in *Spodoptera littoralis* and its activation by a neurohormone. *Journal of Biological Chemistry*, 265: 1381-1387.
161. Pelosi, P., Zhou, J. J., Ban, L. P. and Calvello, M. (2006) Soluble proteins in insect chemical communication. *Cellular and Molecular Life Sciences*, 63: 1658-1676.
162. Abraham, D., Löfstedt, C. and Picimbon, J.F. (2005) Molecular characterization and evolution of pheromone-binding protein genes in *Agrotis* moths. *Insect Biochemistry and Molecular Biology*, 35: 1100-1111.
163. Maibèche-Coisné, M., Jacquin-Joly, E., François, M.C. and Nagnan-Le Meillour, P. (1998) Molecular cloning of two pheromone binding proteins in the cabbage armyworm *Mamestra brassicae*. *Insect Biochemistry and Molecular Biology*, 28: 815-818.
164. Prestwich, G.D., Du, G. and Laforest, S. (1995) How is pheromone specificity encoded in proteins? *Chemical Senses*, 20: 461-469.

165. Mohanty, S., Zubkov, S. and Gronenborn, A.M. (2004) The Solution NMR Structure of *Antheraea polyphemus* PBP Provides New Insight into Pheromone Recognition by Pheromone-binding Proteins. *Journal of Molecular Biology*, 337: 443-451.
166. Teal, P.E.A., Heath, R.R., Dueben, B.D., Coffelt, J.A. and Vick, K.W. (1995) Production and release of (Z,E)-9,12-tetradecadienal by sex pheromone glands of females of *Plodia interpunctella* (Lepidoptera: Pyralidae). *Journal of Chemical Ecology*, 21: 787-799.
167. Bette S., B.H. and J., K. (2002) Probing a pheromone binding protein of the silkworm *Antheraea polyphemus* by endogenous tryptophan fluorescence. *Insect Biochemistry and Molecular Biology*, 32: 241-246.
168. Maïbèche- Coisné, M., Sobrio, F., Delaunay, T., Lettere, M., Dubroca, J., Jaquin-Joly, E., and Nagnan-Le Meillour, P. (1997) Pheromone binding proteins of the moth *Mamestra brassicae*: specificity of ligand binding. *Insect Biochemistry and Molecular Biology*, 27: 213-221.
169. Lautenschlager, C., Leal, W. S., and Clardy, J. (2007) *Bombyx mori* pheromone - binding protein binding non-pheromone ligands: implications for pheromone recognition. *Structure*, 15: 1148-1154.
170. Leal, W.S., Chen, A. M., Erickson, M. L. (2005) Selective and pH-dependent binding of a moth pheromone to a pheromone- binding protein. *Journal of Chemical Ecology*, 31: 2493-2499.
171. Raming, K., Krieger, J. and Breer, H. (1990) Primary structure of a pheromone binding protein from *Antheraea pernyi*: homologies with other ligand-carrying proteins. *Journal of Comparative Physiology*, 160: 503-509.
172. Maida, R., Steinbrecht, A., Ziegelberger, G. and Pelosi, P. (1993) The pheromone binding protein of *Bombyx mori*. Purification, characterization and immunocytochemical localization. *Insect Biochemistry and Molecular Biology*, 23: 243-253.
173. Györgyi, T.K., Roby-Shemkovitz, A. J. and Lerner, M.R. (1988) Characterization and cDNA cloning of the pheromone-binding protein from the tobacco hornworm, *Manduca sexta*: a tissue-specific developmentally regulated protein. *Proceedings of the National Academy of Sciences of USA*, 85: 9851-9855.
174. Zhang, S., Maida, R. and Steinbrecht, R.A. (2001) Immunolocalization of odorant binding proteins in Noctuid moths (Insecta, Lepidoptera). *Chemical Senses*, 26: 885- 896.
175. Schneider, D., Schulz, S., Priesner, E., Ziesmann, J. and Francke, W. (1998) Autodetection and chemistry of female and male pheromone in both sexes of the tiger moth *Panaxia quadripunctaria*. *Journal of Comparative Physiology A*, 182: 153-161.
176. Todd, J.L. and Baker, T.C. (1993) Response of single antennal neurons of female cabbage loopers to behaviorally active attractants. *Naturwissenschaften*, 80: 183-186.
177. Kalinova, B., Hoskovec, M., Liblikas, I., Unelius, C. R. and Hansson, B.S. (2001) Detection of sex pheromone components in *Manduca sexta* (L.). *Chemical Senses*, 26: 1175-1186.

178. Newcomb, R.D., Sirey, T. M., Rassam, M. and Greenwood, D.R. (2002) Pheromone binding proteins of *Epiphyas postvittana* (Lepidoptera: Tortricidae) are encoded at a single locus. *Insect Biochemistry and Molecular Biology*, 32: 1543-1554.
179. Konstantopoulou, M.A., Pratsinis, H., Kletsas, D. and Mazomenos, B.E. (2006) Pheromone-binding protein and general odorant binding protein of *Sesamia nonagrioides*: Sex- and diel-dependent expression. *Entomologia Experimentalis et Applicata*, 119: 129-136.
180. Acín, P., Carrascal, M., Abián, J., Guerrero, A. and Quero, C. (2008) Expression of differential antennal proteins in males and females of an important crop pest, *Sesamia nonagrioides*. *Insect Biochemistry and Molecular Biology*, doi:10.1016/j.ibmb.2008.09.007.
181. Feng, L. and Prestwich, G.D. (1997) Expression and characterization of a lepidopteran general odorant binding protein. *Insect Biochemistry and Molecular Biology*, 27: 405-412.
182. Dickens J. C., S., J. W., and Light, D. M. (1993) Green leaf volatiles enhance sex attractant pheromone of the tobacco budworm, *Heliothis virescens* (Fabricius) (Lepidoptera: Noctuidae). *Chemoecology*, 4: 175-177.
183. Light, D.M., Flath, R. A., Buttery, R. G., Zalom, F. G., Rice, R. E., Dickens, J. C. and Jang, E.B. (1993) Host-plant green-leaf volatiles synergize the synthetic sex pheromones of the corn earworm and codling moth (Lepidoptera). *Chemoecology*, 4: 145-152.
184. Dickens, J.C., Smith, J. W. and Light, D.M. (1993) Green leaf volatiles enhance sex attractant pheromone of the tobacco budworm, *Heliothis virescens* (Fabricius) (Lepidoptera: Noctuidae). *Chemoecology*, 4: 175-177.
185. Renwick, J.A.A. (1989) Chemical ecology of oviposition in phytophagous insects. *Experientia*, 45: 223-228.
186. Merlin, C., Rosell, G., Carot-Sans, G., François, M.C., Bozzolan, F., Pelletier, J., Jacquin-Joly, E., Guerrero, A. and Maibèche-Coisne, M. (2007) Antennal esterase cDNAs from two pest moths, *Spodoptera littoralis* and *Sesamia nonagrioides*, potentially involved in odourant degradation. *Insect Molecular Biology*, 16: 73-81.
187. Sarto i Monteys, V. and Aguilar, L. (2005) The Castniid Palm Borer, *Paysandisia archon* (Burmeister, 1880) (Lep. Castniidae), in Europe: Comparative biology, pest status and possible control methods. *Nachrichten entomologischen Vereins Apollo* 26:61-94.
188. Aguilar, L., Miller, J.Y. and Sarto i Monteys, V. (2001) A new lepidopteran family for the European fauna. *SHILAP Revista lepidopterología*, 29: 86-87.
189. Sarto i Monteys, V. and Aguilar, L. (2001) *Paysandisia archon* (Burmeister, 1880), Castniidae, also in France. *SHILAP Revista lepidopterología*, 29: 280.
190. Espinosa, P., Di Miuccio, P. and Russo, G. (2003) *Paysandisia archon*, una minaccia per le nostre palme. *Informatore Agrario*, 7: 61.
191. Patton, S. and Perry, M. (2002) *Paysandisia archon* -The First British Record. *Atropos*, 18: 28.

192. Drescher, J. and Jaubert, R. (2003) *Paysandisia archon* continue sa progression. *PHM Revue Horticole*, 445: 49-51.
193. Millar, J.E. and Haynes, K.F. (1998) *Methods in Chemical Ecology: Chemical Methods*. Kluwer Academic Publisher, Massachusetts.
194. Andersson, J., Borg-Karlsson, A.K. and Wiklund, C. (2003) Functional significance of citral as an aphrodisiac in the butterfly *Pieris napi*. Behavioural experiments and EAG tests. *Journal of Chemical Ecology*, 29: 1489-1499.
195. Vane-Wright, R.I. and Boppré, M. (1993) Visual and chemical signalling in butterflies: functional and phylogenetic perspectives. *Philosophical Transactions of the Royal Society of London B*, 340: 197-205.
196. Den Otter, C.J., Schuil, H.A. and Sander-Van Oosten, A. (1978) Reception of host-plant odours and female sex pheromone in *Adoxophyes orana* (Lepidoptera: Tortricidae): Electrophysiology and morphology. *Entomologia Experimentalis et Applicata*, 24: 370-378.
197. Kafka, W.A. (1987) Similarity of reaction spectra and odor discrimination:single receptor cell recordings in *Antheraea polyphemus* (Saturniidae). *Journal of Comparative Physiology*, 161: 867-880.
198. Behan, M. and Schoonhoven, L.M. (1978) Chemoreception of an oviposition deterrent associated with eggs in *Pieris brassicae*. *Entomologia Experimentalis et Applicata*, 24: 163-179.
199. Honda, K. (1982) Exocrine substances of the white cabbage butterfly, *Pieris rapae* crucivora (Lepidoptera:Pieridae). *Applied Entomology and Zoology*, 17: 325-331.
200. Rebouças, L.M.C., Do Caraciolo, M.S.B., Sant'Ana, A.E.G., Pickett, J.A., Wadhams, L.J. and Pow, E.M. (1999) Composição química da glândula abdominal da fêmea da mariposa *Castnia licus* (Drury) (Lepidoptera:Castniidae): possíveis feromônios e precursores. *Química Nova*, 22: 645-648.
201. Sturaro, A., Parvoli, G. and Doretti, L. (1992) Artifacts produced by Porapak Q sorbent tubes on solvent desorption. *Chromatographia*, 33: 53-57.
202. Leal, W.S., Kuwahara, Y., Suzuki, T. and Kurosa, K. (1989) The alarm pheromone of the mite *Suidasia medanensis* OUDEMANS, 1924 (Acariformes, Suisasiidae). *Agricultural Chemistry*, 53: 2703-2709.
203. Nishida, R. (1995) *Oviposition stimulants of swallowtail butterflies*, In *Swallowtail Butterflies: Their Ecology and Evolutionary Biology*, Scientific Publishers, Gainsville, FL. pp 17-26.
204. Nishida, R. (2005) Chemosensory basis of host recognition in butterflies- multi-component system of oviposition stimulants and deterrents. *Chemical Senses*, 30: i293-i294.
205. Schlüter, K., Juckusch, B.M. and Rothkegel, M. (1997) Profilins as regulators of actin dynamics. *Biochimica et Biophysica Acta-Molecular Cell Research*, 1359: 97-109.

206. Kuwahara, Y. (1979) Scent Scale Substances of male *Pieris melete* MÉNÈTRIÈS (Pieridae:Lepidoptera). *Applied Entomology and Zoology*, 14: 350-355.
207. Hillier, N.K. and Vickers, N.J. (2004) The role of heliothine hairpencil compounds in female *Heliothis virescens* (Lepidoptera:Noctuidae) behavior and mate acceptance. *Chemical Senses*, 29: 499-511.
208. Wedell, N. (2005) Female receptivity in butterflies and moths. *Journal of Experimental Biology*, 208: 3433-3440.
209. Nieberding, C.M., de Vos, H., Schneider, M.V., Lassance, J.M., Estramil; N., Andersson, J., Bang, J., Hedenstrom, E., Löfstedt,C. and Brakefield, P.M. (2008) The male sex pheromone of the butterfly *Bicyclus anynana*: Towards an evolutionary analysis. *PlosOne*, 3: 1-12.
210. Schneider, D. and Seibt, U. (1969) Sex pheromone of the queen butterfly: Electroantennogram responses. *Science*, 164: 1173-1174.
211. Grant, G.G. (1971) Electroantennogram responses to the scent brush secretions of several male moths. *Annual Entomology Society of America*, 64: 1428-1431.
212. Burger, B.V., Nell, A. E., Smit, D., Spies, H. S. C., Mackenroth, W. M., Groche, D. and Atkinson, P.R. (1993) Constituents of wing gland and abdominal hairpencil secretions of male african sugarcane borer, *Eldana saccharine* Walker (Lepidoptera: Pyralidae). *Journal of Chemical Ecology*, 19: 2255-2277.
213. Lecomte, C., Thibout, E., Pierre, D. and Auger, J. (1998) Transfer, perception, and activity of male pheromone of *Acrolepiopsis assectella* with special reference to conspecific male sexual inhibition. *Journal of Chemical Ecology*, 24: 655-671.
214. Hendricks, D.E. (1976) Tobacco budworm: Disruption of courtship behavior with a component of the synthetic sex pheromone. *Environmental Entomology*, 5: 978-980.
215. Hirai, K., Shorey, H. H. and Gaston, L.K. (1978) Competition among courting male moths: Male-to-male inhibitory pheromone. *Science*, 202: 644-645.
216. Kemp, D. and Wiklund, C. (2001) Fighting without weaponry: a review of male-male contest competition in butterflies. *Behavioral Ecology and Sociobiology*, 49: 429-442.
217. Romero, F. (1998) Comportamiento, oviposición y distribución de *Duboisvalia simulans* (Boisduval) en Venezuela (Lepidoptera: Castniidae). *Boletín de Entomología Venezolana*, 13: 91-92.
218. Schulz, S., Beccaloni, G, Brown Jr, k.S., Boppré, M., Freitas, A.V.L, Ockenfels, P. and Trigo, J.R. (2004) Semiochemicals derived from pyrrolizidine alkaloids in male ithomiine butterflies (Lepidoptera:Nymphalidae:Ithomiinae). *Biochemical Systematics and Ecology*, 32: 699-713.
219. Eisner, T. and Meinwald, J. (1987) *Alkaloid-derived pheromones and sexual selection in Lepidoptera*. In Prestwich, G.D. and Blomquist, G.J. (eds), *Pheromone Biochemistry*. Academic, Orlando, FL. pp 251-269.

220. Trigo, J.R., Barata, L.E.S. and Brown, K.S. (1994) Stereochemical inversion of pyrrolizidine alkaloids by *Mechanitis polymnia* (Lepidoptera:Nymphalidae:Ithomiinae):specificity and evolutionary significance. *Journal of Chemical Ecology*, 20: 2883-2899.
221. Zagatti, P., Kunesch, G., Ramiandrasoa, F., Malosse, C., Hall, D.R., Lester, R. and Nesbitt, B.F. (1987) Sex pheromone of rice moth, *Corcyra cephalonica* STANTON I. Identification of male pheromone. *Journal of Chemical Ecology*, 13: 1561-1573.
222. Arn, H., Toth, M. and Priesner, E. (2002) *List of sex pheromones of Lepidoptera and related attractants*, Paris: 012 BSROP publ.
223. Francke, W., Karalius, V., Plass, E., Lehmann, L., Dos Santos, A., Buda, V., Borg-Karlson, A.K. and Mozuraitis, R. (2004) New type of Sesiidae sex pheromone identified from the hornet moth *Sesia apiformis*. *Journal of Chemical Ecology*, 30: 805-817.
224. Heath, R.R., Landolt, P.J., Leppla, N.C. and Dueben, B.D. (1988) Identification of a male produced pheromone of *Anticarsia gemmatalis* (Hübner) (Lepidoptera: Noctuidae) attractive to conspecific males. *Journal of Chemical Ecology*, 14: 1121-1130.
225. Jacobson, M., Adler, V.E. and Baumhover, A.H. (1984) A male tobacco budworm pheromone inhibitory to courtship. *Journal of Environmental Sciences of Health A*, 19: 469-476.
226. Ollivier, L. and Frérot, B. (2006) *Paysandisia archon* Vous avez dit phéromone? *Phytoma*, 594: 30-32.
227. Andersson, J., Borg-Karlson, A.K., Vongvanich, N. and Wiklund, C. (2007) Male sex pheromone release and female mate choice in a butterfly. *The Journal of Experimental Biology*, 210: 964-970.
228. Renwick, J.A.A. and Radke, C.D. (1987) Chemical stimulants and deterrents regulating acceptance or rejection of crucifers by cabbage butterflies. *Journal of Chemical Ecology*, 13: 1771-1776.
229. Nishida, R. and Fumiaki, H. (1989) Oviposition stimulants of an Aristolochiaceae-feeding swallowtail butterfly, *Atrophaneura alcinous*. *Journal of Chemical Ecology*, 15: 2565-2575.
230. Roessingh, P., Städler, E., Schönl, R. and Feeny, P. (1991) Tarsal contact chemoreceptors of the black swallowtail butterfly, *Papilio polyxenes*: responses to phytochemicals from host and non-host plants. *Physiological Entomology*, 16: 485-495.
231. Tsuchihara, K., Ueno, K., Yamanaka, A., Isono, K., Endo, K., Nishida, R., Yoshihara, K. and Tokunaga, F. (2000) A putative binding protein for lipophilic substances related to butterfly oviposition. *FEBS Letters*, 478: 299-303.