

2009

TESIS DOCTORAL

UTILIDAD DE LA RESONANCIA MAGNÉTICA
EN PACIENTES CON FIBRILACIÓN
AURICULAR TRIBUTARIOS DE
TRATAMIENTO CON ABLACIÓN
PERCUTÁNEA DE LAS VENAS PULMONARES

ROSARIO JESÚS PEREA PALAZÓN

UNIVERSITAT DE BARCELONA
Línea de investigación de biopatología cardiovascular
Dirigida por el Prof. Josep Brugada Terradellas y por la Dra.
Teresa M^a de Caralt Robira

1. BIBLIOGRAFÍA

Referencias bibliográficas

- (1) Bellet S. Clinical disorders of the heart beat. 3rd ed. Philadelphia, PA: Lea & Febiger; 1971.
- (2) Prytowsky EN, Katz AM. Atrial fibrillation. In: Philadelphia, Lipincott-Raven, editors. Textbook of Cardiovascular Medicine. 1998. p. 1661.
- (3) Brugada R, Tapscott T, Czernuszewicz GZ, Marian AJ, Iglesias A, Mont L, et al. Identification of a genetic locus for familial atrial fibrillation. *N Engl J Med* 1997 Mar 27;336(13):905-11.
- (4) Levy S. Epidemiology and classification of atrial fibrillation. *J Cardiovasc Electrophysiol* 1998 Aug;9:S78-S82.
- (5) Sopher SM, Camm AJ. Therapy for atrial fibrillation: control of the ventricular response and prevention of recurrence. *Coron Artery Dis* 1995 Feb;6:106-14.
- (6) Fuster V, Ryden LE, Cannom DS, Crijns HJ, Curtis AB, Ellenbogen KA, et al. ACC/AHA/ESC 2006 guidelines for the management of patients with atrial fibrillation: full text: a report of the American College of Cardiology/American Heart Association Task Force on practice guidelines and the European Society of Cardiology Committee for Practice Guidelines (Writing Committee to Revise the 2001 guidelines for the management of patients with atrial fibrillation) developed in collaboration with the European Heart Rhythm Association and the Heart Rhythm Society. *Europace* 2006 Sep;8(9):651-745.
- (7) Fuster V, Ryden LE, Cannom DS, Crijns HJ, Curtis AB, Ellenbogen KA, et al. ACC/AHA/ESC 2006 Guidelines for the Management of Patients with Atrial Fibrillation: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines and the European Society of Cardiology Committee for Practice Guidelines (Writing Committee to Revise the 2001 Guidelines for the Management of Patients With Atrial Fibrillation): developed in collaboration with the European Heart Rhythm Association and the Heart Rhythm Society. *Circulation* 2006 Aug 15;114(7):e257-e354.
- (8) Kopecky SL, Gersh BJ, McGoon MD, Whisnant JP, Holmes DR, Jr., Ilstrup DM, et al. The natural history of lone atrial fibrillation. A population-based study over three decades. *N Engl J Med* 1987 Sep 10;317(11):669-74.
- (9) Chen YJ, Chen SA, Tai CT, Wen ZC, Feng AN, Ding YA, et al. Role of atrial electrophysiology and autonomic nervous system in patients with supraventricular tachycardia and paroxysmal atrial fibrillation. *J Am Coll Cardiol* 1998 Sep;32(3):732-8.
- (10) Frustaci A, Chimenti C, Bellocchi F, Morgante E, Russo MA, Maseri A. Histological substrate of atrial biopsies in patients with lone atrial fibrillation. *Circulation* 1997 Aug 19;96:1180-4.
- (11) Kopecky SL, Gersh BJ, McGoon MD, Chu CP, Ilstrup DM, Chesebro JH, et al. Lone atrial fibrillation in elderly persons: a marker for cardiovascular risk. *Arch Intern Med* 1999 May 24;159(10):1118-22.
- (12) Jouven X, Desnos M, Guerot C, Ducimetiere P. Idiopathic atrial fibrillation as a risk factor for mortality. The Paris Prospective Study I. *Eur Heart J* 1999 Jun;20(12):896-9.
- (13) Falk RH. Atrial fibrillation. *N Engl J Med* 2001 Apr 5;344(14):1067-78.

- (14) Pappone C, Rosanio S. Evolution of non-pharmacological curative therapy for atrial fibrillation. Where do we stand today? *Int J Cardiol* 2003 Apr;88:135-42.
- (15) Feinberg WM, Blackshear JL, Laupacis A, Kronmal R, Hart RG. Prevalence, age distribution, and gender of patients with atrial fibrillation. Analysis and implications. *Arch Intern Med* 1995 Mar 13;155:469-73.
- (16) Furberg CD, Psaty BM, Manolio TA, Gardin JM, Smith VE, Rautaharju PM. Prevalence of atrial fibrillation in elderly subjects (the Cardiovascular Health Study). *Am J Cardiol* 1994 Aug 1;74:236-41.
- (17) Wolf PA, Abbott RD, Kannel WB. Atrial fibrillation as an independent risk factor for stroke: the Framingham Study. *Stroke* 1991 Aug;22:983-8.
- (18) Maisel WH, Stevenson LW. Atrial fibrillation in heart failure: epidemiology, pathophysiology, and rationale for therapy. *Am J Cardiol* 2003 Mar 20;91:2D-8D.
- (19) Benjamin EJ, Wolf PA, D'Agostino RB, Silbershatz H, Kannel WB, Levy D. Impact of atrial fibrillation on the risk of death: the Framingham Heart Study. *Circulation* 1998 Sep 8;98:946-52.
- (20) Dries DL, Exner DV, Gersh BJ, Domanski MJ, Waclawiw MA, Stevenson LW. Atrial fibrillation is associated with an increased risk for mortality and heart failure progression in patients with asymptomatic and symptomatic left ventricular systolic dysfunction: a retrospective analysis of the SOLVD trials. *Studies of Left Ventricular Dysfunction*. *J Am Coll Cardiol* 1998 Sep;32(3):695-703.
- (21) Redfield MM, Kay GN, Jenkins LS, Mianulli M, Jensen DN, Ellenbogen KA. Tachycardia-related cardiomyopathy: a common cause of ventricular dysfunction in patients with atrial fibrillation referred for atrioventricular ablation. *Mayo Clin Proc* 2000 Aug;75:790-5.
- (22) Le Heuzey JY, Paziaud O, Piot O, Said MA, Copie X, Lavergne T, et al. Cost of care distribution in atrial fibrillation patients: the COCAF study. *Am Heart J* 2004 Jan;147(1):121-6.
- (23) Konings KT, Kirchhof CJ, Smeets JR, Wellens HJ, Penn OC, Allessie MA. High-density mapping of electrically induced atrial fibrillation in humans. *Circulation* 1994 Apr;89(4):1665-80.
- (24) SCHERF D, ROMANO FJ, TERRANOVA R. Experimental studies on auricular flutter and auricular fibrillation. *Am Heart J* 1948 Aug;36(2):241-51.
- (25) Jais P, Haissaguerre M, Shah DC, Chouairi S, Gencel L, Hocini M, et al. A focal source of atrial fibrillation treated by discrete radiofrequency ablation. *Circulation* 1997 Feb 4;95(3):572-6.
- (26) Chen SA, Tai CT, Yu WC, Chen YJ, Tsai CF, Hsieh MH, et al. Right atrial focal atrial fibrillation: electrophysiologic characteristics and radiofrequency catheter ablation. *J Cardiovasc Electrophysiol* 1999 Mar;10(3):328-35.
- (27) Haissaguerre M, Jais P, Shah DC, Takahashi A, Hocini M, Quiniou G, et al. Spontaneous initiation of atrial fibrillation by ectopic beats originating in the pulmonary veins. *N Engl J Med* 1998 Sep 3;339:659-66.
- (28) Hsu LF, Jais P, Keane D, Wharton JM, Deisenhofer I, Hocini M, et al. Atrial fibrillation originating from persistent left superior vena cava. *Circulation* 2004 Feb 24;109(7):828-32.
- (29) Lin WS, Tai CT, Hsieh MH, Tsai CF, Lin YK, Tsao HM, et al. Catheter ablation of paroxysmal atrial fibrillation initiated by non-pulmonary vein ectopy. *Circulation* 2003 Jul 1;107(25):3176-83.

- (30) Schmitt C, Ndrepepa G, Weber S, Schmieder S, Weyerbrock S, Schneider M, et al. Biatrial multisite mapping of atrial premature complexes triggering onset of atrial fibrillation. *Am J Cardiol* 2002 Jun 15;89(12):1381-7.
- (31) Schwartzman D, Bazaz R, Nosbisch J. Common left pulmonary vein: a consistent source of arrhythmogenic atrial ectopy. *J Cardiovasc Electrophysiol* 2004 May;15(5):560-6.
- (32) Jais P, Hocini M, Macle L, Choi KJ, Deisenhofer I, Weerasooriya R, et al. Distinctive electrophysiological properties of pulmonary veins in patients with atrial fibrillation. *Circulation* 2002 Nov 5;106(19):2479-85.
- (33) Shah D, Haissaguerre M, Jais P, Hocini M. Nonpulmonary vein foci: do they exist? *Pacing Clin Electrophysiol* 2003 Jul;26(7 Pt 2):1631-5.
- (34) Kumagai K, Nakashima H, Saku K. The HMG-CoA reductase inhibitor atorvastatin prevents atrial fibrillation by inhibiting inflammation in a canine sterile pericarditis model. *Cardiovasc Res* 2004 Apr 1;62(1):105-11.
- (35) MOE GK, ABILDSKOV JA. Atrial fibrillation as a self-sustaining arrhythmia independent of focal discharge. *Am Heart J* 1959 Jul;58(1):59-70.
- (36) Mandapati R, Skanes A, Chen J, Berenfeld O, Jalife J. Stable microreentrant sources as a mechanism of atrial fibrillation in the isolated sheep heart. *Circulation* 2000 Jan 18;101(2):194-9.
- (37) Lazar S, Dixit S, Marchlinski FE, Callans DJ, Gerstenfeld EP. Presence of left-to-right atrial frequency gradient in paroxysmal but not persistent atrial fibrillation in humans. *Circulation* 2004 Nov 16;110(20):3181-6.
- (38) Sanders P, Berenfeld O, Hocini M, Jais P, Vaidyanathan R, Hsu LF, et al. Spectral analysis identifies sites of high-frequency activity maintaining atrial fibrillation in humans. *Circulation* 2005 Aug 9;112(6):789-97.
- (39) Nattel S. Electrophysiologic remodeling: are ion channels static players or dynamic movers? *J Cardiovasc Electrophysiol* 1999 Nov;10:1553-6.
- (40) Goette A, Honeycutt C, Langberg JJ. Electrical remodeling in atrial fibrillation. Time course and mechanisms. *Circulation* 1996 Dec 1;94:2968-74.
- (41) Ausma J, Litjens N, Lenders MH, Duimel H, Mast F, Wouters L, et al. Time course of atrial fibrillation-induced cellular structural remodeling in atria of the goat. *J Mol Cell Cardiol* 2001 Dec;33:2083-94.
- (42) Li D, Fareh S, Leung TK, Nattel S. Promotion of atrial fibrillation by heart failure in dogs: atrial remodeling of a different sort. *Circulation* 1999 Jul 6;100:87-95.
- (43) Hoit BD, Shao Y, Gabel M, Walsh RA. Left atrial mechanical and biochemical adaptation to pacing induced heart failure. *Cardiovasc Res* 1995 Apr;29:469-74.
- (44) Hoit BD, Shao Y, Gabel M. Left atrial systolic and diastolic function accompanying chronic rapid pacing-induced atrial failure. *Am J Physiol* 1998 Jul;275(1 Pt 2):H183-H189.
- (45) Khan A, Moe GW, Nili N, Rezaei E, Eskandarian M, Butany J, et al. The cardiac atria are chambers of active remodeling and dynamic collagen turnover during evolving heart failure. *J Am Coll Cardiol* 2004 Jan 7;43:68-76.

- (46) Allessie M, Ausma J, Schotten U. Electrical, contractile and structural remodeling during atrial fibrillation. *Cardiovasc Res* 2002 May;54(2):230-46.
- (47) Bailey GW, Braniff BA, Hancock EW, Cohn KE. Relation of left atrial pathology to atrial fibrillation in mitral valvular disease. *Ann Intern Med* 1968 Jul;69(1):13-20.
- (48) Aime-Sempe C, Folliguet T, Rucker-Martin C, Krajewska M, Krajewska S, Heimburger M, et al. Myocardial cell death in fibrillating and dilated human right atria. *J Am Coll Cardiol* 1999 Nov 1;34(5):1577-86.
- (49) Polontchouk L, Haefliger JA, Ebelt B, Schaefer T, Stuhlmann D, Mehlhorn U, et al. Effects of chronic atrial fibrillation on gap junction distribution in human and rat atria. *J Am Coll Cardiol* 2001 Sep;38(3):883-91.
- (50) Everett TH, Li H, Mangrum JM, McRury ID, Mitchell MA, Redick JA, et al. Electrical, morphological, and ultrastructural remodeling and reverse remodeling in a canine model of chronic atrial fibrillation. *Circulation* 2000 Sep 19;102:1454-60.
- (51) Schoonderwoerd BA, Ausma J, Crijns HJ, Van Veldhuisen DJ, Blaauw EH, Van Gelder IC. Atrial ultrastructural changes during experimental atrial tachycardia depend on high ventricular rate. *J Cardiovasc Electrophysiol* 2004 Oct;15(10):1167-74.
- (52) Ausma J, Wijffels M, Thone F, Wouters L, Allessie M, Borgers M. Structural changes of atrial myocardium due to sustained atrial fibrillation in the goat. *Circulation* 1997 Nov 4;96:3157-63.
- (53) Ricard P, Levy S, Trigano J, Paganelli F, Daoud E, Man KC, et al. Prospective assessment of the minimum energy needed for external electrical cardioversion of atrial fibrillation. *Am J Cardiol* 1997 Mar 15;79(6):815-6.
- (54) Wijffels MC, Kirchhof CJ, Dorland R, Allessie MA. Atrial fibrillation begets atrial fibrillation. A study in awake chronically instrumented goats. *Circulation* 1995 Oct 1;92:1954-68.
- (55) Solti F, Vecsey T, Kekesi V, Juhasz-Nagy A. The effect of atrial dilatation on the genesis of atrial arrhythmias. *Cardiovasc Res* 1989 Oct;23:882-6.
- (56) Morillo CA, Klein GJ, Jones DL, Guiraudon CM. Chronic rapid atrial pacing. Structural, functional, and electrophysiological characteristics of a new model of sustained atrial fibrillation. *Circulation* 1995 Mar 1;91:1588-95.
- (57) Shi Y, Ducharme A, Li D, Gaspo R, Nattel S, Tardif JC. Remodeling of atrial dimensions and emptying function in canine models of atrial fibrillation. *Cardiovasc Res* 2001 Nov;52:217-25.
- (58) Kumagai K, Nakashima H, Urata H, Gondo N, Arakawa K, Saku K. Effects of angiotensin II type 1 receptor antagonist on electrical and structural remodeling in atrial fibrillation. *J Am Coll Cardiol* 2003 Jun 18;41:2197-204.
- (59) Raitt MH, Kusumoto W, Giraud G, McAnulty JH. Reversal of electrical remodeling after cardioversion of persistent atrial fibrillation. *J Cardiovasc Electrophysiol* 2004 May;15:507-12.
- (60) Cha YM, Dzeja PP, Shen WK, Jahangir A, Hart CY, Terzic A, et al. Failing atrial myocardium: energetic deficits accompany structural remodeling and electrical instability. *Am J Physiol Heart Circ Physiol* 2003 Apr;284:H1313-H1320.

- (61) Barger PM, Kelly DP. Fatty acid utilization in the hypertrophied and failing heart: molecular regulatory mechanisms. *Am J Med Sci* 1999 Jul;318:36-42.
- (62) Ausma J, Coumans WA, Duimel H, Van der Vusse GJ, Allessie MA, Borgers M. Atrial high energy phosphate content and mitochondrial enzyme activity during chronic atrial fibrillation. *Cardiovasc Res* 2000 Sep;47:788-96.
- (63) Allard MF, Schonekess BO, Henning SL, English DR, Lopaschuk GD. Contribution of oxidative metabolism and glycolysis to ATP production in hypertrophied hearts. *Am J Physiol* 1994 Aug;267:H742-H750.
- (64) Stanley WC, Lopaschuk GD, Hall JL, McCormack JG. Regulation of myocardial carbohydrate metabolism under normal and ischaemic conditions. Potential for pharmacological interventions. *Cardiovasc Res* 1997 Feb;33:243-57.
- (65) Dietz JR. Mechanisms of atrial natriuretic peptide secretion from the atrium. *Cardiovasc Res* 2005 Oct 1;68:8-17.
- (66) Tsiofis C, Stougiannos P, Taxiarchou E, Skiadas I, Chatzis D, Thomopoulos C, et al. The interplay between haemodynamic load, brain natriuretic peptide and left atrial size in the early stages of essential hypertension. *J Hypertens* 2006 May;24:965-72.
- (67) Hanna N, Cardin S, Leung TK, Nattel S. Differences in atrial versus ventricular remodeling in dogs with ventricular tachypacing-induced congestive heart failure. *Cardiovasc Res* 2004 Aug 1;63:236-44.
- (68) Miyauchi Y, Zhou S, Okuyama Y, Miyauchi M, Hayashi H, Hamabe A, et al. Altered atrial electrical restitution and heterogeneous sympathetic hyperinnervation in hearts with chronic left ventricular myocardial infarction: implications for atrial fibrillation. *Circulation* 2003 Jul 22;108:360-6.
- (69) Goette A, Staack T, Rocken C, Arndt M, Geller JC, Huth C, et al. Increased expression of extracellular signal-regulated kinase and angiotensin-converting enzyme in human atria during atrial fibrillation. *J Am Coll Cardiol* 2000 May;35:1669-77.
- (70) Levin ER, Gardner DG, Samson WK. Natriuretic peptides. *N Engl J Med* 1998 Jul 30;339:321-8.
- (71) Stockand JD, Meszaros JG. Aldosterone stimulates proliferation of cardiac fibroblasts by activating Ki-RasA and MAPK1/2 signaling. *Am J Physiol Heart Circ Physiol* 2003 Jan;284:H176-H184.
- (72) Rude MK, Duhaney TA, Kuster GM, Judge S, Heo J, Colucci WS, et al. Aldosterone stimulates matrix metalloproteinases and reactive oxygen species in adult rat ventricular cardiomyocytes. *Hypertension* 2005 Sep;46:555-61.
- (73) Nakashima H, Kumagai K, Urata H, Gondo N, Ideishi M, Arakawa K. Angiotensin II antagonist prevents electrical remodeling in atrial fibrillation. *Circulation* 2000 Jun 6;101:2612-7.
- (74) Mesiskli T, Flevari PG, Livanis EG, Bofilis E, Theodorakis GN, Kremastinos DT, et al. cGMP levels following ANP challenge are markers of subsequent successful reversion of lone atrial fibrillation to sinus rhythm. *Pacing Clin Electrophysiol* 1998 Jan;21:168-71.
- (75) Madrid AH, Bueno MG, Rebollo JM, Marin I, Pena G, Bernal E, et al. Use of irbesartan to maintain sinus rhythm in patients with long-lasting persistent atrial fibrillation: a prospective and randomized study. *Circulation* 2002 Jul 16;106:331-6.

- (76) Healey JS, Baranchuk A, Crystal E, Morillo CA, Garfinkle M, Yusuf S, et al. Prevention of atrial fibrillation with angiotensin-converting enzyme inhibitors and angiotensin receptor blockers: a meta-analysis. *J Am Coll Cardiol* 2005 Jun 7;45(11):1832-9.
- (77) Thai H, Castellano L, Juneman E, Phan H, Do R, Gaballa MA, et al. Pretreatment with angiotensin receptor blockade prevents left ventricular dysfunction and blunts left ventricular remodeling associated with acute myocardial infarction. *Circulation* 2006 Oct 31;114(18):1933-9.
- (78) Wong GC, Marcotte F, Rudski LG. Impact of chronic lisinopril therapy on left atrial volume versus dimension in chronic organic mitral regurgitation. *Can J Cardiol* 2006 Feb;22:125-9.
- (79) Anne W, Willems R, Roskams T, Sergeant P, Herijgers P, Holemans P, et al. Matrix metalloproteinases and atrial remodeling in patients with mitral valve disease and atrial fibrillation. *Cardiovasc Res* 2005 Sep 1;67:655-66.
- (80) Anne W, Willems R, Holemans P, Beckers F, Roskams T, Lenaerts I, et al. Self-terminating AF depends on electrical remodeling while persistent AF depends on additional structural changes in a rapid atrially paced sheep model. *J Mol Cell Cardiol* 2007 Aug;43:148-58.
- (81) Boldt A, Wetzel U, Weigl J, Garbade J, Lauschke J, Hindricks G, et al. Expression of angiotensin II receptors in human left and right atrial tissue in atrial fibrillation with and without underlying mitral valve disease. *J Am Coll Cardiol* 2003 Nov 19;42(10):1785-92.
- (82) Madrid AH, Peng J, Zamora J, Marin I, Bernal E, Escobar C, et al. The role of angiotensin receptor blockers and/or angiotensin converting enzyme inhibitors in the prevention of atrial fibrillation in patients with cardiovascular diseases: meta-analysis of randomized controlled clinical trials. *Pacing Clin Electrophysiol* 2004 Oct;27:1405-10.
- (83) Shroff SC, Ryu K, Martovitz NL, Hoit BD, Stambler BS. Selective aldosterone blockade suppresses atrial tachyarrhythmias in heart failure. *J Cardiovasc Electrophysiol* 2006 May;17:534-41.
- (84) Psychari SN, Apostolou TS, Sinos L, Hamodraka E, Liakos G, Kremastinos DT. Relation of elevated C-reactive protein and interleukin-6 levels to left atrial size and duration of episodes in patients with atrial fibrillation. *Am J Cardiol* 2005 Mar 15;95:764-7.
- (85) Watanabe T, Takeishi Y, Hirono O, Itoh M, Matsui M, Nakamura K, et al. C-Reactive protein elevation predicts the occurrence of atrial structural remodeling in patients with paroxysmal atrial fibrillation. *Heart Vessels* 2005 Mar;20:45-9.
- (86) Chung MK, Martin DO, Sprecher D, Wazni O, Kanderian A, Carnes CA, et al. C-reactive protein elevation in patients with atrial arrhythmias: inflammatory mechanisms and persistence of atrial fibrillation. *Circulation* 2001 Dec 11;104:2886-91.
- (87) Pappone C, Santinelli V, Manguso F, Vicedomini G, Gugliotta F, Augello G, et al. Pulmonary vein denervation enhances long-term benefit after circumferential ablation for paroxysmal atrial fibrillation. *Circulation* 2004 Jan 27;109(3):327-34.
- (88) Schauerte P, Scherlag BJ, Pitha J, Scherlag MA, Reynolds D, Lazzara R, et al. Catheter ablation of cardiac autonomic nerves for prevention of vagal atrial fibrillation. *Circulation* 2000 Nov 28;102(22):2774-80.
- (89) Elvan A, Pride HP, Eble JN, Zipes DP. Radiofrequency catheter ablation of the atria reduces inducibility and duration of atrial fibrillation in dogs. *Circulation* 1995 Apr 15;91(8):2235-44.

- (90) Bettoni M, Zimmermann M. Autonomic tone variations before the onset of paroxysmal atrial fibrillation. *Circulation* 2002 Jun 11;105(23):2753-9.
- (91) Lombardi F, Tarricone D, Tundo F, Colombo F, Belletti S, Fiorentini C. Autonomic nervous system and paroxysmal atrial fibrillation: a study based on the analysis of RR interval changes before, during and after paroxysmal atrial fibrillation. *Eur Heart J* 2004 Jul;25(14):1242-8.
- (92) Gami AS, Pressman G, Caples SM, Kanagala R, Gard JJ, Davison DE, et al. Association of atrial fibrillation and obstructive sleep apnea. *Circulation* 2004 Jul 27;110(4):364-7.
- (93) Tsang TS, Gersh BJ, Appleton CP, Tajik AJ, Barnes ME, Bailey KR, et al. Left ventricular diastolic dysfunction as a predictor of the first diagnosed nonvalvular atrial fibrillation in 840 elderly men and women. *J Am Coll Cardiol* 2002 Nov 6;40(9):1636-44.
- (94) Kitzman DW, Edwards WD. Age-related changes in the anatomy of the normal human heart. *J Gerontol* 1990 Mar;45:M33-M39.
- (95) Redfield MM, Jacobsen SJ, Borlaug BA, Rodeheffer RJ, Kass DA. Age- and gender-related ventricular-vascular stiffening: a community-based study. *Circulation* 2005 Oct 11;112:2254-62.
- (96) Thomas L, Levett K, Boyd A, Leung DY, Schiller NB, Ross DL. Compensatory changes in atrial volumes with normal aging: is atrial enlargement inevitable? *J Am Coll Cardiol* 2002 Nov 6;40(9):1630-5.
- (97) Anyukhovsky EP, Sosunov EA, Chandra P, Rosen TS, Boyden PA, Danilo P, Jr., et al. Age-associated changes in electrophysiologic remodeling: a potential contributor to initiation of atrial fibrillation. *Cardiovasc Res* 2005 May 1;66:353-63.
- (98) Kistler PM, Sanders P, Fynn SP, Stevenson IH, Spence SJ, Vohra JK, et al. Electrophysiologic and electroanatomic changes in the human atrium associated with age. *J Am Coll Cardiol* 2004 Jul 7;44:109-16.
- (99) Little WC, Zile MR, Kitzman DW, Hundley WG, O'Brien TX, Degroot RC. The effect of alagebrium chloride (ALT-711), a novel glucose cross-link breaker, in the treatment of elderly patients with diastolic heart failure. *J Card Fail* 2005 Apr;11:191-5.
- (100) Peterson LR, Waggoner AD, de las FL, Schechtman KB, McGill JB, Gropler RJ, et al. Alterations in left ventricular structure and function in type-1 diabetics: a focus on left atrial contribution to function. *J Am Soc Echocardiogr* 2006 Jun;19:749-55.
- (101) Sasson Z, Rasooly Y, Gupta R, Rasooly I. Left atrial enlargement in healthy obese: prevalence and relation to left ventricular mass and diastolic function. *Can J Cardiol* 1996 Mar;12:257-63.
- (102) Tsang TS, Barnes ME, Gersh BJ, Bailey KR, Seward JB. Left atrial volume as a morphophysiologic expression of left ventricular diastolic dysfunction and relation to cardiovascular risk burden. *Am J Cardiol* 2002 Dec 15;90:1284-9.
- (103) Upshaw CB, Jr. Hemodynamic changes after cardioversion of chronic atrial fibrillation. *Arch Intern Med* 1997 May 26;157(10):1070-6.
- (104) van den Berg MP, Tuinenburg AE, Van Veldhuisen DJ, de Kam PJ, Crijns HJ. Cardioversion of atrial fibrillation in the setting of mild to moderate heart failure. *Int J Cardiol* 1998 Jan 5;63(1):63-70.

- (105) Kochiadakis GE, Skalidis EI, Kalebubas MD, Igoumenidis NE, Chrysostomakis SI, Kanoupakis EM, et al. Effect of acute atrial fibrillation on phasic coronary blood flow pattern and flow reserve in humans. *Eur Heart J* 2002 May;23(9):734-41.
- (106) Sanfilippo AJ, Abascal VM, Sheehan M, Oertel LB, Harrigan P, Hughes RA, et al. Atrial enlargement as a consequence of atrial fibrillation. A prospective echocardiographic study. *Circulation* 1990 Sep;82:792-7.
- (107) Gosselink AT, Crijns HJ, Hamer HP, Hillege H, Lie Kl. Changes in left and right atrial size after cardioversion of atrial fibrillation: role of mitral valve disease. *J Am Coll Cardiol* 1993 Nov 15;22(6):1666-72.
- (108) Manning WJ, Silverman DI, Katz SE, Riley MF, Come PC, Doherty RM, et al. Impaired left atrial mechanical function after cardioversion: relation to the duration of atrial fibrillation. *J Am Coll Cardiol* 1994 Jun;23(7):1535-40.
- (109) Mitusch R, Garbe M, Schmucker G, Schwabe K, Stierle U, Sheikhzadeh A. Relation of left atrial appendage function to the duration and reversibility of nonvalvular atrial fibrillation. *Am J Cardiol* 1995 May 1;75(14):944-7.
- (110) Grogan M, Smith HC, Gersh BJ, Wood DL. Left ventricular dysfunction due to atrial fibrillation in patients initially believed to have idiopathic dilated cardiomyopathy. *Am J Cardiol* 1992 Jun 15;69(19):1570-3.
- (111) Kiern JR, Sacrez A, Facello A, Arbogast R, Bareiss P, Roul G, et al. Increase in radionuclide left ventricular ejection fraction after cardioversion of chronic atrial fibrillation in idiopathic dilated cardiomyopathy. *Eur Heart J* 1992 Sep;13(9):1290-5.
- (112) Packer DL, Bardy GH, Worley SJ, Smith MS, Cobb FR, Coleman RE, et al. Tachycardia-induced cardiomyopathy: a reversible form of left ventricular dysfunction. *Am J Cardiol* 1986 Mar 1;57(8):563-70.
- (113) Shinbane JS, Wood MA, Jensen DN, Ellenbogen KA, Fitzpatrick AP, Scheinman MM. Tachycardia-induced cardiomyopathy: a review of animal models and clinical studies. *J Am Coll Cardiol* 1997 Mar 15;29(4):709-15.
- (114) EVANS W, SWANN P. Lone auricular fibrillation. *Br Heart J* 1954 Apr;16(2):189-94.
- (115) Levy S, Maarek M, Coumel P, Guize L, Lekieffre J, Medvedowsky JL, et al. Characterization of different subsets of atrial fibrillation in general practice in France: the ALFA study. The College of French Cardiologists. *Circulation* 1999 Jun 15;99(23):3028-35.
- (116) Coromilas J. Obesity and atrial fibrillation: is one epidemic feeding the other? *JAMA* 2004 Nov 24;292(20):2519-20.
- (117) Frost L, Hune LJ, Vestergaard P. Overweight and obesity as risk factors for atrial fibrillation or flutter: the Danish Diet, Cancer, and Health Study. *Am J Med* 2005 May;118(5):489-95.
- (118) Wang TJ, Parise H, Levy D, D'Agostino RB, Sr., Wolf PA, Vasan RS, et al. Obesity and the risk of new-onset atrial fibrillation. *JAMA* 2004 Nov 24;292(20):2471-7.
- (119) Fox CS, Parise H, D'Agostino RB, Sr., Lloyd-Jones DM, Vasan RS, Wang TJ, et al. Parental atrial fibrillation as a risk factor for atrial fibrillation in offspring. *JAMA* 2004 Jun 16;291(23):2851-5.

- (120) Darbar D, Herron KJ, Ballew JD, Jahangir A, Gersh BJ, Shen WK, et al. Familial atrial fibrillation is a genetically heterogeneous disorder. *J Am Coll Cardiol* 2003 Jun;41(12):2185-92.
- (121) Fioranelli M, Piccoli M, Mileto GM, Sgreccia F, Azzolini P, Risa MP, et al. Analysis of heart rate variability five minutes before the onset of paroxysmal atrial fibrillation. *Pacing Clin Electrophysiol* 1999 May;22(5):743-9.
- (122) Herweg B, Dalal P, Nagy B, Schweitzer P. Power spectral analysis of heart period variability of preceding sinus rhythm before initiation of paroxysmal atrial fibrillation. *Am J Cardiol* 1998 Oct 1;82(7):869-74.
- (123) Maisel WH. Autonomic modulation preceding the onset of atrial fibrillation. *J Am Coll Cardiol* 2003 Oct 1;42(7):1269-70.
- (124) Fetsch T, Bauer P, Engberding R, Koch HP, Lukl J, Meinertz T, et al. Prevention of atrial fibrillation after cardioversion: results of the PAFAC trial. *Eur Heart J* 2004 Aug;25(16):1385-94.
- (125) Israel CW, Gronefeld G, Ehrlich JR, Li YG, Hohnloser SH. Long-term risk of recurrent atrial fibrillation as documented by an implantable monitoring device: implications for optimal patient care. *J Am Coll Cardiol* 2004 Jan 7;43(1):47-52.
- (126) Page RL, Wilkinson WE, Clair WK, McCarthy EA, Pritchett EL. Asymptomatic arrhythmias in patients with symptomatic paroxysmal atrial fibrillation and paroxysmal supraventricular tachycardia. *Circulation* 1994 Jan;89(1):224-7.
- (127) Page RL, Tilsch TW, Connolly SJ, Schnell DJ, Marcello SR, Wilkinson WE, et al. Asymptomatic or "silent" atrial fibrillation: frequency in untreated patients and patients receiving azimilide. *Circulation* 2003 Mar 4;107(8):1141-5.
- (128) A randomized, placebo-controlled trial of propafenone in the prophylaxis of paroxysmal supraventricular tachycardia and paroxysmal atrial fibrillation. UK Propafenone PSVT Study Group. *Circulation* 1995 Nov 1;92(9):2550-7.
- (129) Anderson JL, Gilbert EM, Alpert BL, Henthorn RW, Waldo AL, Bhandari AK, et al. Prevention of symptomatic recurrences of paroxysmal atrial fibrillation in patients initially tolerating antiarrhythmic therapy. A multicenter, double-blind, crossover study of flecainide and placebo with transtelephonic monitoring. Flecainide Supraventricular Tachycardia Study Group. *Circulation* 1989 Dec;80(6):1557-70.
- (130) Lee SH, Chen SA, Tai CT, Chiang CE, Wen ZC, Chen YJ, et al. Comparisons of oral propafenone and sotalol as an initial treatment in patients with symptomatic paroxysmal atrial fibrillation. *Am J Cardiol* 1997 Apr 1;79(7):905-8.
- (131) Clemo HF, Wood MA, Gilligan DM, Ellenbogen KA. Intravenous amiodarone for acute heart rate control in the critically ill patient with atrial tachyarrhythmias. *Am J Cardiol* 1998 Mar 1;81(5):594-8.
- (132) Juul-Moller S, Edvardsson N, Rehnqvist-Ahlberg N. Sotalol versus quinidine for the maintenance of sinus rhythm after direct current conversion of atrial fibrillation. *Circulation* 1990 Dec;82(6):1932-9.
- (133) Kochiadakis GE, Igioumenidis NE, Marketou ME, Kaleboubas MD, Simantirakis EN, Vardas PE. Low dose amiodarone and sotalol in the treatment of recurrent, symptomatic atrial fibrillation: a comparative, placebo controlled study. *Heart* 2000 Sep;84(3):251-7.

- (134) Roy D, Talajic M, Dorian P, Connolly S, Eisenberg MJ, Green M, et al. Amiodarone to prevent recurrence of atrial fibrillation. Canadian Trial of Atrial Fibrillation Investigators. *N Engl J Med* 2000 Mar 30;342(13):913-20.
- (135) LOWN B, MARASINGHAM R, NEUMAN J. New method for terminating cardiac arrhythmias. Use of synchronized capacitor discharge. *JAMA* 1962 Nov 3;182:548-55.
- (136) Levy S, Ricard P, Lau CP, Lok NS, Camm AJ, Murgatroyd FD, et al. Multicenter low energy transvenous atrial defibrillation (XAD) trial results in different subsets of atrial fibrillation. *J Am Coll Cardiol* 1997 Mar 15;29(4):750-5.
- (137) Levy S, Ricard P, Gueunoun M, Yapo F, Trigano J, Mansouri C, et al. Low-energy cardioversion of spontaneous atrial fibrillation. Immediate and long-term results. *Circulation* 1997 Jul 1;96(1):253-9.
- (138) RABBINO MD, LIKOFF W, DREIFUS LS. COMPLICATIONS AND LIMITATIONS OF DIRECT-CURRENT COUNTERSHOCK. *JAMA* 1964 Nov 2;190:417-20.
- (139) Fitzpatrick AP, Kourouyan HD, Siu A, Lee RJ, Lesh MD, Epstein LM, et al. Quality of life and outcomes after radiofrequency His-bundle catheter ablation and permanent pacemaker implantation: impact of treatment in paroxysmal and established atrial fibrillation. *Am Heart J* 1996 Mar;131(3):499-507.
- (140) Saksena S, Prakash A, Hill M, Krol RB, Munsif AN, Mathew PP, et al. Prevention of recurrent atrial fibrillation with chronic dual-site right atrial pacing. *J Am Coll Cardiol* 1996 Sep;28(3):687-94.
- (141) Saksena S, Delfaut P, Prakash A, Kaushik RR, Krol RB. Multisite electrode pacing for prevention of atrial fibrillation. *J Cardiovasc Electrophysiol* 1998 Aug;9(8 Suppl):S155-S162.
- (142) Tse HF, Lau CP, Yomtov BM, Ayers GM. Implantable atrial defibrillator with a single-pass dual-electrode lead. *J Am Coll Cardiol* 1999 Jun;33(7):1974-80.
- (143) Tse HF, Lau CP, Yu CM, Lee KL, Michaud GF, Knight BP, et al. Effect of the implantable atrial defibrillator on the natural history of atrial fibrillation. *J Cardiovasc Electrophysiol* 1999 Sep;10(9):1200-9.
- (144) Wellens HJ, Lau CP, Luderitz B, Akhtar M, Waldo AL, Camm AJ, et al. Atrioverter: an implantable device for the treatment of atrial fibrillation. *Circulation* 1998 Oct 20;98(16):1651-6.
- (145) Swerdlow CD, Schissel W, Dijkman B, Jung W, Sheth NV, Olson WH, et al. Detection of atrial fibrillation and flutter by a dual-chamber implantable cardioverter-defibrillator. For the Worldwide Jewel AF Investigators. *Circulation* 2000 Feb 29;101(8):878-85.
- (146) Cox JL, Schuessler RB, Lappas DG, Boineau JP. An 8 1/2-year clinical experience with surgery for atrial fibrillation. *Ann Surg* 1996 Sep;224(3):267-73.
- (147) Cox JL, Schuessler RB, D'Agostino HJ, Jr., Stone CM, Chang BC, Cain ME, et al. The surgical treatment of atrial fibrillation. III. Development of a definitive surgical procedure. *J Thorac Cardiovasc Surg* 1991 Apr;101(4):569-83.
- (148) Cox JL, Boineau JP, Schuessler RB, Kater KM, Lappas DG. Five-year experience with the maze procedure for atrial fibrillation. *Ann Thorac Surg* 1993 Oct;56(4):814-23.
- (149) Cox JL. Minimally-invasive Maze procedure. *Circulation* 100: Suppl I, I-778. 1999.
Ref Type: Abstract

- (150) Packer DL, Asirvatham S, Munger TM. Progress in nonpharmacologic therapy of atrial fibrillation. *J Cardiovasc Electrophysiol* 2003 Dec;14(12 Suppl):S296-S309.
- (151) Hocini M, Sanders P, Jais P, Hsu LF, Takahashi Y, Rotter M, et al. Techniques for curative treatment of atrial fibrillation. *J Cardiovasc Electrophysiol* 2004 Dec;15(12):1467-71.
- (152) Pappone C, Santinelli V. The who, what, why, and how-to guide for circumferential pulmonary vein ablation. *J Cardiovasc Electrophysiol* 2004 Oct;15(10):1226-30.
- (153) Cappato R, Calkins H, Chen SA, Davies W, Iesaka Y, Kalman J, et al. Worldwide survey on the methods, efficacy, and safety of catheter ablation for human atrial fibrillation. *Circulation* 2005 Mar 8;111(9):1100-5.
- (154) Oral H, Scharf C, Chugh A, Hall B, Cheung P, Good E, et al. Catheter ablation for paroxysmal atrial fibrillation: segmental pulmonary vein ostial ablation versus left atrial ablation. *Circulation* 2003 Nov 11;108(19):2355-60.
- (155) Wazni OM, Marrouche NF, Martin DO, Verma A, Bhargava M, Saliba W, et al. Radiofrequency ablation vs antiarrhythmic drugs as first-line treatment of symptomatic atrial fibrillation: a randomized trial. *JAMA* 2005 Jun 1;293(21):2634-40.
- (156) Tse HF, Lau CP. Impact of duration of cryothermal application on clinical efficacy of pulmonary vein isolation using transvenous cryoablation. *Pacing Clin Electrophysiol* 2005 Aug;28(8):839-43.
- (157) Hsieh MH, Tai CT, Tsai CF, Yu WC, Lee SH, Lin YK, et al. Pulmonary vein electrogram characteristics in patients with focal sources of paroxysmal atrial fibrillation. *J Cardiovasc Electrophysiol* 2000 Sep;11(9):953-9.
- (158) Nathan H, Eliakim M. The junction between the left atrium and the pulmonary veins. An anatomic study of human hearts. *Circulation* 1966 Sep;34(3):412-22.
- (159) Saito T, Waki K, Becker AE. Left atrial myocardial extension onto pulmonary veins in humans: anatomic observations relevant for atrial arrhythmias. *J Cardiovasc Electrophysiol* 2000 Aug;11(8):888-94.
- (160) Bliss DF, Hutchins GM. The dorsal mesocardium and development of the pulmonary veins in human embryos. *Am J Cardiovasc Pathol* 1995;5(1):55-67.
- (161) Moubarak JB, Rozwadowski JV, Strzalka CT, Buck WR, Tan WS, Kish GF, et al. Pulmonary veins-left atrial junction: anatomic and histological study. *Pacing Clin Electrophysiol* 2000 Nov;23(11 Pt 2):1836-8.
- (162) Blom NA, Gittenberger-De Groot AC, DeRuiter MC, Poelmann RE, Mentink MM, Ottenkamp J. Development of the cardiac conduction tissue in human embryos using HNK-1 antigen expression: possible relevance for understanding of abnormal atrial automaticity. *Circulation* 1999 Feb 16;99(6):800-6.
- (163) Chen SA, Hsieh MH, Tai CT, Tsai CF, Prakash VS, Yu WC, et al. Initiation of atrial fibrillation by ectopic beats originating from the pulmonary veins: electrophysiological characteristics, pharmacological responses, and effects of radiofrequency ablation. *Circulation* 1999 Nov 2;100(18):1879-86.
- (164) Gerstenfeld EP, Guerra P, Sparks PB, Hattori K, Lesh MD. Clinical outcome after radiofrequency catheter ablation of focal atrial fibrillation triggers. *J Cardiovasc Electrophysiol* 2001 Aug;12(8):900-8.

- (165) Haissaguerre M, Jais P, Shah DC, Garrigue S, Takahashi A, Lavergne T, et al. Electrophysiological end point for catheter ablation of atrial fibrillation initiated from multiple pulmonary venous foci. *Circulation* 2000 Mar 28;101(12):1409-17.
- (166) Lin WS, Prakash VS, Tai CT, Hsieh MH, Tsai CF, Yu WC, et al. Pulmonary vein morphology in patients with paroxysmal atrial fibrillation initiated by ectopic beats originating from the pulmonary veins: implications for catheter ablation. *Circulation* 2000 Mar 21;101(11):1274-81.
- (167) Satoh T, Zipes DP. Unequal atrial stretch in dogs increases dispersion of refractoriness conducive to developing atrial fibrillation. *J Cardiovasc Electrophysiol* 1996 Sep;7(9):833-42.
- (168) GARREY WE, TOWNSEND SE. Neural responses and reactions of the heart of a human embryo. *Am J Physiol* 1948 Feb 1;152(2):219-24.
- (169) MOE GK, RHEINBOLDT WC, ABILDSKOV JA. A COMPUTER MODEL OF ATRIAL FIBRILLATION. *Am Heart J* 1964 Feb;67:200-20.
- (170) WEST TC, LANDA JF. Minimal mass required for induction of a sustained arrhythmia in isolated atrial segments. *Am J Physiol* 1962 Feb;202:232-6.
- (171) Haissaguerre M, Jais P, Shah DC, Arentz T, Kalusche D, Takahashi A, et al. Catheter ablation of chronic atrial fibrillation targeting the reinitiating triggers. *J Cardiovasc Electrophysiol* 2000 Jan;11(1):2-10.
- (172) Mansour M, Mandapati R, Berenfeld O, Chen J, Samie FH, Jalife J. Left-to-right gradient of atrial frequencies during acute atrial fibrillation in the isolated sheep heart. *Circulation* 2001 May 29;103(21):2631-6.
- (173) Skanes AC, Mandapati R, Berenfeld O, Davidenko JM, Jalife J. Spatiotemporal periodicity during atrial fibrillation in the isolated sheep heart. *Circulation* 1998 Sep 22;98(12):1236-48.
- (174) Wu TJ, Ong JJ, Chang CM, Doshi RN, Yashima M, Huang HL, et al. Pulmonary veins and ligament of Marshall as sources of rapid activations in a canine model of sustained atrial fibrillation. *Circulation* 2001 Feb 27;103(8):1157-63.
- (175) Haissaguerre M, Shah DC, Jais P, Hocini M, Yamane T, Deisenhofer I, et al. Electrophysiological breakthroughs from the left atrium to the pulmonary veins. *Circulation* 2000 Nov 14;102(20):2463-5.
- (176) Oral H, Knight BP, Tada H, Ozaydin M, Chugh A, Hassan S, et al. Pulmonary vein isolation for paroxysmal and persistent atrial fibrillation. *Circulation* 2002 Mar 5;105(9):1077-81.
- (177) Pappone C, Oreto G, Rosanio S, Vicedomini G, Tocchi M, Gugliotta F, et al. Atrial electroanatomic remodeling after circumferential radiofrequency pulmonary vein ablation: efficacy of an anatomic approach in a large cohort of patients with atrial fibrillation. *Circulation* 2001 Nov 20;104:2539-44.
- (178) Mangrum JM, Mounsey JP, Kok LC, DiMarco JP, Haines DE. Intracardiac echocardiography-guided, anatomically based radiofrequency ablation of focal atrial fibrillation originating from pulmonary veins. *J Am Coll Cardiol* 2002 Jun 19;39(12):1964-72.
- (179) Nakashima H, Kumagai K, Noguchi H, Tojo H, Yasuda T, Saku K. Evaluation of the recurrence of atrial fibrillation after pulmonary venous ablation. *J Cardiol* 2002 Sep;40(3):87-94.

- (180) Marrouche NF, Dresing T, Cole C, Bash D, Saad E, Balaban K, et al. Circular mapping and ablation of the pulmonary vein for treatment of atrial fibrillation: impact of different catheter technologies. *J Am Coll Cardiol* 2002 Aug 7;40(3):464-74.
- (181) Pappone C, Rosanio S, Oreto G, Tocchi M, Gugliotta F, Vicedomini G, et al. Circumferential radiofrequency ablation of pulmonary vein ostia: A new anatomic approach for curing atrial fibrillation. *Circulation* 2000 Nov 21;102(21):2619-28.
- (182) Pappone C, Oreto G, Lamberti F, Vicedomini G, Loricchio ML, Shpun S, et al. Catheter ablation of paroxysmal atrial fibrillation using a 3D mapping system. *Circulation* 1999 Sep 14;100(11):1203-8.
- (183) Wu CC, Tai CT, Lin YK, Tsao HM, Yu WC, Chen SA. Pulmonary vein dissection during mapping of atrial fibrillation. *J Cardiovasc Electrophysiol* 2001 Apr;12(4):505.
- (184) Ho SY, Sanchez-Quintana D, Cabrera JA, Anderson RH. Anatomy of the left atrium: implications for radiofrequency ablation of atrial fibrillation. *J Cardiovasc Electrophysiol* 1999 Nov;10(11):1525-33.
- (185) Ravenel JG, McAdams HP. Pulmonary venous infarction after radiofrequency ablation for atrial fibrillation. *AJR Am J Roentgenol* 2002 Mar;178(3):664-6.
- (186) Robbins IM, Colvin EV, Doyle TP, Kemp WE, Loyd JE, McMahon WS, et al. Pulmonary vein stenosis after catheter ablation of atrial fibrillation. *Circulation* 1998 Oct 27;98(17):1769-75.
- (187) Yang M, Akbari H, Reddy GP, Higgins CB. Identification of pulmonary vein stenosis after radiofrequency ablation for atrial fibrillation using MRI. *J Comput Assist Tomogr* 2001 Jan;25(1):34-5.
- (188) Ren JF, Marchlinski FE, Callans DJ, Gerstenfeld EP, Dixit S, Lin D, et al. Increased intensity of anticoagulation may reduce risk of thrombus during atrial fibrillation ablation procedures in patients with spontaneous echo contrast. *J Cardiovasc Electrophysiol* 2005 May;16(5):474-7.
- (189) Pappone C, Oral H, Santinelli V, Vicedomini G, Lang CC, Manguso F, et al. Atrio-esophageal fistula as a complication of percutaneous transcatheter ablation of atrial fibrillation. *Circulation* 2004 Jun 8;109(22):2724-6.
- (190) Scanavacca MI, D'avila A, Parga J, Sosa E. Left atrial-esophageal fistula following radiofrequency catheter ablation of atrial fibrillation. *J Cardiovasc Electrophysiol* 2004 Aug;15(8):960-2.
- (191) Mesas CE, Pappone C, Lang CC, Gugliotta F, Tomita T, Vicedomini G, et al. Left atrial tachycardia after circumferential pulmonary vein ablation for atrial fibrillation: electroanatomic characterization and treatment. *J Am Coll Cardiol* 2004 Sep 1;44(5):1071-9.
- (192) Pappone C, Manguso F, Vicedomini G, Gugliotta F, Santinelli O, Ferro A, et al. Prevention of iatrogenic atrial tachycardia after ablation of atrial fibrillation: a prospective randomized study comparing circumferential pulmonary vein ablation with a modified approach. *Circulation* 2004 Nov 9;110(19):3036-42.
- (193) Dehen L, Vilmer C, Humiliere C, Corcos T, Pentousis D, Ollivaud L, et al. Chronic radiodermatitis following cardiac catheterisation: a report of two cases and a brief review of the literature. *Heart* 1999 Mar;81(3):308-12.
- (194) Nahass GT. Acute radiodermatitis after radiofrequency catheter ablation. *J Am Acad Dermatol* 1997 May;36(5 Pt 2):881-4.

- (195) Tsao HM, Wu MH, Yu WC, Tai CT, Lin YK, Hsieh MH, et al. Role of right middle pulmonary vein in patients with paroxysmal atrial fibrillation. *J Cardiovasc Electrophysiol* 2001 Dec;12(12):1353-7.
- (196) Alfke H, Wagner HJ, Klose KJ. A case of an anomalous pulmonary vein of the right middle lobe. *Cardiovasc Intervent Radiol* 1995 Nov;18(6):406-9.
- (197) Rey C, Vaksmann G, Francart C. Anomalous unilateral single pulmonary vein mimicking partial anomalous pulmonary venous return. *Cathet Cardiovasc Diagn* 1986;12(5):330-3.
- (198) Arentz T, Jander N, von RJ, Blum T, Furmaier R, Gornandt L, et al. Incidence of pulmonary vein stenosis 2 years after radiofrequency catheter ablation of refractory atrial fibrillation. *Eur Heart J* 2003 May;24(10):963-9.
- (199) Saad EB, Rossillo A, Saad CP, Martin DO, Bhargava M, Erciyes D, et al. Pulmonary vein stenosis after radiofrequency ablation of atrial fibrillation: functional characterization, evolution, and influence of the ablation strategy. *Circulation* 2003 Dec 23;108(25):3102-7.
- (200) Taylor GW, Kay GN, Zheng X, Bishop S, Ideker RE. Pathological effects of extensive radiofrequency energy applications in the pulmonary veins in dogs. *Circulation* 2000 Apr 11;101(14):1736-42.
- (201) Yu WC, Hsu TL, Tai CT, Tsai CF, Hsieh MH, Lin WS, et al. Acquired pulmonary vein stenosis after radiofrequency catheter ablation of paroxysmal atrial fibrillation. *J Cardiovasc Electrophysiol* 2001 Aug;12(8):887-92.
- (202) Gepstein L, Hayam G, Ben Haim SA. A novel method for nonfluoroscopic catheter-based electroanatomical mapping of the heart. In vitro and in vivo accuracy results. *Circulation* 1997 Mar 18;95(6):1611-22.
- (203) Jongbloed MR, Bax JJ, Lamb HJ, Dirksen MS, Zeppenfeld K, van der Wall EE, et al. Multislice computed tomography versus intracardiac echocardiography to evaluate the pulmonary veins before radiofrequency catheter ablation of atrial fibrillation: a head-to-head comparison. *J Am Coll Cardiol* 2005 Feb 1;45(3):343-50.
- (204) Kato R, Lickfett L, Meining G, Dickfeld T, Wu R, Juang G, et al. Pulmonary vein anatomy in patients undergoing catheter ablation of atrial fibrillation: lessons learned by use of magnetic resonance imaging. *Circulation* 2003 Apr 22;107(15):2004-10.
- (205) Wood MA, Wittkamp M, Henry D, Martin R, Nixon JV, Shepard RK, et al. A comparison of pulmonary vein ostial anatomy by computerized tomography, echocardiography, and venography in patients with atrial fibrillation having radiofrequency catheter ablation. *Am J Cardiol* 2004 Jan 1;93(1):49-53.
- (206) Shah DC, Haissaguerre M, Jais P, Hocini M, Yamane T, Deisenhofer I, et al. Curative catheter ablation of paroxysmal atrial fibrillation in 200 patients: strategy for presentations ranging from sustained atrial fibrillation to no arrhythmias. *Pacing Clin Electrophysiol* 2001 Oct;24(10):1541-58.
- (207) Hirata T, Wolfe SB, Popp RL, Helmen CH, Feigenbaum H. Estimation of left atrial size using ultrasound. *Am Heart J* 1969 Jul;78(1):43-52.
- (208) Lang RM, Bierig M, Devereux RB, Flachskampf FA, Foster E, Pellikka PA, et al. Recommendations for chamber quantification: a report from the American Society of Echocardiography's Guidelines and Standards Committee and the Chamber Quantification Writing Group, developed in conjunction with the European Association of Echocardiography, a branch of the European Society of Cardiology. *J Am Soc Echocardiogr* 2005 Dec;18(12):1440-63.

- (209) Lester SJ, Ryan EW, Schiller NB, Foster E. Best method in clinical practice and in research studies to determine left atrial size. *Am J Cardiol* 1999 Oct 1;84(7):829-32.
- (210) Kircher B, Abbott JA, Pau S, Gould RG, Himelman RB, Higgins CB, et al. Left atrial volume determination by biplane two-dimensional echocardiography: validation by cine computed tomography. *Am Heart J* 1991 Mar;121(3 Pt 1):864-71.
- (211) Rodevan O, Bjørnerheim R, Ljosland M, Maehle J, Smith HJ, Ihlen H. Left atrial volumes assessed by three- and two-dimensional echocardiography compared to MRI estimates. *Int J Card Imaging* 1999 Oct;15(5):397-410.
- (212) Erbel R, Schweizer P, Lambertz H, Henn G, Meyer J, Krebs W, et al. Echoventriculography -- a simultaneous analysis of two-dimensional echocardiography and cineventriculography. *Circulation* 1983 Jan;67(1):205-15.
- (213) Wong M, Shah PM, Taylor RD. Reproducibility of left ventricular internal dimensions with M mode echocardiography: effects of heart size, body position and transducer angulation. *Am J Cardiol* 1981 May;47(5):1068-74.
- (214) Jenkins C, Bricknell K, Marwick TH. Use of real-time three-dimensional echocardiography to measure left atrial volume: comparison with other echocardiographic techniques. *J Am Soc Echocardiogr* 2005 Sep;18(9):991-7.
- (215) Keller AM, Gopal AS, King DL. Left and right atrial volume by freehand three-dimensional echocardiography: in vivo validation using magnetic resonance imaging. *Eur J Echocardiogr* 2000 Mar;1(1):55-65.
- (216) Maddukuri PV, Vieira ML, DeCastro S, Maron MS, Kuvvin JT, Patel AR, et al. What is the best approach for the assessment of left atrial size? Comparison of various unidimensional and two-dimensional parameters with three-dimensional echocardiographically determined left atrial volume. *J Am Soc Echocardiogr* 2006 Aug;19(8):1026-32.
- (217) Jongbloed MR, Schalij MJ, Zeppenfeld K, Oemrawsingh PV, van der Wall EE, Bax JJ. Clinical applications of intracardiac echocardiography in interventional procedures. *Heart* 2005 Jul;91(7):981-90.
- (218) Kalman JM, Fitzpatrick AP, Olgin JE, Chin MC, Lee RJ, Scheinman MM, et al. Biophysical characteristics of radiofrequency lesion formation in vivo: dynamics of catheter tip-tissue contact evaluated by intracardiac echocardiography. *Am Heart J* 1997 Jan;133(1):8-18.
- (219) Cummings JE, Schweikert RA, Saliba WI, Burkhardt JD, Brachmann J, Gunther J, et al. Assessment of temperature, proximity, and course of the esophagus during radiofrequency ablation within the left atrium. *Circulation* 2005 Jul 26;112(4):459-64.
- (220) Marrouche NF, Martin DO, Wazni O, Gillinov AM, Klein A, Bhargava M, et al. Phased-array intracardiac echocardiography monitoring during pulmonary vein isolation in patients with atrial fibrillation: impact on outcome and complications. *Circulation* 2003 Jun 3;107(21):2710-6.
- (221) Verma A, Marrouche NF, Yamada H, Grimm RA, Cummings J, Burkhardt JD, et al. Usefulness of intracardiac Doppler assessment of left atrial function immediately post-pulmonary vein antrum isolation to predict short-term recurrence of atrial fibrillation. *Am J Cardiol* 2004 Oct 1;94(7):951-4.

- (222) Yamanaka K, Fujita M, Doi K, Tsuneyoshi H, Yamazato A, Ueno K, et al. Multislice computed tomography accurately quantifies left atrial size and function after the MAZE procedure. *Circulation* 2006 Jul 4;114:I5-I9.
- (223) Marom EM, Herndon JE, Kim YH, McAdams HP. Variations in pulmonary venous drainage to the left atrium: implications for radiofrequency ablation. *Radiology* 2004 Mar;230(3):824-9.
- (224) Schwartzman D, Lacomis J, Wigginton WG. Characterization of left atrium and distal pulmonary vein morphology using multidimensional computed tomography. *J Am Coll Cardiol* 2003 Apr 16;41(8):1349-57.
- (225) Kistler PM, Earley MJ, Harris S, Abrams D, Ellis S, Sporton SC, et al. Validation of three-dimensional cardiac image integration: use of integrated CT image into electroanatomic mapping system to perform catheter ablation of atrial fibrillation. *J Cardiovasc Electrophysiol* 2006 Apr;17(4):341-8.
- (226) Tops LF, Bax JJ, Zeppenfeld K, Jongbloed MR, Lamb HJ, van der Wall EE, et al. Fusion of multislice computed tomography imaging with three-dimensional electroanatomic mapping to guide radiofrequency catheter ablation procedures. *Heart Rhythm* 2005 Oct;2(10):1076-81.
- (227) Pilleul F, Merchant N. MRI of the pulmonary veins: comparison between 3D MR angiography and T1-weighted spin echo. *J Comput Assist Tomogr* 2000 Sep;24(5):683-7.
- (228) Tsao HM, Yu WC, Cheng HC, Wu MH, Tai CT, Lin WS, et al. Pulmonary vein dilation in patients with atrial fibrillation: detection by magnetic resonance imaging. *J Cardiovasc Electrophysiol* 2001 Jul;12(7):809-13.
- (229) Wittkampf FH, Vonken EJ, Derkzen R, Loh P, Velthuis B, Wever EF, et al. Pulmonary vein ostium geometry: analysis by magnetic resonance angiography. *Circulation* 2003 Jan 7;107(1):21-3.
- (230) Therkelsen SK, Groenning BA, Svendsen JH, Jensen GB. Atrial and ventricular volume and function in persistent and permanent atrial fibrillation, a magnetic resonance imaging study. *J Cardiovasc Magn Reson* 2005;7(2):465-73.
- (231) Fleck T, Wolf F, Bader T, Lehner R, Aigner C, Stix G, et al. Atrial function after ablation procedure in patients with chronic atrial fibrillation using steady-state free precession magnetic resonance imaging. *Ann Thorac Surg* 2007 Nov;84(5):1600-4.
- (232) Jayam VK, Dong J, Vasamreddy CR, Lickfett L, Kato R, Dickfeld T, et al. Atrial volume reduction following catheter ablation of atrial fibrillation and relation to reduction in pulmonary vein size: an evaluation using magnetic resonance angiography. *J Interv Card Electrophysiol* 2005 Jul;13:107-14.
- (233) Tsao HM, Wu MH, Huang BH, Lee SH, Lee KT, Tai CT, et al. Morphologic remodeling of pulmonary veins and left atrium after catheter ablation of atrial fibrillation: insight from long-term follow-up of three-dimensional magnetic resonance imaging. *J Cardiovasc Electrophysiol* 2005 Jan;16(1):7-12.
- (234) Hudsmith LE, Petersen SE, Francis JM, Robson MD, Neubauer S. Normal human left and right ventricular and left atrial dimensions using steady state free precession magnetic resonance imaging. *J Cardiovasc Magn Reson* 2005;7(5):775-82.
- (235) Anderson JL, Horne BD, Pennell DJ. Atrial dimensions in health and left ventricular disease using cardiovascular magnetic resonance. *J Cardiovasc Magn Reson* 2005;7(4):671-5.

- (236) Tseng WY, Liao TY, Wang JL. Normal systolic and diastolic functions of the left ventricle and left atrium by cine magnetic resonance imaging. *J Cardiovasc Magn Reson* 2002;4(4):443-57.
- (237) Calkins H. Three dimensional mapping of atrial fibrillation: techniques and necessity. *J Interv Card Electrophysiol* 2005 Aug;13 Suppl 1:53-9.
- (238) Dong J, Dickfeld T, Dalal D, Cheema A, Vasamreddy CR, Henrikson CA, et al. Initial experience in the use of integrated electroanatomic mapping with three-dimensional MR/CT images to guide catheter ablation of atrial fibrillation. *J Cardiovasc Electrophysiol* 2006 May;17(5):459-66.
- (239) Dong J, Calkins H, Solomon SB, Lai S, Dalal D, Lardo AC, et al. Integrated electroanatomic mapping with three-dimensional computed tomographic images for real-time guided ablations. *Circulation* 2006 Jan 17;113(2):186-94.
- (240) Nagel E, Underwood R, Pennell D, Sechtem UP, Neubauers S, Wickline S, et al. New developments in non-invasive cardiac imaging: critical assessment of the clinical role of cardiac magnetic resonance imaging. *Eur Heart J* 1998 Sep;19(9):1286-93.
- (241) Pennell DJ, Sechtem UP, Higgins CB, Manning WJ, Pohost GM, Rademakers FE, et al. Clinical indications for cardiovascular magnetic resonance (CMR): Consensus Panel report. *Eur Heart J* 2004 Nov;25(21):1940-65.
- (242) Poustchi-Amin M, Gutierrez FR, Brown JJ, Miowitz SA, Narra VR, Takahashi N, et al. Performing cardiac MR imaging: an overview. *Magn Reson Imaging Clin N Am* 2003 Feb;11(1):1-18.
- (243) Higgins CB. Cardiac imaging. *Radiology* 2000 Oct;217(1):4-10.
- (244) Lipton MJ, Boxt LM, Hijazi ZM. Role of the radiologist in cardiac diagnostic imaging. *AJR Am J Roentgenol* 2000 Dec;175(6):1495-506.
- (245) Poustchi-Amin M, Gutierrez FR, Brown JJ, Miowitz SA, Narra VR, Takahashi N, et al. Performing cardiac MR imaging: an overview. *Magn Reson Imaging Clin N Am* 2003 Feb;11(1):1-18.
- (246) Rademakers FE. Cardiac physiology. In: Higgins CB, de Ross A, editors. *Cardiovascular MRI and MRA*. Lippincott Williams and Wilkins; 2003. p. 38-50.
- (247) Bluemke DA, Boxerman JL, Atalar E, McVeigh ER. Segmented K-space cine breath-hold cardiovascular MR imaging: Part 1. Principles and technique. *AJR Am J Roentgenol* 1997 Aug;169(2):395-400.
- (248) Scheffler K, Lehnhardt S. Principles and applications of balanced SSFP techniques. *Eur Radiol* 2003 Nov;13(11):2409-18.
- (249) Lotz J, Meier C, Leppert A, Galanski M. Cardiovascular flow measurement with phase-contrast MR imaging: basic facts and implementation. *Radiographics* 2002 May;22(3):651-71.
- (250) Wilke NM, Jerosch-Herold M, Zenovich A, Stillman AE. Magnetic resonance first-pass myocardial perfusion imaging: clinical validation and future applications. *J Magn Reson Imaging* 1999 Nov;10(5):676-85.
- (251) Rogers WJ, Jr., Kramer CM, Geskin G, Hu YL, Theobald TM, Vido DA, et al. Early contrast-enhanced MRI predicts late functional recovery after reperfused myocardial infarction. *Circulation* 1999 Feb 16;99(6):744-50.

- (252) Zerhouni EA, Parish DM, Rogers WJ, Yang A, Shapiro EP. Human heart: tagging with MR imaging--a method for noninvasive assessment of myocardial motion. *Radiology* 1988 Oct;169(1):59-63.
- (253) Alley MT, Shifrin RY, Pelc NJ, Herfkens RJ. Ultrafast contrast-enhanced three-dimensional MR angiography: state of the art. *Radiographics* 1998 Mar;18(2):273-85.
- (254) Sechtem U, Pflugfelder PW, Gould RG, Cassidy MM, Higgins CB. Measurement of right and left ventricular volumes in healthy individuals with cine MR imaging. *Radiology* 1987 Jun;163(3):697-702.
- (255) Chuang ML, Hibberd MG, Salton CJ, Beaudin RA, Riley MF, Parker RA, et al. Importance of imaging method over imaging modality in noninvasive determination of left ventricular volumes and ejection fraction: assessment by two- and three-dimensional echocardiography and magnetic resonance imaging. *J Am Coll Cardiol* 2000 Feb;35(2):477-84.
- (256) Jauhainen T, Jarvinen VM, Hekali PE, Poutanen VP, Penttila A, Kupari M. MR gradient echo volumetric analysis of human cardiac casts: focus on the right ventricle. *J Comput Assist Tomogr* 1998 Nov;22(6):899-903.
- (257) Helbing WA, Rebergen SA, Maliepaard C, Hansen B, Ottenkamp J, Reiber JH, et al. Quantification of right ventricular function with magnetic resonance imaging in children with normal hearts and with congenital heart disease. *Am Heart J* 1995 Oct;130(4):828-37.
- (258) Pattynama PM, Lamb HJ, Van d, V, van der Geest RJ, van der Wall EE, de RA. Reproducibility of MRI-derived measurements of right ventricular volumes and myocardial mass. *Magn Reson Imaging* 1995;13(1):53-63.
- (259) Mohiaddin RH, Hasegawa M. Measurement of atrial volumes by magnetic resonance imaging in healthy volunteers and in patients with myocardial infarction. *Eur Heart J* 1995 Jan;16(1):106-11.
- (260) Ishimoto N, Ito M, Kinoshita M. Signal-averaged P-wave abnormalities and atrial size in patients with and without idiopathic paroxysmal atrial fibrillation. *Am Heart J* 2000 Apr;139(4):684-9.
- (261) Jarvinen V, Kupari M, Hekali P, Poutanen VP. Assessment of left atrial volumes and phasic function using cine magnetic resonance imaging in normal subjects. *Am J Cardiol* 1994 Jun 1;73(15):1135-8.
- (262) Jarvinen VM, Kupari MM, Hekali PE, Poutanen VP. Right atrial MR imaging studies of cadaveric atrial casts and comparison with right and left atrial volumes and function in healthy subjects. *Radiology* 1994 Apr;191(1):137-42.
- (263) Jarvinen VM, Kupari MM, Poutanen VP, Hekali PE. A simplified method for the determination of left atrial size and function using cine magnetic resonance imaging. *Magn Reson Imaging* 1996;14(3):215-26.
- (264) Hauser TH, McClenen S, Katsimbris G, Josephson ME, Manning WJ, Yeon SB. Assessment of left atrial volume by contrast enhanced magnetic resonance angiography. *J Cardiovasc Magn Reson* 2004;6(2):491-7.
- (265) Poutanen T, Ikonen A, Vainio P, Jokinen E, Tikanoja T. Left atrial volume assessed by transthoracic three dimensional echocardiography and magnetic resonance imaging: dynamic changes during the heart cycle in children. *Heart* 2000 May;83(5):537-42.

- (266) Mansour M, Holmvang G, Sosnovik D, Migrino R, Abbara S, Ruskin J, et al. Assessment of pulmonary vein anatomic variability by magnetic resonance imaging: implications for catheter ablation techniques for atrial fibrillation. *J Cardiovasc Electrophysiol* 2004 Apr;15(4):387-93.
- (267) Carreras F, Guillaumet E, Pujadas S, Lopez-Salguero R, Ligero C, Leta R, et al. [Morphologic study of the pulmonary veins by magnetic resonance angiography]. *Rev Esp Cardiol* 2005 Oct;58(10):1226-9.
- (268) Ho SY, Cabrera JA, Tran VH, Farre J, Anderson RH, Sanchez-Quintana D. Architecture of the pulmonary veins: relevance to radiofrequency ablation. *Heart* 2001 Sep;86(3):265-70.
- (269) Ho SY. Pulmonary vein ablation in atrial fibrillation: does anatomy matter? *J Cardiovasc Electrophysiol* 2003 Feb;14(2):156-7.
- (270) Tritto M, De PR, Zardini M, Spadacini G, Oliveira M, Salerno-Uriarte JA. Electrical connections between pulmonary veins in humans: evidence after radiofrequency ablation of the venoatrial junction. *Circulation* 2001 Aug 14;104(7):E30-E31.
- (271) Keane D. New catheter ablation techniques for the treatment of cardiac arrhythmias. *Card Electrophysiol Rev* 2002 Dec;6(4):341-8.
- (272) Keane D, Ruskin J. Pulmonary vein isolation for atrial fibrillation. *Rev Cardiovasc Med* 2002;3(4):167-75.
- (273) Schmidt B, Ernst S, Ouyang F, Chun KR, Broemel T, Bansch D, et al. External and endoluminal analysis of left atrial anatomy and the pulmonary veins in three-dimensional reconstructions of magnetic resonance angiography: the full insight from inside. *J Cardiovasc Electrophysiol* 2006 Sep;17(9):957-64.
- (274) Hocini M, Jais P, Sanders P, Takahashi Y, Rotter M, Rostock T, et al. Techniques, evaluation, and consequences of linear block at the left atrial roof in paroxysmal atrial fibrillation: a prospective randomized study. *Circulation* 2005 Dec 13;112(24):3688-96.
- (275) Dittrich HC, Pearce LA, Asinger RW, McBride R, Webel R, Zabalgoitia M, et al. Left atrial diameter in nonvalvular atrial fibrillation: An echocardiographic study. *Stroke Prevention in Atrial Fibrillation Investigators*. *Am Heart J* 1999 Mar;137:494-9.
- (276) Scharf C, Sneider M, Case I, Chugh A, Lai SW, Pelosi F, Jr., et al. Anatomy of the pulmonary veins in patients with atrial fibrillation and effects of segmental ostial ablation analyzed by computed tomography. *J Cardiovasc Electrophysiol* 2003 Feb;14(2):150-5.
- (277) Stambler BS, Fenelon G, Shepard RK, Clemo HF, Guiraudon CM. Characterization of sustained atrial tachycardia in dogs with rapid ventricular pacing-induced heart failure. *J Cardiovasc Electrophysiol* 2003 May;14:499-507.
- (278) Lemola K, Desjardins B, Sneider M, Case I, Chugh A, Good E, et al. Effect of left atrial circumferential ablation for atrial fibrillation on left atrial transport function. *Heart Rhythm* 2005 Sep;2:923-8.
- (279) Beukema WP, Elvan A, Sie HT, Misier AR, Wellens HJ. Successful radiofrequency ablation in patients with previous atrial fibrillation results in a significant decrease in left atrial size. *Circulation* 2005 Oct 4;112:2089-95.
- (280) Lemola K, Sneider M, Desjardins B, Case I, Chugh A, Hall B, et al. Effects of left atrial ablation of atrial fibrillation on size of the left atrium and pulmonary veins. *Heart Rhythm* 2004 Nov;1:576-81.

- (281) Tops LF, Bax JJ, Zeppenfeld K, Jongbloed MR, van der Wall EE, Schalij MJ. Effect of radiofrequency catheter ablation for atrial fibrillation on left atrial cavity size. *Am J Cardiol* 2006 Apr 15;97:1220-2.
- (282) Reant P, Lafitte S, Jais P, Serri K, Weerasooriya R, Hocini M, et al. Reverse remodeling of the left cardiac chambers after catheter ablation after 1 year in a series of patients with isolated atrial fibrillation. *Circulation* 2005 Nov 8;112:2896-903.
- (283) Khan IA. Atrial stunning: basics and clinical considerations. *Int J Cardiol* 2003 Dec;92(2-3):113-28.
- (284) Stefanadis C, Dernellis J, Toutouzas P. A clinical appraisal of left atrial function. *Eur Heart J* 2001 Jan;22(1):22-36.
- (285) Thomas L, McKay T, Byth K, Marwick TH. Abnormalities of left atrial function after cardioversion: an atrial strain rate study. *Heart* 2007 Jan;93(1):89-95.
- (286) Tsang TS, Barnes ME, Abhayaratna WP, Cha SS, Gersh BJ, Langins AP, et al. Effects of quinapril on left atrial structural remodeling and arterial stiffness. *Am J Cardiol* 2006 Mar 15;97:916-20.
- (287) Hagens VE, Van Veldhuisen DJ, Kamp O, Rienstra M, Bosker HA, Veeger NJ, et al. Effect of rate and rhythm control on left ventricular function and cardiac dimensions in patients with persistent atrial fibrillation: results from the RAtc Control versus Electrical Cardioversion for Persistent Atrial Fibrillation (RACE) study. *Heart Rhythm* 2005 Jan;2:19-24.
- (288) Westenberg JJ, van der Geest RJ, Lamb HJ, Versteegh MI, Braun J, Doornbos J, et al. MRI to evaluate left atrial and ventricular reverse remodeling after restrictive mitral annuloplasty in dilated cardiomyopathy. *Circulation* 2005 Aug 30;112:I437-I442.
- (289) Handke M, Harloff A, Hetzel A, Olszewski M, Bode C, Geibel A. Left atrial appendage flow velocity as a quantitative surrogate parameter for thromboembolic risk: determinants and relationship to spontaneous echocontrast and thrombus formation--a transesophageal echocardiographic study in 500 patients with cerebral ischemia. *J Am Soc Echocardiogr* 2005 Dec;18(12):1366-72.
- (290) Kizer JR, Bella JN, Palmieri V, Liu JE, Best LG, Lee ET, et al. Left atrial diameter as an independent predictor of first clinical cardiovascular events in middle-aged and elderly adults: the Strong Heart Study (SHS). *Am Heart J* 2006 Feb;151(2):412-8.
- (291) Tsang TS, Abhayaratna WP, Barnes ME, Miyasaka Y, Gersh BJ, Bailey KR, et al. Prediction of cardiovascular outcomes with left atrial size: is volume superior to area or diameter? *J Am Coll Cardiol* 2006 Mar 7;47(5):1018-23.
- (292) Benjamin EJ, D'Agostino RB, Belanger AJ, Wolf PA, Levy D. Left atrial size and the risk of stroke and death. The Framingham Heart Study. *Circulation* 1995 Aug 15;92(4):835-41.
- (293) Beruezo A, Tamborero D, Mont L, Benito B, Tolosana JM, Sitges M, et al. Pre-procedural predictors of atrial fibrillation recurrence after circumferential pulmonary vein ablation. *Eur Heart J* 2007 Apr;28(7):836-41.
- (294) Kirchhof P, Auricchio A, Bax J, Crijns H, Camm J, Diener HC, et al. Outcome parameters for trials in atrial fibrillation: executive summary. *Eur Heart J* 2007 Nov;28(22):2803-17.
- (295) Petersen P, Kastrup J, Brinch K, Godtfredsen J, Boysen G. Relation between left atrial dimension and duration of atrial fibrillation. *Am J Cardiol* 1987 Aug 1;60(4):382-4.

- (296) D'Andrea A, Caso P, Romano S, Scarafale R, Riegler L, Salerno G, et al. Different effects of cardiac resynchronization therapy on left atrial function in patients with either idiopathic or ischaemic dilated cardiomyopathy: a two-dimensional speckle strain study. *Eur Heart J* 2007 Nov;28(22):2738-48.
- (297) Yu CM, Fang F, Zhang Q, Yip GW, Li CM, Chan JY, et al. Improvement of atrial function and atrial reverse remodeling after cardiac resynchronization therapy for heart failure. *J Am Coll Cardiol* 2007 Aug 21;50(8):778-85.
- (298) Anwar AM, Geleijnse ML, Soliman OI, Nemes A, ten Cate FJ. Left atrial Frank-Starling law assessed by real-time, three-dimensional echocardiographic left atrial volume changes. *Heart* 2007 Nov;93(11):1393-7.
- (299) Anwar AM, Soliman OI, Geleijnse ML, Nemes A, Vletter WB, ten Cate FJ. Assessment of left atrial volume and function by real-time three-dimensional echocardiography. *Int J Cardiol* 2008 Jan 11;123(2):155-61.
- (300) Thomas L. Assessment of atrial function. *Heart Lung Circ* 2007 Jun;16(3):234-42.
- (301) Cheitlin MD, Armstrong WF, Aurigemma GP, Beller GA, Bierman FZ, Davis JL, et al. ACC/AHA/ASE 2003 guideline update for the clinical application of echocardiography: summary article: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines (ACC/AHA/ASE Committee to Update the 1997 Guidelines for the Clinical Application of Echocardiography). *Circulation* 2003 Sep 2;108(9):1146-62.
- (302) Barkhausen J, Ruehm SG, Goyen M, Buck T, Laub G, Debatin JF. MR evaluation of ventricular function: true fast imaging with steady-state precession versus fast low-angle shot cine MR imaging: feasibility study. *Radiology* 2001 Apr;219(1):264-9.
- (303) Plein S, Bloomer TN, Ridgway JP, Jones TR, Bainbridge GJ, Sivananthan MU. Steady-state free precession magnetic resonance imaging of the heart: comparison with segmented k-space gradient-echo imaging. *J Magn Reson Imaging* 2001 Sep;14(3):230-6.
- (304) Gentles PJ, Sauer WH, Gerstenfeld EP, Lin D, Dixit S, Zado E, et al. Reversal of left ventricular dysfunction following ablation of atrial fibrillation. *J Cardiovasc Electrophysiol* 2007 Jan;18:9-14.
- (305) Thomas L, Boyd A, Thomas SP, Schiller NB, Ross DL. Atrial structural remodelling and restoration of atrial contraction after linear ablation for atrial fibrillation. *Eur Heart J* 2003 Nov;24:1942-51.
- (306) Verma A, Kilicaslan F, Adams JR, Hao S, Beheiry S, Minor S, et al. Extensive ablation during pulmonary vein antrum isolation has no adverse impact on left atrial function: an echocardiography and cine computed tomography analysis. *J Cardiovasc Electrophysiol* 2006 Jul;17(7):741-6.
- (307) Thomas MD, Kalra PR, Jones A, Struthers AD, More RS. Time course for recovery of atrial mechanical and endocrine function post DC cardioversion for persistent atrial fibrillation. *Int J Cardiol* 2005 Jul 20;102(3):487-91.
- (308) Abhayaratna WP, Seward JB, Appleton CP, Douglas PS, Oh JK, Tajik AJ, et al. Left atrial size: physiologic determinants and clinical applications. *J Am Coll Cardiol* 2006 Jun 20;47(12):2357-63.
- (309) Casaclang-Verzosa G, Gersh BJ, Tsang TS. Structural and functional remodeling of the left atrium: clinical and therapeutic implications for atrial fibrillation. *J Am Coll Cardiol* 2008 Jan 1;51(1):1-11.

- (310) Victal OA, Teerlink JR, Gaxiola E, Wallace AW, Najar S, Camacho DH, et al. Left ventricular volume reduction by radiofrequency heating of chronic myocardial infarction in patients with congestive heart failure. *Circulation* 2002 Mar 19;105(11):1317-22.
- (311) Zipes DP. Atrial fibrillation. A tachycardia-induced atrial cardiomyopathy. *Circulation* 1997 Feb 4;95(3):562-4.
- (312) Thomas L, Thomas SP, Hoy M, Boyd A, Schiller NB, Ross DL. Comparison of left atrial volume and function after linear ablation and after cardioversion for chronic atrial fibrillation. *Am J Cardiol* 2004 Jan 15;93(2):165-70.
- (313) Jais P, Peng JT, Shah DC, Garrigue S, Hocini M, Yamane T, et al. Left ventricular diastolic dysfunction in patients with so-called lone atrial fibrillation. *J Cardiovasc Electrophysiol* 2000 Jun;11(6):623-5.
- (314) Bode F, Katchman A, Woosley RL, Franz MR. Gadolinium decreases stretch-induced vulnerability to atrial fibrillation. *Circulation* 2000 May 9;101(18):2200-5.
- (315) Nazir SA, Lab MJ. Mechanoelectric feedback in the atrium of the isolated guinea-pig heart. *Cardiovasc Res* 1996 Jul;32(1):112-9.
- (316) Ravelli F, Allessie M. Effects of atrial dilatation on refractory period and vulnerability to atrial fibrillation in the isolated Langendorff-perfused rabbit heart. *Circulation* 1997 Sep 2;96(5):1686-95.
- (317) Sideris DA, Toumanidis ST, Tselepatiotis E, Kostopoulos K, Stringli T, Kitsiou T, et al. Atrial pressure and experimental atrial fibrillation. *Pacing Clin Electrophysiol* 1995 Sep;18(9 Pt 1):1679-85.
- (318) Thomas SP, Nicholson IA, Nunn GR, Rees A, Trieu L, Daly MP, et al. Effect of atrial radiofrequency ablation designed to cure atrial fibrillation on atrial mechanical function. *J Cardiovasc Electrophysiol* 2000 Jan;11(1):77-82.
- (319) Avitall B, Urbonas A, Urboniene D, Millard S, Helms R. Time course of left atrial mechanical recovery after linear lesions: normal sinus rhythm versus a chronic atrial fibrillation dog model. *J Cardiovasc Electrophysiol* 2000 Dec;11(12):1397-406.
- (320) Chang SH, Tsao HM, Wu MH, Tai CT, Chang SL, Wongcharoen W, et al. Morphological changes of the left atrial appendage after catheter ablation of atrial fibrillation. *J Cardiovasc Electrophysiol* 2007 Jan;18(1):47-52.
- (321) Packer DL, Stevens CL, Curley MG, Bruce CJ, Miller FA, Khandheria BK, et al. Intracardiac phased-array imaging: methods and initial clinical experience with high resolution, under blood visualization: initial experience with intracardiac phased-array ultrasound. *J Am Coll Cardiol* 2002 Feb 6;39(3):509-16.
- (322) Saad EB, Marrouche NF, Saad CP, Ha E, Bash D, White RD, et al. Pulmonary vein stenosis after catheter ablation of atrial fibrillation: emergence of a new clinical syndrome. *Ann Intern Med* 2003 Apr 15;138(8):634-8.
- (323) Mansour M, Ruskin J, Keane D. Efficacy and safety of segmental ostial versus circumferential extra-ostial pulmonary vein isolation for atrial fibrillation. *J Cardiovasc Electrophysiol* 2004 May;15(5):532-7.
- (324) Pappone C, Rosanio S, Augello G, Gallus G, Vicedomini G, Mazzone P, et al. Mortality, morbidity, and quality of life after circumferential pulmonary vein ablation for atrial fibrillation: outcomes from a controlled nonrandomized long-term study. *J Am Coll Cardiol* 2003 Jul 16;42(2):185-97.

- (325) Silva RM, Mont L, Berrueto A, Fosch X, Wayar L, Alvarenga N, et al. [Radiofrequency ablation in the treatment of focal atrial fibrillation using circumferential mapping and segmentary disconnection of pulmonary veins]. *Rev Esp Cardiol* 2003 Apr;56(4):361-7.
- (326) Vasamreddy CR, Jayam V, Bluemke DA, Calkins H. Pulmonary vein occlusion: an unanticipated complication of catheter ablation of atrial fibrillation using the anatomic circumferential approach. *Heart Rhythm* 2004 May;1(1):78-81.
- (327) Karch MR, Zrenner B, Deisenhofer I, Schreieck J, Ndreppepa G, Dong J, et al. Freedom from atrial tachyarrhythmias after catheter ablation of atrial fibrillation: a randomized comparison between 2 current ablation strategies. *Circulation* 2005 Jun 7;111(22):2875-80.
- (328) Packer DL, Keelan P, Munger TM, Breen JF, Asirvatham S, Peterson LA, et al. Clinical presentation, investigation, and management of pulmonary vein stenosis complicating ablation for atrial fibrillation. *Circulation* 2005 Feb 8;111(5):546-54.
- (329) Firstenberg MS, Greenberg NL, Smedira NG, Prior DL, Scalia GM, Thomas JD, et al. Doppler echo evaluation of pulmonary venous-left atrial pressure gradients: human and numerical model studies. *Am J Physiol Heart Circ Physiol* 2000 Aug;279(2):H594-H600.
- (330) Ren JF, Marchlinski FE, Callans DJ, Zado ES. Intracardiac Doppler echocardiographic quantification of pulmonary vein flow velocity: an effective technique for monitoring pulmonary vein ostia narrowing during focal atrial fibrillation ablation. *J Cardiovasc Electrophysiol* 2002 Nov;13(11):1076-81.
- (331) Saad EB, Cole CR, Marrouche NF, Dresing TJ, Perez-Lugones A, Saliba WI, et al. Use of intracardiac echocardiography for prediction of chronic pulmonary vein stenosis after ablation of atrial fibrillation. *J Cardiovasc Electrophysiol* 2002 Oct;13(10):986-9.
- (332) Ernst S, Ouyang F, Goya M, Lober F, Schneider C, Hoffmann-Riem M, et al. Total pulmonary vein occlusion as a consequence of catheter ablation for atrial fibrillation mimicking primary lung disease. *J Cardiovasc Electrophysiol* 2003 Apr;14(4):366-70.
- (333) Perez-Lugones A, Schwartzman PR, Schweikert R, Tchou PJ, Saliba W, Marrouche NF, et al. Three-dimensional reconstruction of pulmonary veins in patients with atrial fibrillation and controls: morphological characteristics of different veins. *Pacing Clin Electrophysiol* 2003 Jan;26(1 Pt 1):8-15.
- (334) Qureshi AM, Prieto LR, Latson LA, Lane GK, Mesia CI, Radvansky P, et al. Transcatheter angioplasty for acquired pulmonary vein stenosis after radiofrequency ablation. *Circulation* 2003 Sep 16;108(11):1336-42.
- (335) Sohn RH, Schiller NB. Left upper pulmonary vein stenosis 2 months after radiofrequency catheter ablation of atrial fibrillation. *Circulation* 2000 Apr 4;101(13):E154-E155.
- (336) Ueda M, Tada H, Kurosaki K, Itoi K, Koyama K, Naito S, et al. Pulmonary vein morphology before and after segmental isolation in patients with atrial fibrillation. *Pacing Clin Electrophysiol* 2005 Sep;28(9):944-53.
- (337) Maksimovic R, Scholten MF, Cademartiri F, Jordaens LJ, Pattynama PM. Sixteen multidetector row computed tomography of pulmonary veins: 3-months' follow-up after treatment of paroxysmal atrial fibrillation with cryothermal ablation. *Eur Radiol* 2005 Jun;15(6):1122-7.
- (338) Ghaye B, Szapiro D, Dacher JN, Rodriguez LM, Timmermans C, Devillers D, et al. Percutaneous ablation for atrial fibrillation: the role of cross-sectional imaging. *Radiographics* 2003 Oct;23 Spec No:S19-S33.

- (339) Dill T, Neumann T, Ekinci O, Breidenbach C, John A, Erdogan A, et al. Pulmonary vein diameter reduction after radiofrequency catheter ablation for paroxysmal atrial fibrillation evaluated by contrast-enhanced three-dimensional magnetic resonance imaging. *Circulation* 2003 Feb 18;107(6):845-50.
- (340) Arentz T, Weber R, Jander N, Burkle G, von RJ, Blum T, et al. Pulmonary haemodynamics at rest and during exercise in patients with significant pulmonary vein stenosis after radiofrequency catheter ablation for drug resistant atrial fibrillation. *Eur Heart J* 2005 Jul;26(14):1410-4.