

UNIVERSITAT AUTÒNOMA DE BARCELONA



***In vitro* embryo production from prepubertal
goat oocytes in different culture media**

Sondes Hammami

Facultat de Veterinària

Departament de Ciència Animal i dels Aliments

Tesis Doctoral

Bellaterra, Junio del 2014

La **Dra. Dolors Izquierdo Tugas**, professora agregada del Departament de Ciència Animal i dels Aliments de la Universitat Autònoma de Barcelona,

I la **Dra. Roser Morató Molet**, investigadora Juan de la Cierva, amb destí al Departament de Biologia de la Universitat de Girona, Biotecnologia de la Reproducció Animal i Humana de la Universitat de Girona,

CERTIFIQUEN

Que **Sondes Hammami** ha realitzat sota la seva direcció el present treball d'investigació que porta per títol “*In vitro embryo production from prepubertal goat oocytes in different culture media* ” per optar al grau de Doctor per la Universitat Autònoma de Barcelona. Aquest treball ha estat finançat pel Ministeri de Ciència i Innovació, a través del projecte AGL 2011-23784, el grup de recerca consolidat SGR2009SGR0621 i una beca atorgada pel Ministeri d'Assumptes Exteriors i de Cooperació.

I per a que així consti, firmen el present certificat

Dra. Dolors Izquierdo i Tugas

Dra. Roser Morató Molet

Bellaterra, Maig del 2014

Aquest treball ha estat finançat pel Ministerio de Ciencia e Innovación, a través del projecte AGL2011-23784 i el grup de recerca consolidat SGR2009SGR0621.

À ma mère,
À mon frère
À mes soeurs

Abstract

The prepubertal goat ovary shows a large number of small oocytes with a compromised competence to develop up to blastocyst stage following IVM, IVF and IVC. Culture conditions used to support *in vitro* embryo production appears to be an important factor to suit the requirements of these oocytes and to increase the blastocyst yield. In the present thesis work, we have carried out two studies to evaluate the effects of addition of different supplements to *in vitro* maturation and culture media on embryo development and the quality of resultant blastocyst in prepubertal goat oocytes.

In oocytes from prepubertal females, different types and concentrations of substances including gonadotrophic hormones, growth factors, protein sources and antioxidants, such as insulin-transferrin-selenium (ITS) and L-ascorbic acid (AA), have been shown to improve oocyte cytoplasmic maturation and embryo development. The aim of the first study was to assess the effects of adding low level of hormones and/or ITS plus AA to IVM medium through the developmental competence and embryo quality of small oocytes from prepubertal goats. Specifically, we tested four maturation media: conventional IVM medium (CM), CM+ITS+AA and low level of hormones (named as Growth Medium; GM), CM with low level of hormones (modified CM; mCM) and CM+ITS+AA and normal level of hormones (modified GM; mGM). Cumulus-Oocyte Complexes (COCs) were classified into two categories according to oocyte diameter: small (<125 μm) and large ($\geq 125 \mu\text{m}$) oocytes. Large oocytes were matured in CM and small oocytes were matured in different combination of CM, GM, mCM or mGM. After IVM, oocytes were fertilized and after 24h presumptive zygotes were cultured for 8 days. Results of this study showed that using different combinations of CM, GM, mCM, and mGM for IVM of prepubertal goat small oocytes did not have a beneficial effect on the percentage of blastocysts. The high blastocyst rate was only related to oocyte diameter. However, the culture of small oocytes in GM improved the quality of blastocysts and enhanced their survival post-warming compared to CM (53.3% vs 30.8%, respectively; $P < 0.05$).

The second study examined the effect of activin-A during IVM and embryo culture on the blastocyst rate of prepubertal goat oocytes. Activin is a member of the transforming growth factor β superfamily that plays a functional role in the process of cellular proliferation and differentiation. In experiment 1, with a view to search for optimal

concentration of recombinant human activin-A during IVM, three concentrations (0, 10 and 100 ng/mL) were tested on oocytes meiotic maturation and subsequent *in vitro* embryo development. The experiment 2 was performed to evaluate the effect of adding 10 ng/mL activin-A at the IVM and/or IVC media on embryo development and blastocyst quality. Results of this study demonstrated that the addition of activin-A to IVM yielded similar percentages of maturation and blastocyst. Moreover, the addition of 10 ng/mL of activin-A throughout the whole embryo culture significantly increased rates of development to the blastocyst stage as compared with the control group ($19.5 \pm 2.21\%$ vs $13.1 \pm 2.37\%$, respectively; $P < 0.05$). Blastocyst quality was not improved in the presence of activin-A neither during IVM evaluated after Hoechst 33342 staining, nor during IVM and/ or IVC evaluated by differential staining.

In conclusion, IVM of prepubertal goat small oocytes in GM would be useful to improve the quality of blastocysts produced *in vitro* and the presence of activin-A during IVC enhances embryo development of prepubertal goat oocytes.

Resumen

Los ovarios de cabras prepúberes proporcionan una población de ovocitos de pequeño diámetro con una menor competencia para el desarrollo hasta blastocisto después de la maduración, fecundación y posterior cultivo *in vitro* (MIV, FIV y CIV). Las condiciones del cultivo *in vitro* aportan los requerimientos necesarios a los ovocitos e influyen sobre su posterior desarrollo embrionario. En esta tesis hemos llevado a cabo dos estudios para evaluar los efectos de la adición de diferentes suplementos a los medios de MIV y CIV sobre el desarrollo embrionario y la calidad de los blastocistos obtenidos a partir de ovocitos de cabras prepúberes.

En ovocitos de hembras prepúberes, la adición de sustancias, incluyendo las hormonas gonadotrópicas, factores de crecimiento, fuentes de proteínas y antioxidantes, como son el complejo insulina-transferrina-selenio (ITS) y el L-ácido ascórbico (AA), han demostrado producir una mejora en la maduración citoplasmática de los ovocitos y su subsiguiente desarrollo embrionario *in vitro*. El objetivo del primer estudio fue evaluar los efectos de la adición de AA, ITS y un bajo nivel hormonal en el medio de MIV sobre el desarrollo embrionario y la calidad de los blastocistos obtenidos. En concreto, hemos utilizado 4 medios maduración: Medio Tradicional (MT), Medio de Crecimiento (MC): MT + ITS + AA + bajo nivel hormonal, Medio Tradicional modificado (MTm): MT + bajo nivel hormonal y Medio de Crecimiento modificado (MCm): MT + ITS + AA + nivel hormonal normal. Los complejos-ovocito-cúmulus (COCs) se clasificaron en dos grupos según el diámetro ovocitario: pequeños ($<125\ \mu\text{m}$) y grandes ($\geq 125\ \mu\text{m}$). Los ovocitos $\geq 125\ \mu\text{m}$ fueron madurados en MT mientras que los ovocitos $<125\ \mu\text{m}$ fueron madurados en combinaciones de MT, MC, MTm y MCm. Después de la MIV, los ovocitos fueron fecundados *in vitro* y posteriormente los embriones fueron CIV durante 8 días. En este estudio se observó que el desarrollo embrionario de los ovocitos $<125\ \mu\text{m}$ no mejoró en las diferentes combinaciones de medios de MIV estudiadas (MT, MC, MTm y MCm). El mayor desarrollo hasta blastocisto se obtuvo en el grupo de embriones procedentes de ovocitos $\geq 125\ \mu\text{m}$. Sin embargo, cuando los ovocitos de menor diámetro se maduraron en MC, se observó una mejoría significativa en la calidad de los blastocistos obtenidos, optimizando la supervivencia de los embriones tras la descongelación comparado con el MT (53.3% vs 30.8%, respectivamente; $P < 0.05$).

En el segundo estudio se analizó el efecto de la suplementación de los medios de MIV y CIV con activina-A sobre el desarrollo embrionario. La activina-A es un miembro de la superfamilia del factor de crecimiento transformante β (TGF- β) y cumple un papel importante en la proliferación y diferenciación celular. Con el fin de determinar el efecto de la activina-A sobre los ovocitos de cabras prepúberes, se realizaron 2 experimentos, analizando la maduración nuclear de los ovocitos, su posterior desarrollo embrionario y calidad de los blastocistos obtenidos. En el primer experimento se estudió la suplementación del medio de MIV con varias concentraciones de activina-A (0, 10 y 100 ng/mL) y en el segundo experimento se evaluó la adición de 10 ng/mL de activina-A a los medios de MIV y CIV. Los resultados de este estudio demostraron que la suplementación del medio de MIV con diferentes concentraciones de activina-A proporciona similares porcentajes de maduración nuclear y de blastocistos comparado con la MIV sin activina-A. Además, la adición de 10 ng/mL de activina-A durante el CIV aumentó significativamente las tasas de desarrollo hasta blastocisto en comparación con el grupo control ($19.5 \pm 2.21\%$ vs $13.1 \pm 2.37\%$, respectivamente; $P < 0.05$). Sin embargo, no se observó una mejora en la calidad de los blastocistos cuando se añadió activina-A en el medio de MIV (evaluado por la tinción de Hoechst 33342) y/o en el de CIV (evaluado por la tinción diferencial).

En conclusión, la maduración *in vitro* de los ovocitos de cabras prepúberes de diámetro $< 125 \mu\text{m}$ en medio de crecimiento sería una alternativa útil para mejorar la calidad de los blastocistos producidos *in vitro*, mientras que la presencia de activina-A durante el cultivo *in vitro* incrementa el porcentaje de embriones desarrollados hasta el estadio de blastocisto.

Index

Chapter 1: Introduction	15
Chapter 2: Literature Review	23
2.1. <i>In vitro</i> maturation (IVM).....	25
2.1.1. Nuclear maturation	25
2.1.2. Cytoplasmic maturation	26
2.1.3. Factors affecting oocyte quality	26
2.1.3.1. Effect of follicle and oocyte size	27
2.1.3.2. Age of the donor	29
2.1.3.3. Media and supplements used for IVM	30
2.1.3.3.1. Hormones	31
2.1.3.3.2. Antioxidants	32
2.1.3.3.3. Insulin-Transferrin-Selenium	34
2.1.3.3.4. Activin-A	35
2.1.3.3.5. Serum and other substances	37
2.2. <i>In vitro</i> fertilization (IVF).....	39
2.3. <i>In vitro</i> culture (IVC).....	40
2.4. Assessment of embryo quality	43
Chapter 3: Objectives	47
Chapter 4: Developmental competence and embryo quality of small oocytes from pre-pubertal goats cultured in IVM medium supplemented with low level of hormones, insulin-transferrin-selenium and ascorbic acid	51
Chapter 5: <i>In vitro</i> developmental competence of prepubertal goat oocytes cultured with recombinant activin-A	59
Chapter 6: General Discussion	69
Chapter 7: General Conclusions	77
Chapter 8: References	81
Chapter 9: Acknowledgments	107

List of abbreviations

μg	Microgram
μm	Micrometer
AA	Ascorbic Acid
AI	Artificial Insemination
ARTs	Assisted Reproductive Technologies
AI-TI	Anaphase I- Telophase I
BCB	Brilliant Cresyl Blue
CM	Conventional Medium
COCs	Cumulus–Oocyte Complexes
DNA	Desoxyribonucleic Acid
E ₂ -17 β	Estradiol-17 β
EGF	Epidermal Growth Factor
ET	Embryo Transfer
FCS	Fetal Calf Serum
FF	Follicular Fluid
FSH	Follicle Stimulating Hormone
GDF9	Growth-Differentiation Factor 9
GH	Growth Hormone
GM	Growth Medium
GnRH	Gonadotrophin-Releasing Hormone
GV	Germinal Vesicle
GVBD	Germinal Vesicle Breakdown
ICM	Inner Cell Mass
ICSI	Intracytoplasmic Sperm Injection
IGF-I	Insulin-Like Growth Factor-I
ITS	Insulin-Transferrin-Selenium
IU	International Unit
IVC	<i>In Vitro</i> Culture
IVEP	<i>In Vitro</i> Embryo Production
IVF	<i>In Vitro</i> Fertilization
IVM	<i>In Vitro</i> Maturation
IVP	<i>In Vitro</i> Production

JIVET	Juvenile <i>In Vitro</i> Embryo Technology
LH	Luteinizing Hormone
LOPU	Laparoscopic Ovum Pick-Up
MI	Metaphase I
MII	Metaphase II
mL	Milliliter
mm	Millimeter
mM	Millimolar
MOET	Multiple Ovulation and Embryo Transfer
MPF	Maturation Promoting Factor
ng	Nanogram
OPU	Ovum Pick-Up
PN	Pronuclei
PUFAs	Polyunsaturated Fatty Acids
RNA	Ribonucleic Acid
ROS	Reactive Oxygen Species
SR	Serum Replacer
TE	Trophectoderm
TGF β	Transforming Growth Factor β
ZP	Zona Pellucida

INTRODUCTION

Introduction

In recent years, demand for goat products has increased in both developing and developed countries; the population of goats in the world has increased by nearly 100% over the past four decades. In 2012 there were more than 921 million goats around the world: Asia had the greatest number, followed by Africa, South America, Europe, North America and Oceania (<http://www.fao.org/>). Goats have always had a major role to play in developing countries due to their adaptability under harsh and marginal rural conditions, predominantly in the tropical and subtropical zones of the world.

A growing interest in assisted reproductive technologies (ARTs) has occurred. ARTs are being used for enhancement of reproductive performance and genetic improvement in sheep and goats. In addition, they can have substantial contribution in preservation of endangered species or breeds (Cognié *et al.*, 2004). While their applications are widespread in cattle, in small ruminants it is almost restricted to artificial insemination. Nonetheless, during the last 30 years, considerable progress has been made in sheep and goat embryo technologies.

Furthermore, *in vitro* embryo production (IVEP) in small ruminants provides an excellent source of low cost embryos for basic research on developmental biology and physiology and for commercial application of the emerging biotechnologies such as nuclear transfer, transgenesis, embryo sexing and stem cells. Thus, goats offer a good model for the development of these technologies (reviewed by Cognié *et al.*, 2003).

In the laboratory, embryos can be routinely produced and developed up to the blastocyst stage using three subsequent techniques: *in vitro* maturation (IVM) of oocytes recovered from ovaries, followed by sperm capacitation and *in vitro* fertilization (IVF) of matured oocytes and then *in vitro* culture (IVC) of the fertilized oocytes up to the blastocyst stage that can be transferred to recipient females or cryopreserved for future use. However, *in vitro* embryo production still is not very efficient due to several limiting factors affecting the outcome of each step of the process.

The use of prepubertal females as oocyte donors for *in vitro* embryo production (juvenile *in vitro* embryo technology or JIVET) offers considerable potential for accelerating genetic gain in domestic livestock through reduced generation interval as compared to classical approach (Armstrong *et al.*, 1997). However, reduced

developmental competence in oocytes of juvenile animals as compared to their adult counterpart when matured *in vitro*, was addressed in numerous studies on farm animals including bovine (Presicce *et al.*, 1997), ovine (Ledda *et al.*, 1997), caprine (Leoni *et al.*, 2009) and porcine (Marchal *et al.*, 2001).

Evidence suggests that the low number of embryos obtained from prepubertal females is mainly due to the low competence of these oocytes, and this competence is defined by Sirard *et al.*, (2006) as the ability to: 1) resume meiosis; 2) cleave following fertilization; 3) develop to the blastocyst stage; 4) induce pregnancy; and 5) bring offspring to term in good health, and it is acquired progressively during oocyte growth (Sorensen and Wassarman, 1976; Eppig, 1996).

Embryo development is influenced by events occurring during oocyte maturation. For successful IVM, oocytes must undergo nuclear and cytoplasmic maturation. Therefore the efficiency of embryo development from these oocytes is mainly attributed to limitations induced by two factors: failure to select the most competent oocytes for *in vitro* production and an inadequate understanding of oocyte maturation and culture conditions to provide an optimized environment, necessary to support the physiological changes associated with oocytes maturation and required for attaining developmental competence (Rizos *et al.*, 2002a; Rizos *et al.*, 2002b; Morton *et al.*, 2008).

To overcome the reduced maturational competence of oocytes from juvenile females, several factors and strategies have been shown to affect the oocyte competence such as: selection follicles and oocytes diameters (Anguita *et al.*, 2007; Romaguera *et al.*, 2010a), donor's age (Ledda *et al.*, 1997; Baldassarre and Karatzas, 2004), hormonal stimulation (Koeman *et al.*, 2003), *in vitro* maturation protocol (Wu *et al.*, 2006) and supplementation to oocyte maturation media (Mertens *et al.*, 2005; Romaguera *et al.*, 2010b; Córdova *et al.*, 2011).

Numerous research groups have reported a positive relationship between follicle size and the ability of its oocyte to develop to blastocyst stage in cows (Lonergan *et al.*, 1994), ewes (Cognié *et al.*, 1998), sows (Motlik *et al.*, 1984), buffalos (Raghu *et al.*, 2002a) and goats (Martino *et al.*, 1994; Crozet *et al.*, 1995). The developmental competence of an oocyte is obtained progressively as follicular diameter increases to maximum size, suggesting that the follicle size and oocyte diameter are closely related, and as both increase oocyte competence is gradually acquired within the ovary.

According to oocyte diameter, previous studies in our laboratory concluded that a higher blastocyst rate is obtained in oocytes larger than 135 μm (12.5%) compared to oocytes of 125 to 135 μm diameter (1.95%) after IVF (Anguita *et al.*, 2007). However, when ICSI was used to fertilize oocytes, Jiménez-Macedo *et al.*, (2006) did not find differences between these two oocytes categories.

A range of oocyte maturation media and additives have been tried to accomplish with the *in vitro* maturation of oocytes, and their developmental competence. Reports in the literature have described the effect of maturing oocytes *in vitro* in culture medium supplemented with different types and concentrations of substances including gonadotrophic hormones, antioxidants, growth factors and protein sources. The addition of gonadotropins (FSH and LH) and estradiol-17 β (E₂-17 β), to the maturation medium significantly has improved maturation rates (nuclear, cytoplasmic maturation and cumulus expansion) in mammalian oocytes (Sanbuissho and Threlfall, 1990; Mattioli *et al.*, 1991; Singh *et al.*, 1993; Guler *et al.*, 2000; Abdoon *et al.*, 2001; Roberts *et al.*, 2005; Alvarez *et al.*, 2009), has promoted the ability of the oocytes to form a male pronucleus after *in vitro* fertilization (Thibault *et al.*, 1975; Funahashi and Day, 1993) and their developmental competence up to blastocyst stage (Men *et al.*, 2002). The concentration of gonadotropins used during the process of IVM of oocytes has been showed to determine the results of oocyte maturation and developmental competence. In bovine, Liu *et al.*, (2011) reported that extremely high concentrations of gonadotropins have detrimental effects on oocyte nuclear maturation and embryo development and increase apoptosis in cumulus cells. In porcine, Bing *et al.*, (2001) have reported that using a low concentration of FSH (1 $\mu\text{g}/\text{mL}$) is not sufficient to induce full nuclear maturation of oocytes, compared with 10 $\mu\text{g}/\text{mL}$ FSH. On the other hand, some studies have shown that lower concentrations of hormones are enough for IVM in mice (Peluso, 1988; Byskov *et al.*, 1997). Also, in prepubertal gilts Wu *et al.*, (2006) have observed a significant increase in blastocyst yield derived from oocytes obtained from small follicles by reducing 250-fold the concentration of the hormones in a protocol of IVM medium (2-step culture system). Results of those studies suggest the importance of judicious use of gonadotropins to enhance *in vitro* oocyte maturation and embryonic development, depending on the species and the donor age.

During *in vitro* culture, cells are exposed to higher concentrations of oxygen than those that occur *in vivo* and this causes the constant production of free radicals. High levels of free radicals cause damage to cell components by lipid peroxidation, protein modification, and DNA damage resulting in impaired cell function and subsequently can affect oocyte maturation and further embryonic development (Cetica *et al.*, 2001). Thus, to optimize embryo production, oocytes need to be protected against oxidative stress during *in vitro* culture. Several antioxidants have been added to maturation media (Ali *et al.*, 2003). Insulin-transferrin-selenium (ITS) combination is routinely used in culture systems to promote the development of oocytes because of its protecting effects from oxidative damage by reducing free radical production and inhibiting lipid peroxidation, and the promoting effect of the uptake of glucose and amino acids (Tatemoto *et al.*, 2004; Ebert *et al.*, 2006). Studies performed in several species during *in vitro* maturation, showed that ITS addition during IVM, improved cytoplasmic maturation and decreased polyspermic fertilization rate in porcine oocytes (Jeong *et al.*, 2008). Similarly, in bovine, supplementation with ITS (Jaakma *et al.*, 1997; Bowles and Lishman, 1998) showed a higher blastocyst rate. In mice, an increase on blastocyst rate (Zhang and Armstrong, 1990) as well as their quality (Jeong *et al.*, 2008) was observed when IVM medium was supplemented with ITS. Moreover, the ascorbic acid is an antioxidant that has been shown to support cytoplasmic maturation of porcine oocytes relating to developmental competence after fertilization by alleviating oxidative stress (Tatemoto *et al.*, 2001). In addition, the presence of ascorbic acid in the culture medium has been shown to significantly reduce the level of apoptosis in mouse oocyte-associated granulosa cells although no improvement of oocyte developmental competence was detected (Eppig *et al.*, 2000). Ascorbic acid addition plus the ITS complex to the culture medium also enhances the blastocyst production of gilts (Wu *et al.*, 2006) and bovine oocytes (Córdova *et al.*, 2010).

A second approach to improve the developmental competence of oocytes after their recovery from the follicle has been the addition of growth promoting substances to *in vitro* maturation and culture media, such as epidermal growth factor (EGF), growth hormone (GH), inhibin, activin and follistatin. Those growth factors are generally added to the basal medium to increase cell proliferation and to stimulate specific cell functions (Van der Valk *et al.*, 2010) despite modest improvements in development have been achieved in this way. Activin-A is an important member of the transforming growth

factor β (TGF β) superfamily. In the ovaries of vertebrates, activin-A is expressed predominantly in the granulosa cell layer of follicles, suggesting important roles in processes such as folliculogenesis, steroid hormone production, and oocyte maturation as paracrine or autocrine factors (Peng and Mukai, 2000). Studies *in vitro* showed that exposure of oocytes during maturation and/or culture to different concentrations of exogenous activin-A, stimulates the resumption of meiosis of oocytes in several species (mice (Itoh *et al.*, 1990); cattle (Stock *et al.*, 1997; Park *et al.*, 2010; Trigal *et al.*, 2011); rhesus monkey (Alak *et al.*, 1996) and human (Alak *et al.*, 1998)). Moreover, a positive effect of activin-A on embryo development were shown when the protein was added at the earliest stages of bovine (Yoshioka and Kamomae, 1996, Yoshioka *et al.*, 1998a, Lee *et al.*, 2009) embryo culture.

The basal components of the TGF β superfamily signalling pathways are the type II and type I transmembrane serine and threonine kinase receptors and the cytoplasmic Smad proteins (Heldin *et al.*, 1997). Activins signal through type I and type II receptor proteins, both of which are serine/threonine kinases. Subsequently, signals such as Smad proteins are phosphorylated (Peng and Mukai, 2000). Activin receptors are represented by two isoforms: activin receptors type IA (ActR-IA), IB (ActR-IB), IIA (ActR-IIA) and IIB (ActR-IIB). Both protein and mRNA for activin-A and activin receptors type I and II have been localized in both oocyte and granulosa cells of follicles at various developmental stages in a variety of species, including human (Eramaa *et al.*, 1995; Mather *et al.*, 1997; Sidis *et al.*, 1998), rodent (Zhao *et al.*, 2001), porcine (Van den Hurk and Van de Pavert, 2001), and bovine (Hulshof *et al.*, 1997) follicles and caprine ovary and follicles (Silva *et al.*, 2004; 2006). These facts suggest that activin-A may play a role in development of prepubertal goat oocytes, and investigation about the mechanism of action of this protein in oocytes from prepubertal goat during *in vitro* maturation and during embryo *in vitro* culture is required because, to date, no studies on the effects of activin-A on *in vitro* developmental potential of prepubertal goat oocytes have been reported.

In summary, juvenile donors will have an important place in livestock-improvement as well as in biotechnology programs. However, further investigations on molecular and cellular approaches will increase our understanding of cytoplasmic maturation in prepubertal oocytes allowing a more efficient *in vitro* production of embryos by IVM/IVF, ICSI and nuclear transfer, and the production of offspring from juveniles to

reduce the generation interval and increase the rate of genetic progress in breeding schemes.

All these points above mentioned need to be addressed in future studies to find the most efficient *in vitro* embryo production to achieve high rates of oocyte maturation and embryo development from prepubertal goats.

Continuing efforts of previous research performed in our laboratory with the aim to improve the *in vitro* embryo yield of prepubertal goat oocytes, the present thesis work is designed to evaluate the effects of addition of different supplements to *in vitro* maturation and culture media on embryo development.

LITERATURE REVUE

Literature Review

Over recent years, considerable research into *in vitro* embryo production (IVEP) technology has been undertaken in an attempt to determine which conditions are needed during the three subsequent steps: *in vitro* maturation (IVM), *in vitro* fertilization (IVF) and *in vitro* culture (IVC). These *in vitro* processes are intended to mimic the processes *in vivo*, for improving animal breeding programs and biomedical field. These steps will be discussed in more detail in subsequent sections of this thesis.

2.1. *In vitro* maturation (IVM)

Full maturation involves both nuclear and cytoplasmic events that confer on the oocyte the capacity for supporting normal fertilization and early embryonic development (Thibault and Gerard, 1973; Moor and Trounson, 1977). Mammalian oocytes arrested at the germinal vesicle (GV) stage acquire the capacity to resume meiosis during growth of the ovarian follicle from the primary to the preovulatory stage. As removed from antral follicles of females ovaries, cumulus–oocyte complexes (COCs) can be induced to undergo *in vitro* the sequence of events found during *in vivo* maturation. Matured oocytes reach metaphase II stage (nuclear maturation), accompanying by the molecular and structural changes that allow them to support normal fertilization and embryonic development (cytoplasmic maturation) (Bevers *et al.*, 1997; Watson, 2006).

2.1.1. Nuclear maturation

Oocytes that are collected from growing follicles for IVEP are blocked at the prophase of the first meiotic division. As soon as they are removed from follicles and transferred to a culture medium, the oocytes can spontaneously resume the meiotic division (Edwards, 1965). The resumption of meiosis is characterized by dissolution of the nuclear membrane and a condensation of the chromatin, a nuclear stage referred as germinal vesicle breakdown (GVBD). Chromosomes are then arranged in metaphase I (MI) stage which is followed by anaphase I to telophase I (AI-TI) transition and oocytes progress to metaphase II (MII) where they remain until their fertilization. Nuclear maturation can be visualized by the extrusion of the first polar body. Studies *in vitro* (Samake *et al.*, 2000; Bormann *et al.*, 2003), showed that under specific conditions, pre-

selected oocytes can resume the nuclear maturation reaching 70–90% of the oocytes the MII stage. Prepubertal goat oocytes recovered from follicles larger than 3 mm (Martino *et al.*, 1994), were able to reach the MII stage at a rate of 72 % after IVM.

2.1.2. Cytoplasmic maturation

Embryo development is strongly influenced by events occurring during oocyte maturation. IVM seems to be the limiting factor, because even after selection of a homogenous population of cumulus-oocyte complexes, only 35% will attain full cytoplasmic maturation and possess the competence to produce a viable, transferable blastocyst (Blondin and Sirard, 1995). Cytoplasmic maturation covers all morphologic and molecular events accompanying nuclear maturation after LH surge in preovulatory follicles and preparing oocyte cytoplasm to successful fertilization and embryo development. As reviewed by Mermillod *et al.*, (2006), cytoplasmic maturation includes well known morphological modifications, such as the migration of cortical granules in the cortical region of the ooplasm. These granules are stored during oocyte growth and release their enzymatic content in the perivitellin space after fertilization. Cytoplasmic maturation also includes the accumulation of mRNA, proteins, alterations to Golgi complexes, mitochondrial accumulation in the ooplasm, and cumulus expansion (Sirard, 2001). Cumulus expansion can importantly be used to microscopically assess the *in vitro* maturation rate of oocytes (Gupta *et al.*, 2005). This store of maternal RNA instill upon the oocyte a capacity to decondense and remodel the sperm head, to form the pronuclei (PN), zygote formation, early embryogenesis and normal fetal development (Mermillod *et al.*, 1999; Watson, 2006).

2.1.3. Factors affecting oocyte quality

Good quality oocyte maturation is an essential prerequisite for embryo development. Naturally the oocyte quality is determined by the oocyte's ability to mature, be fertilized and give rise to normal offspring after gestation (Sirard *et al.*, 2006). This capacity is acquired during folliculogenesis, as the oocyte grows and during the period of oocyte maturation (Krisner, 2004). Proper oocyte selection in the laboratory is crucial for successful embryo production. A study performed in bovine (Blondin and Sirard, 1995)

showed that the best indicators of an immature oocyte ability to undergo maturation and embryonic development were the presence of an intact complement of cumulus cell layers surrounding the oocyte and a homogenous appearing cytoplasm. Also, the follicular and oocyte sizes, as well as several factors, such as age of the donor animal, and the media used for maturing the oocytes are some factors that have been linked to the maturational competence of oocytes and have therefore been proposed as selection criteria for oocyte quality, as a key factor in determining the development of embryos to the blastocyst stage (Lonergan *et al.*, 2003).

2.1.3.1. Effect of follicle and oocyte size

The size of the follicle seems to be an important factor in the selection of potential oocytes (reviewed by Sirard *et al.*, 2006), involving RNA or protein stores as factors involved in oocyte competence. Increased developmental competence of oocytes has been associated with increased follicular diameter as reported in several studies in various species. In cattle, Lonergan *et al.*, (1994) reported a higher proportion of blastocysts obtained from follicles > 6 mm compared to 2–6 mm follicles. Similarly, in calves, Kauffold *et al.*, (2005) showed an increase in blastocyst production in oocytes coming from follicles with diameter > 8 mm than from follicles of < 8 mm. It has been suggested that the reason for the differences between the follicular diameters on oocyte quality is due to their content. The follicular fluid (FF) constitutes the microenvironment of the oocyte during follicular maturation and contains molecules involved in nuclear and cytoplasmic maturation, ovulation and fertilization (Yoshida *et al.*, 1992). Thereby, Ali *et al.*, (2004) illustrated the effect of follicle diameter by the use of bovine follicular fluid obtained from large (> 8 mm) and small follicles (2–5 mm) as a supplement of *in vitro* maturation media of bovine oocytes. Results of this study showed that following fertilization and embryo culture, more oocytes reached the blastocyst stage when oocytes were cultured with FF from large follicles compared with FF derived from small follicles. Similarly, in buffalo (Raghu *et al.*, 2002a), sheep (adult: Cognié *et al.*, 1998; prepubertal: Ledda *et al.*, 1999b), goats (adult: De Smedt *et al.*, 1994; Crozet *et al.*, 1995; prepubertal: Romaguera *et al.*, 2010a) and pigs (Marchal *et al.*, 2002) has been reported that the acquisition of meiotic competence as well as the ability to develop up to the blastocyst stage is acquired sequentially as the follicle

enlarges. In prepubertal goats, Romaguera *et al.*, (2010a) demonstrated that oocytes from follicles of ≥ 3 mm showed greater mean oocyte diameter, higher percentages of fragmented DNA cells, higher cleavage rates and greater developmental competence to the blastocyst stage than oocytes from follicles of < 3 mm.

The comparison of the oocyte diameter is often used as a marker for oocyte maturity or meiotic competence, able to attain their full developmental competence to blastocysts *in vitro*, as there is an intensive synthesis of RNA during this phase that causes an increase in size. According to Brevini *et al.*, (2007), during the oocytes growth, messenger RNAs and proteins of maternal origin are accumulated into the oocyte throughout its growth in the ovary, upon fertilization, several mechanisms are activated that control the appropriate use of such material and prepare for the synthesis of new products supporting fertilization and initiating embryo development. The association between the oocyte diameter and its ability to resume and complete meiotic maturation *in vitro* has been described in several farm animals. In bovine, Otoi *et al.*, (1997) classified oocytes in six categories according to oocyte diameter ($< 110 \mu\text{m}$, 110 to $< 115 \mu\text{m}$, 115 to $< 120 \mu\text{m}$, 120 to $< 125 \mu\text{m}$, 125 to $< 130 \mu\text{m}$ and $\geq 130 \mu\text{m}$), and concluded that bovine oocytes have acquired full meiotic competence at a diameter of $115 \mu\text{m}$ but not yet attained full developmental competence to blastocysts, and that oocytes have acquired full developmental competence at a diameter of $120 \mu\text{m}$. In ovine, Shirazi and Sadeghi (2007) reported no significant differences in the percentage of oocytes that reached the MII stage (81, 82, and 84%) with diameters of $< 110 \mu\text{m}$, 110 – $150 \mu\text{m}$, $> 150 \mu\text{m}$, respectively. In buffalo, the rate of *in vitro* blastocyst production was significantly higher in oocytes with diameters greater than $145 \mu\text{m}$ (Raghu *et al.*, 2002a). In prepubertal goats, oocytes were classified by Anguita *et al.*, (2007) and Jiménez-Macedo *et al.*, (2006), into 4 categories of diameter: $< 110 \mu\text{m}$; 110 to $125 \mu\text{m}$; 125 to $135 \mu\text{m}$ and $> 135 \mu\text{m}$, results of those studies showed that oocytes smaller than $125 \mu\text{m}$ fertilized by IVF (Anguita *et al.*, 2007) and ICSI (Jiménez-Macedo *et al.*, 2006), were unable to develop up to blastocyst stage. Anguita *et al.*, (2007) observed that the oocyte diameter was positively related to the percentage of oocytes at MII after IVM (0, 21, 58 and 78%, respectively) and the percentage of blastocysts obtained at 8 days postinsemination (0, 0, 1.95 and 12.5%, respectively). However, using ICSI to fertilize these oocytes categories, the percentage of ICSI derived blastocysts (blastocysts/injected oocytes) obtained from oocytes of 125 – $135 \mu\text{m}$ diameter had

similar blastocyst development to oocytes larger than 135 μm (15.9 and 11.1%, respectively) (Jiménez-Macedo *et al.*, 2006). This difference between oocyte categories after IVF and ICSI protocols could be explained by the fact that oocytes selected to perform ICSI have completed their nuclear maturation. Also, the lack of polyspermic zygotes by the microinjection could be one other explanation.

2.1.3.2. Age of the donor

There is a general agreement upon the fact that the use of prepubertal females as oocytes donors plays an important role on *in vitro* embryo production programs, by the reduction of the generation interval and consequently increasing the intensity of breeding. However, oocytes derived from juvenile females show a reduced developmental competence as reported in numerous studies on farm species including bovine (Revel *et al.*, 1995; Damiani *et al.*, 1996; Khatir *et al.*, 1998), ovine (Ledda *et al.*, 1997; O'Brien *et al.*, 1997b), and porcine (Marchal *et al.*, 2002). In caprine, using *in vitro* produced zygotes from laparoscopic ovum pick-up (LOPU), both cleavage and blastocyst development rates of embryos from adult donors have been higher than those from prepubertal donors (90 and 16% vs. 82 and 6%, respectively) (Wang *et al.*, 2002a).

Many factors may reduce the development competence of oocytes collected from prepubertal animals. In bovine, the lower developmental ability of oocytes from 3-month-old calves compared with that of cyclic cow oocytes may depend on some defective endocrine environment encountered *in vivo* before the onset of puberty (Revel *et al.*, 1995). Gandolfi *et al.*, (1998) observed differences in size between calf and cow oocytes (118 μm and 123 μm , respectively) and showed that the difference in the developmental competence may be induced because of difference in gene expression abundance between adult and the prepubertal oocytes, showing a reduced protein synthesis in oocytes and cumulus cells from calves. It has been also reported that in oocytes from prepubertal donors, structural changes are delayed and incomplete and may contribute to failures of appropriate zona pellucida (ZP) changes (reviewed by Slavik *et al.*, 2005). In prepubertal goats, it has been revealed functional deficiencies in cytoplasmic maturation of oocytes, such as altered distribution of cortical granules (Velilla *et al.*, 2004) and mitochondria (Velilla *et al.*, 2006), disorganization of microtubule and microfilament (Velilla *et al.*, 2005) and alteration in total RNA content,

p34 (cdc2) and cyclin B1 expression as well as maturation promoting factor (MPF) activity (Anguita *et al.*, 2007; 2008). After *in vitro* fertilization, a low incidence of sperm head decondensation at fertilization (Martino *et al.*, 1995; Mogas *et al.*, 1997b), a high percentage of haploid embryos (Villamediana *et al.*, 2001) and, consequently, a low rate of blastocyst production (Izquierdo *et al.*, 2002) has been also observed.

However, other reports suggest that donor age may not be the only important criterion, since oocytes coming from follicles (> 3mm diameter) of 45 days old goats have the capacity to *in vitro* develop to blastocyst stage as well as oocytes derived from adult goats (Romaguera *et al.*, 2011). Thus, the low embryo development of prepubertal females oocytes could be related to the small number of large follicles in their ovaries, as reported by Martino *et al.*, (1994) since only 1.1% of follicles were larger than 3 mm.

Studies in ovine (Earl *et al.*, 1994; O'Brien *et al.*, 1997a; Ledda *et al.*, 1999a) prove a better effectiveness of using this source of oocytes applying a previous stimulation of prepubertal females with exogeneous gonadotropins. A positive effect of gonadotropins on follicular and oocyte size could be the cause of the lack of differences in embryo development between prepubertal and adult oocytes, which allow the oocytes to develop up to blastocyst stage similarly to that obtained when oocytes from adult sheep. Besides, these oocytes showed higher diameter than oocytes from unstimulated lambs. Also in caprine, following ovarian stimulation and laparoscopic recovery, prepubertal and adult goat oocytes cultured in semi-defined media presented a similar developmental competence to blastocyst stage (Koeman *et al.*, 2003).

For all above mentioned and in order to make use of these prepubertal oocytes more efficiently, it is important to develop culture systems that permit oocytes to acquire the competence for undergoing maturation, fertilization, and development up to blastocyst stage *in vitro* in a similar way than their adult counterparts and those coming from *in vivo*.

2.1.3.3. Media and supplements used for IVM

In vitro maturation is the most critical part of the whole process of *in vitro* embryo production. Due to the heterogeneous nature of immature oocytes (oocytes from a range of follicle stages) used for IVEP, the *in vitro* maturation of oocyte can be influenced by

culture media components and culture conditions used for IVM (Cognié *et al.*, 2004). Several culture media have been proposed for IVM (MEM, Waymouth, Ham-F12, etc.). However, the most widely used seems to be the TCM199 medium, bicarbonate buffered and containing minerals, carbon and energy sources (glucose, glutamine) as well as vitamins and amino acids (reviewed by Mermillod *et al.*, 2006), supplemented with L-glutamine, pyruvate, FSH, LH, estradiol-17 β (E₂-17 β), plus complex fluids (heat-treated serum, follicular fluid) (reviewed by Cognié *et al.*, 2003; Tibary *et al.*, 2005; Paramio, 2010). Nevertheless, there are some research teams using synthetic oviductal fluid (SOF) medium instead of TCM199 in sheep (Shabankareh and Akhondi, 2012) and goat (Ongeri *et al.*, 2001; Bormann *et al.*, 2003; Herrick *et al.*, 2004) oocytes.

2.1.3.3.1. Hormones

During *in vivo* follicular development, the mammalian follicular oocyte is arrested at diplotene stage of meiosis I, and resumption from this arrest depends upon the preovulatory surge of gonadotropins (Eppig, 1996). Gonadotropin hormones and E₂-17 β are generally used in *in vitro* maturation protocols. It has been shown to improve nuclear and cytoplasmic oocyte maturation as well as expansion of the surrounding cumulus cells of several species including porcine (Mattioli *et al.*, 1991; Singh *et al.*, 1993; Schoevers *et al.*, 2003; Alvarez *et al.*, 2009), ovine (Guler *et al.*, 2000), murine (Roberts *et al.*, 2005), bovine (Sanbuissho and Threlfall, 1990), equine (Tremoleda *et al.*, 2003), buffalo (Abdoon *et al.*, 2001) and caprine (Pawshe *et al.*, 1996). Moreover, gonadotropins improve the ability of mammalian oocytes to form the male pronucleus after IVF (Thibault *et al.*, 1975; Funahashi and Day, 1993) and their developmental competence up to blastocyst stage (Men *et al.*, 2002). In goats, blastocyst formation is also improved by media containing 1 IU/ml FSH reporting a rate of 19.4% to 22.6% compared to a 12% blastocyst formation rate in media without FSH (Wang *et al.*, 2007). Other effects of the presence of gonadotropins during IVM have been observed on gene profile expression of granulosa cells from bovine (Endo *et al.*, 2013) and maternal mRNA levels in oocytes from buffalo (Nath *et al.*, 2013), that have been shown to be gonadotropin (FSH and LH) dependents.

The concentration on gonadotropins used during the process of IVM of oocytes has been showed to determine the results of oocyte maturation and developmental

competence. In bovine, Liu *et al.*, (2011) reported that extremely high concentrations of gonadotropins have detrimental effects on oocyte nuclear maturation and embryo development and increase apoptosis in cumulus cells. In porcine, the use of a low concentration of FSH (1 µg/mL) seems not be sufficient to induce full nuclear maturation of oocytes, compared with 10 µg/mL FSH (Bing *et al.*, 2001). However, and in contrast, some studies have shown that lower concentrations of hormones are enough for IVM of mice oocytes (Peluso, 1988; Byskov *et al.*, 1997). Also in gilt, Wu *et al.*, (2006) have observed a significant increase in blastocyst yield derived from oocytes obtained from small follicles by reducing 250-fold the concentration of the hormones in the protocol of IVM (2-step culture system).

Estradiol-17β is also involved in *in vitro* oocyte maturation in numerous species, including cattle (Sirotkin, 1992), sheep (Guler *et al.*, 2000) and goats (Pawshe *et al.*, 1996). In bovine, the supplementing of IVM medium with 1 µg/mL E₂-17β enhanced oocyte maturational success in some cases (Sirotkin, 1992), but was deleterious in others (Beker-van Woudenberg *et al.*, 2004). In prepubertal goats, supplementation with high concentrations of E₂-17β (10 and 100 µg/mL) during IVM was found to be inhibitory on the progression to metaphase II of oocytes collected from prepubertal goats as compared with 1 µg/mL (Lv *et al.*, 2010).

Results of those studies suggest the importance of judicious use of gonadotropins to enhance *in vitro* oocyte maturation and embryonic development, depending on the dose, specie and the donor age. Gonadotropic concentration commonly added to oocyte maturation of prepubertal and adult goats IVM medium is 10 µg/mL LH, 10 µg/mL FSH and 1 µg/mL E₂-17β (Mogas *et al.*, 1997a; Cognié *et al.*, 2004).

2.1.3.3.2. Antioxidants

The manipulation of gametes and embryos in an *in vitro* environment when performing assisted reproductive techniques carries the risk of exposure of these cells to supraphysiological levels of reactive oxygen species (ROS) (Agarwal *et al.*, 2006), especially when the oxygen concentration employed in most IVM protocols is approximately 20%. When the physiological balance between ROS production and antioxidant defences is lost during exposure to stressful stimuli, oxidative stress

subsequently results in damage to nucleic acids, proteins and lipids despite that they are produced endogenously and derived from external sources (Liu *et al.*, 2003). Studies performed *in vitro* showed that oxidative stress during *in vitro* culture provokes alterations in bovine (Fatehi *et al.*, 2005), murine (Choi *et al.*, 2007), porcine (Tatemoto *et al.*, 2000) and goat (Rodriguez-Gonzalez *et al.*, 2003) oocytes, that impair their maturation and developmental competence.

Within cells, one of the means to control excessive ROS formation is their degradation by antioxidant enzymes. For the past few years, the regulatory role of antioxidants, during oocyte maturation and embryonic development has been a subject of great research interest for their potential use to improve oocyte nuclear and cytoplasmic competence and quality of *in vitro* produced embryos. An antioxidant is any substance that, when present at low concentrations compared to that of an oxidizable substrate can significantly delay or prevent the oxidation process (Halliwell and Gutteridge, 1989). Several previous studies have reported that the presence of non-enzymatic antioxidants, such as β -mercaptoethanol (Choe *et al.*, 2010), vitamin A (Rajesh *et al.*, 2010), vitamin C (Tao *et al.*, 2010), vitamin E (Tareq *et al.*, 2012), cysteine/cysteamine (Zhou *et al.*, 2008), hypotaurine (Mizushima and Fukui, 2001), taurine (Manjunatha *et al.*, 2009) or melatonin (Cebrian-Serrano *et al.*, 2013), enhance embryo development ability or quality after IVC. For instance, improvements in oocyte maturation, embryo development and quality, and embryo survival ability have been observed when oocytes and embryos are treated with these antioxidants in pig (Kitagawa *et al.*, 2004; Hossein *et al.*, 2007), cattle (Olson and Seidel, 2000; Takahashi *et al.*, 2002; Wongsrikeao *et al.*, 2007; Hosseini *et al.*, 2009; Rooke *et al.*, 2012, Korhonen *et al.*, 2012), mouse (Lane *et al.*, 2002; Wang *et al.*, 2002b) and sheep (Miclea *et al.*, 2012). In addition, gene expression profile has been shown to be altered when porcine embryos are cultured with vitamin C (Huang *et al.*, 2011). Likewise, the presence of flavonoids or melatonin in the culture medium has also reported to affect the gene expression profile in bovine (Lee *et al.*, 2011) and mouse (Wang *et al.*, 2013a) embryos. Thiol compounds, such as cysteamine, 2-mercaptoethanol, cysteine, cystin and glutathione (GSH), added to IVM media protect the oocytes from ROS (De Matos *et al.*, 2002). These thiols also increase intracytoplasmic GSH concentration which has a positive effect on fertilization and male pronucleus formation in oocytes from adult (De Matos *et al.*, 2002) and prepubertal (Bai *et al.*, 2008; Rodriguez-Gonzalez *et al.*, 2003) females. In prepubertal

goats, Rodriguez-Gonzalez *et al.*, (2003) showed that the addition of 100 μM of cysteamine to the maturation medium improves embryo development. The relationship between glutathione and development competence has also been previously demonstrated in bovine and sheep oocytes (De Matos *et al.*, 1995; 2002).

One of the most important antioxidant in extracellular fluids, ascorbic acid (vitamin C), plays a key role in many biological processes such as the protection of lipid structures against peroxidation and the biosynthesis of collagen and other components of the extracellular matrix (Buettner, 1993; Rose and Bode, 1993). The antioxidant properties of ascorbic acid are attributed to its capacity to significantly reduce the damage from ROS, forming ascorbate as a stable free radical (Buettner, 1993). Although it is reasonable to hypothesise that ascorbic acid mediates this ROS-decreasing through altering the expression of genes related with antioxidant response. Several works emphasize the antioxidant effect of ascorbic acid in different species. In this way, studies in porcine (Tatemoto *et al.*, 2001) showed that oocytes treated with 250 μM ascorbic acid during IVM displayed enhanced abilities to undergo the male pronucleus formation and to develop to the blastocyst stage after *in vitro* fertilization by prevention of oxidative stress. Similarly, after parthenogenetic activation, ascorbic acid promoted the subsequent development of porcine cumulus-denuded oocytes (Tao *et al.*, 2010), and prevented cumulus cell DNA fragmentation (Tao *et al.*, 2004). Also in mice, ascorbic acid has been shown to exhibit anti-apoptotic effects, when addition of ascorbic acid (0.5 mM) into the culture medium with no serum supplementation significantly reduced the level of apoptosis in the oocyte-associated granulosa cells (Eppig *et al.*, 2000).

2.1.3.3.3. Insulin-Transferrin-Selenium

Insulin is a polypeptide hormone that promotes the uptake of glucose and amino acids and may have mitogenic effects (Spicer and Echternkamp, 1995). Selenium (Se) is an essential trace element. The predominant biochemical action of Se in both humans and animals is to serve as an anti-oxidant, via the Se-dependent enzyme glutathione peroxidase, and thus protect cellular membranes and organelles from peroxidative damage (Moslemi and Tavanbakhsh, 2011). Iron-mediated toxicity has been attributed to iron (Fe), which reacts with oxygen to generate free radicals that damage

macromolecules and cause cell death (Chamnongpol *et al.*, 2002). Transferrin binds iron, making the Fe ions less toxic, but still available to the cell. Thus, transferrin exercises its supportive effect on *in vitro* embryo production via its chelating effects (reviewed by Bowles and Lishman, 1998).

Studies performed *in vitro* reported that the combination of Insulin–transferrin–selenium (ITS) had a positive effect on nuclear and cytoplasmic maturation that make an improvement of the post-fertilization and embryonic development in pig oocytes (Hu *et al.*, 2011). In buffalo, the yield of blastocysts was higher in IVM media containing ITS as compared to control group (Raghu *et al.*, 2002b). In bovine, Córdova *et al.*, (2010) showed that the presence of ITS and ascorbic acid during the first 12h of IVM rendered significantly higher blastocyst rates. Whereas in lamb oocytes the supplementation of maturation medium with ITS and ascorbic acid have shown no effects on the percentage of *in vitro* embryos produced (Catalá *et al.*, 2013). Moreover, the combination of L-ascorbic acid with insulin-transferrin-selenium (ITS) in a growth-supporting maturation medium of gilt oocytes (Wu *et al.*, 2006) has showed to improve the competence of oocytes isolated from small follicles.

2.1.3.3.4. Activin-A

The transforming growth factor-beta (TGF- β) superfamily is a large group of peptide growth and differentiation factors that have important functions in many physiological processes, including reproduction (Bilezikjian *et al.*, 2006). The activins and inhibins are members of the TGF- β superfamily and, along with follistatin, a high affinity binding protein of activin, form a group of interrelated factors originally isolated for their role in regulating the release of follicle-stimulating hormone (FSH) (Phillips, 2005). As described by Knight and Glister (2003) (Figure 1), all three proteins are synthesized and secreted by granulosa cells in an FSH-responsive manner. Local actions of activins include promotion of granulosa cell proliferation, FSH receptor expression, higher levels of P450 aromatase expression and estrogen production, and inhibition of LH-induced androgen production by theca cells. Thus, granulosa cells are likely to be the main source of paracrine factors, and are crucial for oocyte maturation. Follistatin is structurally unrelated to the activins and inhibins, but binds with high affinity to the β -subunits and neutralize the activity of inhibin (Phillips, 2005). The inhibin α -subunit

exerts negative feedback on FSH secretion, and inhibits FSH activity by suppressing its receptor expression in granulosa cells.




Figure 1: Schematic diagram summarizing potential intrafollicular roles of inhibins, activins and follistatin (Knight and Glister, 2003).

Various tissues express protein and mRNA for activin-A and activin receptors, among them the ovary is one of the representative tissues that expresses activin. Likewise, the expression of protein and mRNA for activin-A and activin receptors has been localized in both oocyte and granulosa cells of follicles at various developmental stages of primates (Roberts *et al.*, 1993; Sidis *et al.*, 1998); swine (Van den Hurk and Van de Pavert, 2001); bovine (Gandolfi *et al.*, 1995; Hulshof *et al.*, 1997; Izadyar *et al.*, 1998); ovine (Tisdall *et al.*, 1994; Thomas *et al.*, 2003); and caprine (Silva *et al.*, 2004). The presence of activin receptors on oocytes and granulosa cells suggests that activin and its binding protein, follistatin, may regulate oocyte maturation.

Various studies reported that supplementing IVM media with exogenous activin-A improves oocyte developmental potential. The increase of oocyte maturation has been reported in prepubertal rats (Itoh *et al.*, 1990; Sadatsuki *et al.*, 1993), monkeys (Alak *et al.*, 1996), cows (Stock *et al.*, 1997) and humans (Alak *et al.*, 1998). In bovine, it has been observed a dose dependent effect of activin-A. Thus, Silva and Knight (1998) showed an enhance in the blastocyst rate with a dose of 500 ng/mL activin-A in the IVM medium, whereas Stock *et al.*, (1997) obtained the best blastocyst rate after adding 10 ng/mL to the IVM media. While other studies performed in bovine (Van Tol *et al.*,

1994) and porcine (Coskun and Lin, 1994) oocytes reported no effect of activin-A on oocyte maturation. Also in bovine, IVM in the presence of 10 ng/mL activin-A did not change the proportion of cleaved oocytes that reached the blastocyst stage (Izadyar *et al.*, 1996). The reasons for these differences are likely related to a fundamental difference between culture conditions and size and maturation stage of COCs.

2.1.3.3.5. Serum and other substances

In vitro oocyte maturation (IVM) procedures frequently include supplements of animal origin, such as serum and follicular fluid, and it is considered as non-defined culture medium because of its unknown composition. Their actions are not fully understood, but it is believed that they provide proteins and/or some growth factors that contribute to the success of *in vitro* maturation and subsequent development. Despite the undefined and variable nature of serum composition, the supplementation of maturation media with serum is practiced widely. In goat, maturation media are generally supplemented with 10–20% heat-treated serum, among them: fetal bovine or calf serum (FCS) (Rho *et al.*, 2001; Cognié *et al.*, 2004; Katska-Ksiazkiewicz *et al.*, 2007; Khatun *et al.*, 2011), steer serum (SS) (Jiménez-Macedo *et al.*, 2006; Romaguera *et al.*, 2010b), estrous goat serum (EGS) (Keskin-tepe *et al.*, 1994; Teotia *et al.*, 2001; Baldassarre *et al.*, 2002; Izquierdo *et al.*, 2002), estrous sheep serum (ESS) (Tajik and Esfandabadi, 2003), serum of peritoneal fluid from rabbit and goat (Malik and Lohan, 1999), and bovine serum albumin (BSA)+EGS (Rajikin *et al.*, 1994). The follicular fluid is also used as a supplement in the IVM media in goats and sheep (Cognié *et al.*, 2004). The supplementation of IVM media with follicular fluid from non-atretic or gonadotropin stimulated large follicles (>4 mm) resulted in beneficial effects in both sheep and goat oocytes (reviewed by Tibary *et al.*, 2005). In goat, Mogas *et al.*, (1997a) tested EGS (at different times of estrus), FCS and SS and did not find any significant differences on maturation and embryo production. However, the presence of serum or follicular fluid in culture media introduces a variation from batch to batch thus contributing to the lack of reproducibility often observed in IVEP laboratories. Furthermore, serum and follicular fluid contain many components including hormones, trace elements, and growth factors.

Culture media supplemented with different growth factors have been used to study their potential role on maturation and embryonic development in a growing number of studies. Among the latter, epidermal growth factor (EGF) has been implicated in oocyte maturation and subsequent development in several species. In sheep, *in vitro* matured oocytes in the presence of EGF had greater cumulus cell expansion and higher fertilization rates (Guler *et al.*, 2000) and increases blastocyst formation in FSH-treated ewes (Grazul-Bilska *et al.*, 2003). In goat, the expression of EGF-R both in oocytes and in follicular cells in goat cumulus-oocyte complexes (Gall *et al.*, 2004), suggested that EGF may regulate the oocyte growth and may be involved in nuclear and cytoplasmic maturation. In prepubertal gilt, addition of 10 ng/mL EGF to IVM medium increased the percentage of meiotically matured oocytes (88% vs. 70%) and also the embryos presented a higher number of cells per blastocyst compared with those of control blastocysts (51.1 ± 5.1 % vs 36.0 ± 3.1 %) (Gruppen *et al.*, 1997), showing that EGF plays an important role in both the meiotic and cytoplasmic *in vitro* maturation of porcine oocytes. Another factor to consider is insulin-like growth factor-I (IGF-I) that has also been known to stimulate oocyte maturation and promoting blastocyst development in several species, such as bovine (Matsui *et al.*, 1995), ovine (Guler *et al.*, 2000), porcine (Gruppen *et al.*, 1997) and caprine (Magalhães-Padilha *et al.*, 2012).

Also the positive influence of growth hormone (GH) on oocyte maturation has been well reported. In ovine, Shirazi *et al.*, (2010) showed a positive effect of GH added to *in vitro* maturation medium that resulted in 73% of hatched blastocyst. In bovine, it was observed that GH induced cumulus expansion and promoted subsequent embryonic development in terms of enhancement of the number of cleaved embryos and blastocysts (Izadyar *et al.*, 1996).

A part of hormones and growth factors that are found in the follicular fluid, polyunsaturated fatty acids (PUFAs) constitute the major portion of the fatty acid content of the follicular fluid in small and large follicles (Homa and Brown, 1992). PUFAs concentration in the follicular fluid is associated with the content of the diet and may alter the oocyte maturation process and affect its further development. Actually, studies performed in cattle have shown that *in vitro* supplementation of bovine oocytes with physiological concentrations of α -linolenic acid (ALA; 18:3n-3) or linoleic acid (LA; 18:2n-6) during *in vitro* maturation had diverse effects on maturation and

subsequent embryo development; stimulatory effects were achieved by ALA supplementation (an increased maturation rate and a higher blastocyst yield and production of better quality blastocysts) whereas LA was inhibitory compared with untreated controls (Marei *et al.*, 2009; 2010). However, to our knowledge, no studies of the effect of PUFAs on *in vitro* embryo development in goats have been reported.

2.2. *In vitro* fertilization (IVF)

In vitro fertilization is a complex procedure whose success requires appropriate oocyte maturation, sperm selection, sperm capacitation, and IVF media. In goat, freshly ejaculated semen is usually used for IVF (Cox *et al.*, 1994; Keskindepe *et al.*, 1994; Crozet *et al.*, 1995; Mogas *et al.*, 1997a; Izquierdo *et al.*, 1998; Anguita *et al.*, 2007; Romaguera *et al.*, 2011). Few trials have been described where IVF was carried out using frozen-thawed sperm (Keskindepe *et al.*, 1998; Rho *et al.*, 2001; Bormann *et al.*, 2003).

Under *in vivo* conditions, potentially fertile spermatozoa are separated from immotile spermatozoa, debris and seminal plasma in the female genital tract by active migration through the cervical mucus. So, prior to *in vitro* fertilization, spermatozoa need to be selected and prepared to inseminate the oocytes. The ejaculate comprises of a mixture of seminal plasma, mature and immature spermatozoa, non-reproductive cells, various microorganisms and non-specific debris. In bucks, the most common methods used for separating the sample into motile and non-motile fractions from fresh ejaculate or from the frozen-thawed sperm are the swim-up (Keskindepe *et al.*, 1994; Izquierdo *et al.*, 1998; Katska-Ksiazkiewicz *et al.*, 2004; Jiménez-Macedo *et al.*, 2005; Anguita *et al.*, 2007; Romaguera *et al.*, 2010b) and discontinuous density gradient centrifugation (Pawshe *et al.*, 1996; Rho *et al.*, 2001; Wang *et al.*, 2002a; Bormann *et al.*, 2003), or sephadex filtration (Rho *et al.*, 2001) among others. Greater yields of highly motile spermatozoa can be obtained by swim-up, compared to density gradient centrifugation, but no differences were observed in terms of oocyte penetration and cleavage rate after IVF with fresh goat semen (Palomo *et al.*, 1999).

Once the most viable and motile spermatozoa are selected, sperm capacitation is carried out *in vitro* and using defined media. Capacitation is a crucial process that mammalian

sperm must undergo in order to achieve fertilizing ability. It is defined as the phenomenon leads to "acrosome reaction" causing the release of proteolytic enzymes that may assist sperm penetration into the oocyte. In goat, several capacitating agents have been used to capacitate spermatozoa and to yield good fertilization and cleavage rates, such as heat-inactivated estrous from sheep (De Smedt *et al.*, 1992), goat serum (Koeman *et al.*, 2003; Katska-Ksiazkiewicz *et al.*, 2004), heparin using fresh semen (Izquierdo *et al.*, 1998; Jiménez-Macedo *et al.*, 2005) or frozen-thawed sperm (De Souza *et al.*, 2013), heparin and ionomycin (Wang *et al.*, 2002a; Urdaneta *et al.*, 2004), calcium ionophore (Pereira *et al.*, 2000) and heparin and caffeine (Younis *et al.*, 1991).

Regarding the fertilization media used in goats, these include: modified Defined Media (mDM) (Crozet *et al.*, 1995) Tyrode's Albumin Lactate Pyruvate (TALP) medium (Parrish *et al.*, 1986) supplemented with hypotaurine (Mogas *et al.*, 1997b; Izquierdo *et al.*, 1998), and Synthetic Oviductal Fluid (SOF) medium (Rho *et al.*, 2001). Incubation of sperm selected by swim-up in TALP with heparin (50 mg/mL) for 45 min resulted in excellent fertilization rates (Palomo *et al.*, 1999; Katska *et al.*, 2002; Katska-Ksiazkiewicz *et al.*, 2004). Similarly, mDM plus heparin for sperm capacitation and TALP medium with hypotaurine for oocyte fertilization provided the highest proportion of penetrated oocytes (Izquierdo *et al.*, 1998).

2.3. *In vitro* culture (IVC)

The last stage of *in vitro* embryo production is the culture of the presumptive zygotes in culture media where they undergo a number of divisions until the blastocyst stage 6-7 days after *in vitro* fertilization in ruminant species (Gardner *et al.*, 1994). This period of postfertilization culture is the period having the greatest impact on the blastocyst quality (Rizos *et al.*, 2002a). Several major developmental events take place, including the first cleavage division; the switching on of the embryonic genome; the compaction of the morula and the blastocyst formation. The blastocyst stage involves the differentiation of two types of cells, the inner cell mass (ICM), which after further differentiation gives rise to the fetus, and the trophoctoderm (TE), which ultimately contributes to the placenta formation (Watson, 1992). Clearly, any modifications of the culture environment, which could affect any or all of these processes, could have a major effect

on the quality of the embryo. Some of these factors that could influence on the rate of embryo development and the quality of the embryo are described below.

The *in vitro* oxygen concentration under which embryo development occurs has been found to modify the development up to the blastocyst stage. The oxygen tension that most mammalian embryos encounter in the reproductive tract, range from 3.5 to 8% (Fischer and Bavister, 1993). The positive effect of low oxygen tension on embryo culture has been reported for many species. For instance, the culture of *in vitro* fertilized pig embryos in low oxygen increased the number of cells (Booth *et al.*, 2005) and embryo production rate (Karja *et al.*, 2004) to greater than those cultured in 20% oxygen. In bovine, embryo development and survival rates after cryopreservation were higher when 5% oxygen was applied during IVC compared to 20% oxygen in air (Rizos *et al.*, 2001). Those results suggested that greater O₂ tension during culture is detrimental for embryo development, probably due to the accumulation of reactive oxygen species (ROS).

Different culture systems have been studied to support pre-implantation development of embryos up to the blastocyst stage. Given that female reproductive tract secretions have several amino acids that can be used as energetic substrate by the embryo, the use of amino acids in culture media improves embryo development probably through an antioxidant action and reducing the stress and cell fragmentation caused by *in vitro* embryo culture. On the other hand, in the undefined culture systems, serum is one of the main components. It can provide many beneficial factors to the embryo such as amino acids, vitamins, growth factors, and energetic substrates; however, it may also contaminate the culture media with embryotoxic factors (Bavister, 1995).

Diverse culture media have been successfully used for small ruminant embryo development such as TCM199 (Wani *et al.*, 2012), B2 (Katska-Ksianzkiewicz *et al.*, 2007) and “Sydney IVF Blastocyst” medium (Beilby *et al.*, 2011). However, the most widely used medium is the synthetic oviduct fluid (SOF). Tervit *et al.*, (1972; 1974) were among the first to report successful culture of ruminant zygotes to the blastocyst stage *in vitro* using SOF medium, which was based on the composition of ovine oviduct fluid. Subsequently, changes to the original composition have been made with some modification (Takahashi and First, 1992). Some laboratories routinely supplement SOF medium with serum. Studies in bovine have shown that serum has a biphasic effect; the

presence of serum can inhibit the early cleavage divisions, while it can have an accelerating effect later in development, resulting in the appearance of blastocysts earlier in culture (Lonergan *et al.*, 1999; Gutiérrez-Adán *et al.*, 2001). Thus, some researchers supplement IVC medium with 5-10% FCS (Cognié *et al.*, 2003; Jiménez-Macedo *et al.*, 2005; 2006; Anguita *et al.*, 2007) at 2-3 days post-insemination to promote a higher viability after transfer of such IVP embryos (Cognié, 1999). Other studies only add BSA to SOF media (Leoni *et al.*, 2007; Shabankareh and Akhondi, 2012; Wang *et al.*, 2013b).

Currently, several animal and human studies attest a beneficial effect of addition of growth factors to culture media. However, there is still ambiguity regarding the exact role of growth factors in embryonic development, the optimal dose of growth factors to be added to the culture media, and the combinatorial effect of growth factors in embryonic development (Hegde and Behr, 2012).

Previous research has shown that growth factors such as EGF and IGF-I induced a positive effect on preimplantation development by stimulating metabolism and growth of embryos. In bovine, Palma *et al.*, (1997) demonstrated that culture media containing high concentrations of IGF-I improved the development of embryos produced *in vitro*, and significantly reduced apoptosis (Makarevich and Markkula, 2002). Furthermore, combination of EGF and IGF-I showed an additive effect, and higher rates of embryos developed into blastocysts (Sakagami *et al.*, 2012). In murine embryos, it has been reported that a high dose of EGF resulted in an improvement on blastocyst rate and cell number (Głabowski *et al.*, 2005). Similarly, Mtango *et al.*, (2003) demonstrated that the addition of growth factors and growth hormone (GH) to the culture media of bovine embryos had favorable effects on *in vitro* embryo development, freezing sensitivity and post-thawing survival, hatching rate and total cell number of blastocysts. Other growth factors may also have beneficial effects on embryo development. Regarding the use of hormones, mouse blastocysts that have been cultured with growth hormone showed a higher chance for implantation after transfer to the recipient (Fukaya *et al.*, 1998). Moreover, ghrelin is a widespread hormone that several studies have linked with reproductive physiology (Garcia-Garcia *et al.*, 2007; Tena-Sempere, 2008). In sheep (Wang *et al.*, 2013b), the blastocyst rate, total cell number of blastocysts and the expression levels of the GLUT1 and IFNT genes were increased when 50 ng/mL ghrelin was added during IVC to the SOF medium.

Since activin-A is produced by oviduct epithelial cells (Gandolfi *et al.*, 1995), the addition of activin-A to embryo *in vitro* culture may reproduce the environment of the cleavage-stage embryos in the oviduct *in vivo*. Activin-A and activin receptors have been detected in mouse (Lu *et al.*, 1993) and bovine (Yoshioka *et al.*, 1998b) embryos from zygote to the morula stage, suggesting that the protein might play a role in embryogenesis. According to studies performed *in vitro*, activin-A has been shown to have an effect in regulation of the development of bovine embryos (Yoshioka and Kamomae, 1996; Yoshioka *et al.*, 1998a). However, the effect of activin-A on *in vitro* embryo culture is still controversial. That depending on the timing of its addition to the culture medium, exogenous activin-A may promote the morula and blastocyst development when the protein was added at the earliest stages of bovine (Yoshioka and Kamomae, 1996; Yoshioka *et al.*, 1998a; Lee *et al.*, 2009) embryo culture. Similar beneficial effects of activin, when added to the second half of the embryo culture, were reported as increased expanding and hatching rates (Park *et al.*, 2010; Trigal *et al.*, 2011). As described by Park *et al.*, (2010), activin-A, during the later stage of *in vitro* bovine embryo development, can enhance *in vitro* development of embryos by affecting expression levels of genes related to hatching and implantation in defined culture medium, and the development-enhancing effect of activin-A may be associated with timing of activation of appropriate genes within the embryonic genome (Trigal *et al.*, 2011). Also, maternal levels of activin protein in the mouse embryo decline during early cleavage stages but increase, presumably due to embryo transcription, in the compacted morula (Albano *et al.*, 1993). It has been also shown a dose effect when activin-A was supplemented to *in vitro* culture media of IVM/IVF bovine oocytes. Yoshioka *et al.*, (1998a) observed that the percentages of zygotes that developed to the morula and blastocyst stages increased dose-dependently with addition of activin-A, similar to the percentages of expanded blastocysts. Therefore, it may be advisable to test what culture system or culture medium will provide better results for a given lab.

2.4. Assessment of embryo quality

A proper evaluation of blastocyst quality remains an important challenge for every researcher involved in embryology to select the best embryos for transfer, chance to obtain pregnancies and to define its sanitary condition.

The noninvasive methods used for evaluating the embryo quality are the timing of early cleavage (Van Soom *et al.*, 1992), understanding that good quality embryos must exhibit appropriate kinetics and synchrony of division, the evaluation of the timing of blastocyst formation (Van Soom *et al.*, 1997) and the blastocyst morphology (Van Soom *et al.*, 2003). All of those methods are practical to perform and without any consequence for the embryo's viability.

The cryopreservation of embryos is an essential method in assisted reproduction technology (ART) for world-wide embryo transfer (ET) technology that also allows to select the most robust and highest quality embryos will be able to support the freeze-thaw process. Therefore, the ability of embryos to survive cryopreservation is commonly used as an indicator of embryo quality and viability (Rizos *et al.*, 2008). Studies performed in several species reported differences on survival rates and hatching percentages after the cryopreservation of blastocysts depending on the origin of embryos (*in vivo* versus *in vitro*), the stage of embryo development, the age of the donor female, and media for IVEP.

Additionally, research on animal models has demonstrated that stress, and the resultant adaptation to conditions during pre-implantation stages, can affect pregnancy loss and fetal growth. It is therefore important to understand the role of each medium component and to identify possible sources of cellular stress to the embryo that will ultimately affect the function and viability of the fetus.

Serum supplementation has been associated with accumulation of cytoplasmic lipid droplets in IVP embryos (Abe *et al.*, 1999) and can lead to poor cryosurvival (Leibo and Loskutoff, 1993). In ovine, Shirazi *et al.*, (2012) showed that a significantly lower survival rate of vitrified-warmed blastocysts derived from oocytes matured in presence of FCS, might be due to the excess accumulation of cytoplasmic lipid droplets in resulting blastocysts compared to the other group. In bovine, Moore *et al.*, (2007) used a serum replacer (SR) for maturation of oocytes and results showed an improved embryo development *in vitro* as well as better cryotolerance and survival when replacing serum in culture. In the same way, the presence of somatic cells during the IVC was shown to affect the freezability of *in vitro* produced bovine embryos (Gómez *et al.*, 2008). Shehab-El-Deen *et al.*, (2009) demonstrated that the addition of elevated concentrations of saturated fatty acids (palmitic and stearic acid) during IVM of bovine oocyte have

deleterious on embryo quality, featured by a significantly reduced post-warming survival of vitrified blastocysts.

In research, the quality of embryos obtained can be also assessed with invasive techniques. Invasive assessment of embryo quality mostly involves a kind of fixation and staining of the embryo. The assessment of embryo quality is carried out under a stereomicroscope on grounds of morphological criteria. Many measures can be investigated to determine embryo quality such as total cell number, the ICM/TE ratio of blastocysts, the degree of cell fragmentation, and determination of genes related with embryo quality and ability to achieve pregnancy after transfer. A simplified technique for differential staining of inner cell mass and trophectoderm cells of mouse and bovine blastocysts (Thouas *et al.*, 2001) permits the differentiation of the two cell populations as the blue ICM cell nuclei (stained only with Hoechst) that can be clearly distinguished from the pink TE cell nuclei (stained both with propidium iodide and Hoechst). This technique has been applied to embryos from different species, including bovine (Iwasaki *et al.*, 1990; Fouladi-Nashta *et al.*, 2005), ovine (Catalá *et al.*, 2011), porcine (Dang-Nguyen *et al.*, 2011) and buffalo (Selokar *et al.*, 2012) among others. Several factors have been reported to affect the proportion of cell number in each blastocyst compartment. If compare *in vivo* versus *in vitro* produced embryos, it has been reported that the *in vitro* production systems affect the blastocyst quality in terms of total cell number, incidence of cell apoptosis, oxidative stress and up and down regulation of gene expression. In bovine, Knijn *et al.*, (2003) reported a significantly lower total cell number for those embryos produced completely *in vitro* compared to that of embryos developed entirely *in vivo*, as well as a higher incidence of DNA fragmentation.

Until the blastocyst stage of development is attained, the embryo is somewhat autonomous (*i.e.*, does not need contact with the maternal reproductive tract) as evidenced by the fact that blastocysts can be successfully developed *in vitro* in large number using IVF technology and transferred to reproductive synchronized recipients (Mamo *et al.*, 2012). However, the ability to undergo implantation and pregnancies depends on blastocyst quality. Thus, cell number and apoptosis levels are also important parameters that are emerging as useful indicators of embryo development and health (Brison and Schultz, 1997; De la Fuente and King, 1997). This was further supported by Fouladi-Nashta *et al.*, (2005) who demonstrated that counting the total number of blastomeres and the percentage of apoptotic nuclei provides more detailed information

on embryo quality and developmental potential. These authors investigated the effect of serum supplementation on total cell number, ICM/TE ratio and apoptosis index after *in vitro* production of bovine embryos. Results of this study showed that supplementation of serum will possibly increase the number of TE cells leading a reduced ratio of ICM/TE. The high number of TE cells in the serum treated embryos may cause the increased placenta size in the large offspring syndrome.

Another parameter used to assess the quality of embryos is the ratio of programmed cell death or DNA fragmentation. Cell death is present in mammalian embryos, which might be involved in the elimination of abnormal cells and its normal pattern is crucial for further development necessary to maintain the morphogenesis and to allow cell renovation. Cell death was confirmed in goat blastocysts using the terminal deoxynucleotidyl transferase (TdT) mediated dUTP nick-end labeling (TUNEL) assay (Jiménez-Macedo *et al.*, 2007). TUNEL allows in situ assessment of DNA breaks in the nuclei (reviewed by Neuber *et al.*, 2002) and it has been observed a negative relation with the implantation rates.

Likewise, oxygen free radicals can cause DNA strand breaks and other forms of cellular damage (lipid peroxidation) which are known to result in cell cycle arrest or death. Although apoptosis can be considered to be a normal process in preimplantation embryos to eliminate deviating cells, a high incidence of apoptotic cells is correlated with abnormal morphology of the embryo. Moreover, embryonic cell apoptosis has received increasing attention, mostly due to its suggested role in the cellular response to suboptimal developmental conditions (Betts and King, 2001). Thus, increased apoptosis is an important indicator of inadequate *in vitro* conditions for eggs.

Ending, environmentally induced changes in oocyte and embryo biology have both immediate effects on the viability and development of these structures, but can also have pronounced and often persistent effects on the resultant offspring. For all above mentioned and keeping on mind that the ultimate test of the quality of an embryo is its ability to produce live and healthy offspring after transfer to a recipient, it is really important consider and optimize the conditions of the *in vitro* culture systems in order to get embryos able to undergo pregnancies.

O

BJECTIVES

Objectives of the study

The overall aim of this thesis work was to improve the *in vitro* embryo production from oocytes of prepubertal goats. More specifically, the aims were:

I: To examine the effect of supplementing *in vitro* maturation medium with low level of hormones and/or insulin-transferrin-selenium plus ascorbic acid in the developmental competence and embryo quality of small oocytes from prepubertal goats.

II: To evaluate the effect of adding activin-A at the *in vitro* oocytes maturation and *in vitro* embryo culture media on meiotic maturation, embryo development and embryo quality of *in vitro*-produced blastocysts.

Developmental competence and embryo quality of small oocytes from pre-pubertal goats cultured in IVM medium supplemented with low level of hormones, insulin-transferrin-selenium and ascorbic acid

Hammami S¹, Morató R², Romaguera R¹, Roura M¹, Catalá MG¹, Paramio MT¹, Mogas T², Izquierdo D¹.

¹ Departament de Ciència Animal i dels Aliments, Universitat Autònoma de Barcelona, Bellaterra, Spain

² Departament de Medicina i Cirurgia Animals, Universitat Autònoma de Barcelona, Bellaterra, Spain

Reprod Domest Anim. 48 (2013) 339-344.

Developmental Competence and Embryo Quality of Small Oocytes from Pre-pubertal Goats Cultured in IVM Medium Supplemented with Low Level of Hormones, Insulin–Transferrin–Selenium and Ascorbic Acid

S Hammami, R Morató, R Romaguera, M Roura, MG Catalá, MT Paramió, T Mogas and D Izquierdo

Facultat de Veterinària, Universitat Autònoma de Barcelona, Bellaterra, Spain

Contents

The aim of this study was to test the effect of insulin–transferrin–selenium (ITS) and L-ascorbic acid (AA) supplementation and the hormonal level during *in vitro* maturation (IVM) of small oocytes from pre-pubertal goat on the blastocyst yield and quality. Concretely, we used four maturation media: conventional IVM medium (CM), growth medium (GM: CM+ITS+AA and low level of hormones), modified CM (mCM: CM with low level of hormones) and modified GM (mGM: CM+ITS+AA and normal level of hormones). Cumulus–oocyte complexes (COCs) were classified into two categories according to oocyte diameter: <125 µm and ≥125 µm. Large oocytes were matured 24 h in CM (Treatment A). Small oocytes were matured randomly in six experimental groups: Treatment B: 24 h in CM; Treatment C: 12 h in GM and 12 h in CM; Treatment D: 24 h in mGM; Treatment E: 12 h in mGM and 12 h in CM; Treatment F: 12 h in mCM and 12 h in CM; and Treatment G: 12 h in GM and 12 h in mGM. After IVM, oocytes were fertilized and cultured for 8 days. The blastocyst quality was assessed by the survival following vitrification/warming and the mean cell number. When different maturation media were combined, the blastocyst rate did not improve. The large oocytes produced the highest blastocysts yield. However, the culture of small oocytes in GM (53.3%) enhanced the post-warming survival of blastocysts compared to large oocytes matured in CM (35.7%). In conclusion, IVM of pre-pubertal goat small oocytes in GM would be useful to improve the quality of *in vitro*-produced blastocysts.

Introduction

In vitro embryo production using oocytes from pre-pubertal females has been studied for years and has reported a low embryo development (reviewed by Armstrong 2001). In pre-pubertal goats, oocytes smaller than 125 µm were unable to develop into blastocysts after *in vitro* fertilization (IVF) (Anguita et al. 2007) and ICSI (Jimenez-Macedo et al. 2006) procedures. This poor development of *in vitro*-derived zygotes suggests that culture conditions during *in vitro* maturation (IVM) of small oocytes do not allow for a correct cytoplasmic maturation and normal embryo development after fertilization. Thus, over the last few years, several systems supporting IVM of oocytes have been studied to improve the ability of the culture system to stimulate embryonic development and/or blastocyst quality [bovine (Gomez et al. 2008); ovine (Shirazi et al. 2010)]. In pig, Wu et al. (2006) have defined a growth medium (GM) for culturing oocytes derived from small follicles of pre-pubertal gilt ovaries. This medium has been considered as a 'growth-supporting and less maturation-promoting environment', and it is used before the

culture in conventional IVM medium, achieving a blastocyst yield comparable to full-growth oocytes. Comparing this to the conventional IVM medium, the growth medium has a low level of hormones and is supplemented with the combination of insulin–transferrin–selenium (ITS) and L-ascorbic acid. In mice, some studies show that oocytes from pre-antral follicles can complete their growth *in vitro* with low concentrations of hormones (Peluso 1988; Eppig and Schroeder 1989). Moreover, insulin, a polypeptide hormone that promotes the uptake of glucose and amino acids (Lee et al. 2005), transferrin and selenium are essential trace elements and have antioxidant activity (Tatemoto et al. 2004). Thus, ITS supplementation has been routinely used in several IVM systems, including mouse (De la Fuente et al. 1999), bovine (Shamsuddin et al. 1993) and porcine (Wu et al. 2006; Jeong et al. 2008). Finally, ascorbic acid has a vital role as an antioxidant, and Tatemoto et al. (2001) concluded that a critical intracellular concentration of ascorbic acid inside the oocytes would be necessary for their normal cytoplasmic maturation and their embryo developmental competence.

Thus, to improve the *in vitro* embryo production from small pre-pubertal goat oocytes, the objectives of the present study are the following: (i) to examine the effect of GM previous maturation in conventional IVM medium to increase blastocyst rate, (ii) to investigate the effect of GM on blastocyst quality and (iii) to test the effect of adding to the maturation medium a low level of hormones and/or insulin–transferrin–selenium (ITS) and ascorbic acid in a sequential maturation process on embryo development to the blastocyst stage.

Material and Methods

Reagents

All chemicals and media were purchased from Sigma Chemical Co. (St. Louis, MO, USA) unless otherwise indicated.

Oocyte collection, classification and IVM

Pre-pubertal goat (1–2 months old) ovaries were obtained from a slaughterhouse and transported to the laboratory in phosphate-buffered saline (PBS) at 38°C. Cumulus–oocyte complexes (COCs) were recovered by the slicing technique. The COCs with a compact cumulus and homogeneous cytoplasm were selected for IVM and classified into two categories according to their oocyte diameter: <125 µm and ≥125 µm.

The COCs were washed three times in maturation medium. Groups of 25–30 COCs were transferred into 100- μ l drops of IVM medium covered with mineral oil and incubated at 38.5°C and 5% CO₂ in humidified air. The basic medium used for IVM was TCM199 supplemented with 275 μ g/ml sodium pyruvate, 146 μ g/ml L-glutamine, 50 μ g/ml gentamycin, 10% (v/v) donor bovine serum (CanSera; Ontario, Canada) and 100 μ M cysteamine. Four types of IVM media were used: (i) basic medium supplemented with 10 μ g/ml follicle stimulating hormone (FSH), 10 μ g/ml LH, 1 μ g/ml and 17 β -oestradiol (conventional IVM medium, CM), (ii) basic medium supplemented with 0.04 μ g/ml FSH, 0.04 μ g/ml LH, 0.004 μ g/ml 17 β -oestradiol, 100 μ g/ml ascorbic acid and 5 μ l/ml ITS (growth medium, GM), (iii) basic medium supplemented with 0.04 μ g/ml FSH, 0.04 μ g/ml LH and 0.004 μ g/ml 17 β -oestradiol (modified conventional IVM medium, mCM) and (iv) basic medium supplemented with 10 μ g/ml FSH, 10 μ g/ml LH, 1 μ g/ml 17 β -oestradiol, 100 μ g/ml ascorbic acid and 5 μ l/ml ITS (modified growth medium, mGM).

***In vitro* fertilization (IVF) and *in vitro* embryo culture (IVC)**

After IVM, 25–30 oocytes were transferred to 100- μ l microdrops of modified Tyrode's medium (TALP) covered with mineral oil and cocultured with fresh goat sperm in a final concentration of 4×10^6 motile sperm/ml at 38.5°C and 5% CO₂ in humidified air, as described by Romaguera et al. (2010). At 24 h post-insemination (pi), groups of 10–15 presumptive embryos were placed into 20- μ l droplets of synthetic oviductal fluid medium (Takahashi and First 1992) [SOF medium supplemented with 10% foetal calf serum (FCS)] under mineral oil in a humidified atmosphere with 5% CO₂, 5% O₂ and 90% N₂ for 8 days. Cleavage rate was evaluated at 48 h pi, and blastocyst rates were recorded at Days 7, 8 and 9 pi.

Evaluation of blastocyst quality

Blastocyst vitrification and warming

Blastocyst vitrification was carried out using the cryotop method described by Kuwayama et al. (2005). All warming steps were performed as described by Morato et al. (2010). Briefly, the holding medium (HM) for formulating all vitrification-warming solutions consisted of TCM 199 Hepes buffered with 20% FCS. The blastocysts were transferred into equilibration solution (ES) consisting of 7.5% ethylene glycol (EG) and 7.5% dimethylsulphoxide (DMSO) in HM for 10–15 min. Then, the blastocysts were transferred to the vitrification solution (VS) containing 15% DMSO, 15% EG and 0.5 M sucrose dissolved in HM. After incubating for 30–40 s, the blastocysts were loaded onto the cryotop and the sample was quickly plunged into liquid nitrogen. The entire process from exposure to VS to plunging was completed within 1 min. For warming, the cryotop was directly immersed into the warming solution containing 1 M sucrose dissolved in HM. After 1 min, the blastocysts were transferred to the dilution solution, which

contained 0.5 M sucrose dissolved in HM for 3 min. Next, the blastocysts were incubated in HM for 5 min. Finally, the blastocysts were transferred to the SOF medium and incubated at 38.5°C in a 5% CO₂, 5% O₂ and 90% N₂. After vitrification and warming, survival rates were assessed according to the extent of re-expansion of the blastocysts after 3, 24 and 72 h of recovery.

Hoechst staining

To account the cell number, the blastocysts were stained with 1 mg/ml Hoechst 33342 (H3570; Invitrogen) for 5 min at 38°C. Cell count was conducted under a fluorescence microscope.

Experimental design

Effect of growth medium during IVM on subsequent embryo development of pre-pubertal goat small oocytes

The objective was to test the effect of culturing small oocytes from pre-pubertal goat on GM previous to maturation in CM on blastocyst yield. The COCs were classified into two categories based on oocyte diameter: <125 μ m and \geq 125 μ m. The \geq 125- μ m oocytes were matured in CM for 24 h (Treatment A). The <125- μ m oocytes were distributed into two experimental groups: COCs matured in CM (Treatment B) and COCs cultured in GM for 12 h followed by another 12 h in CM (Treatment C). After IVM, all COCs were fertilized and cultured *in vitro*. The cleavage and blastocyst rates were assessed at Days 2, 7, 8 and 9 post-insemination.

Effect of growth medium during IVM on the tolerance to blastocyst vitrification and blastocyst cell number

The objective of this study was to examine the effect of GM during IVM on blastocyst quality measured by survival following vitrification/warming and blastocyst cell number. The experimental groups were the same as those in Experiment 1 (Treatments A, B and C). The blastocysts were vitrified at Day 8 post-insemination and cultured for 72 h after warming to examine their viability *in vitro*. Survival rate was defined as post-warming blastocyst re-expansion and its maintenance in SOF medium for 3, 24 and 72 h. The blastocyst cell number was evaluated at the end of the culture period (Day 9 pi).

Effect of different supplements in the maturation medium on pre-pubertal goat embryo development according to oocyte diameter

In this experiment, the effect of ITS, ascorbic acid and hormonal levels was evaluated to optimize embryo development of small oocytes from pre-pubertal goats. The experimental groups are described in Table 1. The cleavage rate was assessed at 48 h pi. At Days 7, 8 and 9, blastocysts were evaluated and classified according to the extent of blastocoele expansion into one of four groups as follows: (i) non-expanded or early, (ii) expanded, (iii) hatching and (iv) hatched blastocysts.

Table 1. Description of different IVM treatments in terms of oocyte diameter, level of hormones and ITS and ascorbic acid supplementation

Oocyte diameter	Maturation treatments	Time of IVM (24 h)	
		First 12 h	Following 12 h
≥125 µm	A (CM)	Normal level of hormones	Normal level of hormones
< 125 µm	B (CM)	Normal level of hormones	Normal level of hormones
< 125 µm	C (GM/CM)	Low level of hormones ITS and ascorbic acid	Normal level of hormones
< 125 µm	D (mGM)	Normal level of hormones ITS and ascorbic acid	Normal level of hormones ITS and ascorbic acid
< 125 µm	E (mGM/CM)	Normal level of hormones ITS and ascorbic acid	Normal level of hormones
< 125 µm	F (mCM/CM)	Low level of hormones	Normal level of hormones
< 125 µm	G (GM/mGM)	Low level of hormones ITS and ascorbic acid	Normal level of hormones ITS and ascorbic acid

CM, Conventional Medium; GM, Growth Medium; mCM, modified Conventional Medium; mGM, modified Growth Medium; ITS, insulin–transferrin–selenium; IVM, *in vitro* maturation.

Table 2. Effect of Growth Medium on cleavage and blastocyst development of pre-pubertal goats according to oocyte diameter

Oocyte diameter	Maturation treatments	Oocytes, n	Cleavage, n (%)	Blastocysts, n (%)		
				Day 7	Day 8	Day 9
≥125 µm	A (CM)	305	171 (56.06)	36 (11.80)	53 (17.37) ^a	63 (20.65) ^a
< 125 µm	B (CM)	298	114 (38.25)	20 (06.71)	28 (09.39) ^b	31 (10.40) ^b
< 125 µm	C (GM/CM)	309	128 (41.42)	24 (07.76)	28 (09.06) ^b	30 (09.70) ^b
	p value		0.052	0.1826	0.0057	0.0002

Values with different letters within each column differ significantly ($p < 0.05$).
CM, Conventional Medium; GM, Growth Medium.

Statistical analysis

Data were analysed using the Statistical Analysis Systems package (SAS, v9.1: NC, USA). Cleavage rate, blastocysts rate, blastocysts stage and blastocyst cell number were analysed by the Turkey's test. Blastocysts survival rates were analysed by chi-square analysis. For all the statistical analyses, differences were considered significant when $p < 0.05$.

Results

Experiment 1

The results of cleavage rate and embryo development to blastocyst stage of pre-pubertal goat oocytes in relation to their oocyte diameter and maturation medium are presented in Table 2. At 48 h pi, although the oocytes with a diameter ≥125 µm presented a cleavage rate higher than did small oocytes (<125 µm), no statistical differences were observed among the three treatments (56.06%, 38.25% and 41.42%; Treatments A, B and C, respectively). The yield of blastocysts was significantly higher ($p < 0.05$) in the group of large oocytes than in those of small oocytes. There was no difference between blastocyst rate derived from Treatment C and Treatment B of small oocytes.

Experiment 2

The post-warming viability of the vitrified blastocysts and the total cell number of Day 9 blastocysts are shown in Table 3. At 3 h post-warming, a significant increase in the survival rate was observed on the blastocysts derived from small oocytes matured in CM (76.5%), as

compared to those obtained from the large oocytes (64.1%). There were no differences in survival rates at 24 h after warming. The post-warming viability at 72 h was significantly higher in the blastocysts derived from small oocytes matured in GM than in the blastocysts produced in CM, regardless of the size of the oocyte (35.7%, 30.8% and 53.3%; Treatments A, B and C, respectively). The cell number of blastocysts obtained from ≥125-µm oocytes was significantly higher than that obtained from <125-µm oocytes.

Experiment 3

Table 4 shows the effect of different maturation culture conditions on embryo development of pre-pubertal goat oocyte. No significant differences were observed on cleavage rate among the seven treatments. When blastocyst yield was determined on Day 7 and Day 8, there were no significant differences among the large oocytes (Treatment A) and the small oocytes from Treatments B (CM) and F (mCM/CM), but on Day 9, the blastocyst rate from ≥125-µm oocytes was significantly higher than that obtained with the six treatments of <125-µm oocytes. With respect to the small oocytes, on Days 8 and 9, the blastocyst rates in Treatment D (mGM) (6.22% and 7.05%, respectively) and E (mGM/CM) (7.23% and 7.66%, respectively) were significantly lower than those of Treatment B (11.20% and 13.13%, respectively) and Treatment F (12.80% and 14.40%, respectively).

When data were analysed by day and blastocyst stage (Table 5), the proportions of early, expanded, hatching and hatched blastocysts on Days 7 and 9 were not

Table 3. Cryotolerance assessed by blastocoele re-expansion at 3, 24 and 72 h post-warming and mean cell number of pre-pubertal goat blastocysts according to IVM treatment and oocyte diameter

Oocyte diameter	Maturation treatments	Cryotolerance				Mean cell number	
		Blastocysts, n	% Re-expansion after warming			Blastocysts, n	Blastocyst mean cell number (Mean \pm SEM)
			3 h	24 h	72 h		
$\geq 125 \mu\text{m}$	A (CM)	39	64.1 ^a	56.4	35.7 ^a	21	315.71 \pm 24.14 ^a
< 125 μm	B (CM)	17	76.5 ^b	58.8	30.8 ^a	20	243.45 \pm 24.38 ^b
< 125 μm	C (GM/CM)	19	68.4 ^{ab}	57.9	53.3 ^b	19	207.53 \pm 26.91 ^b

Values in the same column with different letters differ significantly ($p < 0.05$).
CM, Conventional Medium; GM, Growth Medium.

Table 4. Effect of different supplements in the maturation medium on pre-pubertal goat embryo development according to oocyte diameter

Oocyte diameter	Maturation treatments	Oocytes, n	Cleavage, n (%)	Blastocyst, n (%)		
				Day 7	Day 8	Day 9
				$\geq 125 \mu\text{m}$	A (CM)	294
< 125 μm	B (CM)	259	105 (40.54)	24 (9.27) ^{ab}	29 (11.20) ^{abc}	34 (13.13) ^b
< 125 μm	C (GM/CM)	254	105 (41.34)	21 (8.27) ^b	26 (10.24) ^{bcd}	29 (11.42) ^{bc}
< 125 μm	D (mGM)	241	88 (36.51)	12 (4.98) ^{cd}	15 (6.22) ^d	17 (7.05) ^c
< 125 μm	E (mGM/CM)	235	102 (43.40)	9 (3.83) ^d	17 (7.23) ^{cd}	18 (7.66) ^c
< 125 μm	F (mCM/CM)	250	108 (43.20)	23 (9.20) ^{abc}	32 (12.80) ^{ab}	36 (14.40) ^b
< 125 μm	G (GM/mGM)	182	67 (36.81)	10 (5.49) ^b	14 (7.69) ^{bcd}	18 (9.89) ^{bc}
	p value		0.0701	0.0011	0.010	0.003

Values in the same column with different letters differ significantly ($p < 0.05$).

CM, Conventional Medium; GM, Growth Medium; mCM, modified Conventional Medium; mGM, modified Growth Medium.

significantly different among treatments. On Day 8, significant differences were found in expanded blastocyst rates between large oocytes (13.73%), which attained the maximum of expanded blastocysts 1 day before, and the small oocytes from Treatments B (CM), C (GM/CM), F (mCM/CM) and G (GM/mGM) (51.72%, 50.00%, 43.75% and 42.86%, respectively). The expanded blastocyst rate derived from Treatment D (mGM) (26.67%) was significantly lower than that of Treatment B and Treatment C.

Discussion

We hypothesized that the presence of low hormone levels, ascorbic acid and ITS (GM) during the first 12 h of IVM of pre-pubertal goat small oocytes could positively influence the *in vitro* embryo development. However, our results showed that the use of GM did not improve the blastocyst percentage, when compared to those matured in conventional IVM medium. In ovine, Catala et al. (2010) also observed that GM failed to significantly increase embryo production of lamb oocytes. These discrepancies with the findings of Wu et al. (2006) on gilt oocytes may be related to variability among biological requirements of immature oocytes depending on the species used.

In relation to blastocyst quality, the ability of blastocysts derived from $\geq 125\text{-}\mu\text{m}$ oocytes to survive the vitrification/warming process at 3 h is lower than that of blastocysts obtained from small oocytes. At Day 8, the decrease in the cryosurvival rate may be explained by the developmental stage of blastocysts derived from large oocytes (hatching and hatched), as compared to the stage

of development of the blastocysts obtained from small oocytes groups (early and expanded). Our findings are in agreement with previous data in bovine (Morato et al. 2010), suggesting that these differences in survival rates of *in vitro*-produced blastocysts depend on their developmental stage. In this study, we could not avoid the heterogeneity of blastocyst stages because of the small number of blastocysts obtained in each one of the replicates. At 72 h post-warming, the re-expansion rate was significantly higher in the blastocysts obtained from small oocytes matured during the first 12 h in GM (Treatment C) than in the blastocysts from oocytes IVM in conventional medium, reflecting a higher oocyte quality. Previous studies in bovine showed that the culture medium supplementation (Rizos et al. 2001; Gomez et al. 2008) affects intrinsic embryo quality in terms of cryosurvival. On the other hand, with respect to the blastocyst total cell number on Day 8, it was found that blastocysts produced from the large oocytes had the highest cell number. As reported by Jiang et al. (1992), the cell number of bovine blastocysts varied depending on the morphological grade and that the later developing blastocysts were of poor quality as judged by the cell number. The total blastocyst cell number did not vary significantly between the small oocytes treatments. Thus, growth medium did not significantly influence blastocyst cell number of small pre-pubertal goat oocytes.

Based on the previous results, we studied the effects of different growth medium components' combinations (low level of hormones, insulin–transferrin–selenium and ascorbic acid) on the cleavage and developmental capacity of pre-pubertal goat oocytes. We found that embryo development of these oocytes is more related

Table 5. Effect of different supplements in the maturation medium on pre-pubertal goat blastocyst stage according to oocyte diameter

Oocyte diameter	Maturation treatments	Day 7				Day 8				Day 9			
		Early n (%)	Expanded n (%)	Hatching n (%)	Hatched n (%)	Early n (%)	Expanded n (%)	Hatching n (%)	Hatched n (%)	Early n (%)	Expanded n (%)	Hatching n (%)	Hatched n (%)
≥125 µm	A (CM)	11/41 (26.83)	18/41 (43.90)	3/41 (07.32)	9/41 (21.95)	7/51 (13.73)	7/51 (13.73) ^c	29/51 (56.86)	2/62 (03.23)	5/62 (08.06)	4/62 (06.45)	51/62 (82.26)	
<125 µm	B (CM)	10/24 (41.67)	7/24 (29.17)	5/24 (20.83)	2/24 (08.33)	2/29 (06.90)	15/29 (51.72) ^a	6/29 (20.69)	0/34 (00.00)	3/34 (08.82)	3/34 (08.82)	28/34 (82.35)	
<125 µm	C (GM/CM)	10/21 (47.62)	6/21 (28.57)	3/21 (14.29)	2/21 (09.52)	2/26 (07.69)	13/26 (50.00) ^d	5/26 (19.23)	0/29 (00.00)	4/29 (13.79)	3/29 (10.34)	22/29 (75.86)	
<125 µm	D (mGM)	3/12 (25.00)	5/12 (41.66)	3/12 (25.00)	1/12 (08.33)	2/15 (13.33)	4/15 (26.67) ^{bc}	7/15 (46.67)	0/17 (00.00)	1/17 (05.88)	1/17 (05.88)	15/17 (88.24)	
<125 µm	E (mGM/CM)	3/10 (30.00)	4/10 (40.00)	1/10 (10.00)	2/10 (20.00)	1/17 (05.88)	7/17 (41.18) ^{abc}	6/17 (35.29)	0/18 (00.00)	2/18 (11.11)	3/18 (16.67)	13/18 (72.22)	
<125 µm	F (mCM/CM)	7/23 (30.43)	9/23 (39.13)	5/23 (21.74)	2/23 (08.70)	3/32 (09.37)	14/32 (43.75) ^{ab}	8/32 (25.00)	1/36 (02.78)	4/36 (11.11)	7/37 (19.44)	24/36 (66.67)	
<125 µm	G (GM/mGM)	4/10 (40.00)	4/40 (40.00)	1/10 (10.00)	1/10 (10.00)	1/14 (07.14)	6/14 (42.86) ^{ab}	5/14 (35.71)	0/18 (00.00)	2/18 (11.11)	4/18 (22.22)	12/18 (66.67)	

Values in the same column with different letters differ significantly ($p < 0.05$).

CM, Conventional Medium; GM, Growth Medium; mGM, modified Conventional Medium; mGM, modified Growth Medium.

to oocyte diameter than to the conditions of IVM medium culture, which confirms our previous results (Jimenez-Macedo et al. 2006; Anguita et al. 2007). In addition, in oocytes <125 µm, the supplementation of conventional medium with ascorbic acid and ITS for 12 and 24 h in the presence of the normal level of hormones (Treatments E and D) significantly decreased the percentage of blastocysts, when compared to the conventional IVM medium (7.66% and 7.05% vs 13.13%, respectively). This result indicates a detrimental effect of ITS and ascorbic acid added to conventional medium. In calf oocytes, Cordova et al. (2010) observed that the presence of ascorbic acid significantly reduced the proportion of embryos developing to the blastocyst stage. In contrast, other authors have shown the beneficial effects of ITS on the nuclear and cytoplasmic maturation of pig oocytes (Jeong et al. 2008) and on the yield of blastocysts in buffalo (Raghu et al. 2002). On the other hand, in the absence of supplementation with ITS and ascorbic acid, no differences were observed in cleavage and blastocyst rates using low or normal levels of hormones during the first 12 h of IVM. The growth of mouse oocytes in a medium without FSH has been demonstrated, and these oocytes acquire competence to undergo maturation, fertilization and embryonic development (Peluso 1988; Eppig and Schroeder 1989). Also, the addition of LH to the medium has been shown to have no effect on the growth of mouse follicles (Murray et al. 2008). These studies suggest that gonadotrophins may not be required for the acquisition of full development competence by growing oocytes.

With respect to the developmental stages (early, expanded, hatching and hatched) of blastocysts on Days 7, 8 and 9, only significant differences were found in the expanded blastocyst rate on Day 8 between the large oocytes and small oocytes from Treatments B (CM), C (GM/CM), F (mCM/CM) and G (GM/mGM). This result can be related to the process of IVM of each oocyte diameter in the patterns of gene expression involved in the expansion and hatching of embryos. In goat, Tan et al. (2009) demonstrated that the oocytes from larger follicles achieve cytoplasmic maturity earlier than those from small follicles to support embryo development. Thus, the ability of oocytes to develop after fertilization is related to the capacity to store the appropriate set of mRNA and proteins that will be required to support early embryo development and to initiate embryonic genome transcriptional activity. Finally, on Day 8, it was observed that expanded blastocysts from small oocytes matured with a high level of hormones and in the presence of ITS and ascorbic acid 12 h or 24 h (Treatments D and E) were not significantly different, when compared to the blastocysts from the large oocytes (Treatment A). This result could be related to the modest number of blastocysts obtained from the small oocytes.

We can conclude that the addition of ascorbic acid, insulin–transferrin–selenium and a low level of hormones during the first 12 h of IVM of pre-pubertal goat oocytes would be useful to comply with the requirements of the intrinsic quality of blastocysts produced *in vitro*, thus enhancing their cryosurvival during the process of cryopreservation. On the other

hand, the capacity of those oocytes to develop to the blastocyst stage *in vitro* seems to be more related to oocyte diameter than to the conditions of the IVM system.

Acknowledgements

This study was supported by the Spanish Ministry of Science and Innovation (Project Number: code: AGL2011-23784) and a predoctoral grant awarded to S.H. from the Spanish Ministry of Foreign Affairs and Cooperation. The authors would also like to thank the English assistance provided by Chuck Simmons, an English instructor of our University.

References

- Anguita B, Jimenez-Macedo AR, Izquierdo D, Mogas T, Paramio MT, 2007: Effect of oocyte diameter on meiotic competence, embryo development, p34 (cdc2) expression and MPF activity in prepubertal goat oocytes. *Theriogenology* **67**, 526–536.
- Armstrong DT, 2001: Effects of maternal age on oocyte developmental competence. *Theriogenology* **55**, 1303–1322.
- Catala MG, Izquierdo D, Romaguera R, Roura M, Paramio MT, 2010: Effect of a growth medium during IVM on embryo development of prepubertal ewe oocytes. *Reprod Fertil Dev* **22**, 323–324.
- Cordova B, Morato R, Izquierdo D, Paramio T, Mogas T, 2010: Effect of the addition of insulin-transferrin-selenium and/or L-ascorbic acid to the *in vitro* maturation of prepubertal bovine oocytes on cytoplasmic maturation and embryo development. *Theriogenology* **74**, 1341–1348.
- De la Fuente R, O'Brien MJ, Eppig JJ, 1999: Epidermal growth factor enhances preimplantation developmental competence of maturing mouse oocytes. *Hum Reprod* **14**, 3060–3068.
- Eppig JJ, Schroeder AC, 1989: Capacity of mouse oocytes from preantral follicles to undergo embryogenesis and development to live young after growth, maturation, and fertilization *in vitro*. *Biol Reprod* **41**, 268–276.
- Gomez E, Rodriguez A, Munoz M, Caamano JN, Hidalgo CO, Moran E, Facal N, Diez C, 2008: Serum free embryo culture medium improves *in vitro* survival of bovine blastocysts to vitrification. *Theriogenology* **69**, 1013–1021.
- Jeong YW, Hossein MS, Bhandari DP, Kim YW, Kim JH, Park SW, Lee E, Park SM, Jeong YI, Lee JY, Kim S, Hwang WS, 2008: Effects of insulin-transferrin-selenium in defined and porcine follicular fluid supplemented IVM media on porcine IVF and SCNT embryo production. *Anim Reprod Sci* **106**, 13–24.
- Jiang HS, Wang WL, Lu KH, Gordon I, Polge C, 1992: Examination of cell numbers of blastocysts derived from IVM IVF and IVC of bovine follicular oocytes. *Theriogenology* **37**, 229. Abstract.
- Jimenez-Macedo AR, Anguita B, Izquierdo D, Mogas T, Paramio MT, 2006: Embryo development of prepubertal goat oocytes fertilised by intracytoplasmic sperm injection (ICSI) according to oocyte diameter. *Theriogenology* **66**, 1065–1072.
- Kuwayama M, Vajta G, Kato O, Leibo SP, 2005: Highly efficient vitrification method for cryopreservation of human oocytes. *Reprod Biomed Online* **11**, 300–308.
- Lee MS, Kang SK, Lee BC, Hwang WS, 2005: The beneficial effects of insulin and metformin on *in vitro* developmental potential of porcine oocytes and embryos. *Biol Reprod* **73**, 1264–1268.
- Morato R, Izquierdo D, Paramio MT, Mogas T, 2010: Survival and apoptosis rates after vitrification in cryotop devices of *in vitro*-produced calf and cow blastocysts at different developmental stages. *Reprod Fertil Dev* **22**, 1141–1147.
- Murray AA, Swales AKE, Smith RE, Molinek MD, Hillier SG, Spears N, 2008: Follicular growth and oocyte competence in the *in vitro* cultured mouse follicle: effects of gonadotrophins and steroids. *Mol Hum Reprod* **14**, 75–83.
- Peluso JJ, 1988: Effect of the peripubertal pattern of LH and FSH secretion on *in vitro* oestradiol-17-Beta secretion and follicular growth within juvenile rat ovaries. *J Reprod Fertil* **84**, 239–245.
- Raghu HM, Nandi S, Reddy SM, 2002: Effect of insulin, transferrin and selenium and epidermal growth factor on development of buffalo oocytes to the blastocyst stage *in vitro* in serum-free, semidefined media. *Vet Rec* **151**, 260–265.
- Rizos D, Ward F, Boland MP, Lonergan P, 2001: Effect of culture system on the yield and quality of bovine blastocysts as assessed by survival after vitrification. *Theriogenology* **56**, 1–16.
- Romaguera R, Casanovas A, Morato R, Izquierdo D, Catala M, Jimenez-Macedo AR, Mogas T, Paramio MT, 2010: Effect of follicle diameter on oocyte apoptosis, embryo development and chromosomal ploidy in prepubertal goats. *Theriogenology* **74**, 364–373.
- Shamsuddin M, Larsson B, Rodriguezmartinez H, 1993: Culture of bovine IVM IVF embryos up to blastocyst stage in defined medium using Insulin, transferrin and selenium or growth factors. *Reprod Domest Anim* **28**, 209–210.
- Shirazi A, Soleimani M, Karimi M, Nazari H, Ahmadi E, Heidari B, 2010: Vitrification of *in vitro* produced ovine embryos at various developmental stages using two methods. *Cryobiology* **60**, 204–210.
- Takahashi Y, First NL, 1992: *In vitro* development of bovine one cell embryos influence of glucose, lactate, pyruvate, amino acids and vitamins. *Theriogenology* **37**, 963–978.
- Tan JH, Wu YG, Lan GC, Han ZB, Han D, Zhou P, Luo MJ, 2009: The rapidity of *in vitro* maturation of goat oocytes correlates with follicle sizes and affects blastocyst development of cloned embryos when used as recipient cytoplasts. *Biol Reprod* **81**, 275. Abstract.
- Tatemoto H, Ootaki K, Shigeta K, Muto N, 2001: Enhancement of developmental competence after *in vitro* fertilization of porcine oocytes by treatment with ascorbic acid 2-O-alpha-glucoside during *in vitro* maturation. *Biol Reprod* **65**, 1800–1806.
- Tatemoto H, Muto N, Sunagawa I, Shinjo A, Nakada T, 2004: Protection of porcine oocytes against cell damage caused by oxidative stress during *in vitro* maturation: role of superoxide dismutase activity in porcine follicular fluid. *Biol Reprod* **71**, 1150–1157.
- Wu D, Cheung QCK, Wen L, Li J, 2006: A growth-maturation system that enhances the meiotic and developmental competence of porcine oocytes isolated from small follicles. *Biol Reprod* **75**, 547–554.

Submitted: 30 Mar 2012; Accepted: 1 Jul 2012

Author's address (for correspondence): Dolores Izquierdo, Departament de Ciència Animal i dels Aliments, Facultat de Veterinària, UAB 08193 Bellaterra, Barcelona, Spain. E-mail: dolores.izquierdo@uab.cat

**In vitro developmental competence of prepubertal goat oocytes
cultured with recombinant activin-A**

Hammami S¹, Izquierdo D¹, Catalá MG¹, Paramio MT¹, Morató R².

¹Departament de Ciència Animal i dels Aliments, Universitat Autònoma de Barcelona,
Bellaterra, Spain

²Departament de Biologia i Biotecnologia de la Reproducció Animal i Humana
(TechnoSperm), Institut de Tecnologia Agroalimentaria, Universitat de Girona,
Girona, Spain

Animal 8 (2014) 94-101

In vitro developmental competence of prepubertal goat oocytes cultured with recombinant activin-A

S. Hammami¹, D. Izquierdo¹, M. G. Catalá¹, M. T. Paramio¹ and R. Morató^{2†}

¹Departament de Ciència Animal i dels Aliments, Facultat de Veterinària, Universitat Autònoma de Barcelona, 08193 Bellaterra, Barcelona, Spain;

²Department of Biology, Biotechnology of Animal and Human Reproduction (TechnoSperm), Institute of Food and Agricultural Technology, University of Girona, 17071 Girona, Spain

(Received 20 June 2013; Accepted 14 September 2013)

The present study was designed to evaluate the effect of activin-A during the *in vitro* oocyte maturation (IVM) and *in vitro* embryo culture (IVC) on nuclear maturation, blastocyst yield and blastocyst quality of prepubertal goat oocytes. In Experiment 1, three groups of oocytes were used during the IVM of prepubertal goat oocytes to determine the optimal concentration of recombinant human activin-A added to the maturation medium. Cumulus–oocyte complexes were matured in an IVM medium containing 0, 10 and 100 ng/ml (groups A0, A10 and A100), fertilized and *in vitro* cultured using standard procedures. In Experiment 2, the addition of 10 ng/ml activin-A at IVM (A10A0), IVC (A0A10) or IVM + IVC (A10A10) was studied and compared with the control group (A0A0). Results of the first experiment demonstrated that the addition of activin-A yielded similar percentages of maturation ($\leq 71.0\%$) and blastocyst formation rates ($\leq 24.9\%$) than the control group (A0). Experiment 2 showed that exposure of prepubertal goat oocytes to an IVC medium containing 10 ng/ml activin-A (A0A10) significantly increased the rates of development to the blastocyst stage, as compared with the control group (A0A0) ($19.5 \pm 2.21\%$ v. $13.1 \pm 2.37\%$, respectively; $P < 0.05$). With regard to the blastocyst quality, total number of cells, inner cell mass (ICM) and trophectoderm of prepubertal goat embryos produced in the presence of activin-A did not differ significantly among experimental groups. In summary, these results indicate that supplementation of the IVC medium with activin-A enhances embryo development of prepubertal goat oocytes.

Keywords: *in vitro* maturation, nuclear stage, *in vitro* culture, blastocysts, goat

Implications

In vitro embryo production in small ruminants provides an excellent source of low-cost embryos for basic research on developmental biology and for commercial application of the emerging biotechnologies such as nuclear transfer, transgenesis, embryo sexing and stem cells. Moreover, this technology would allow a fast multiplication of unique and expensive animals as endangered species. The use of oocytes from prepubertal female goats into breeding programs is advantageous because it can reduce the generation interval and increase the rate of genetic gain. Thus, a challenge is to establish a system that could potentially provide an unlimited source of competent oocytes for biomedical application.

Introduction

The use of prepubertal females as oocyte donors in *in vitro* embryo production allows for an acceleration in genetic

progress by shortening the generation interval. However, reduced developmental competence in oocytes of juvenile animals was addressed in numerous studies on ruminant species, including bovine (Khatir *et al.*, 1998), ovine (Ledda *et al.*, 1997) and caprine (Romaguera *et al.*, 2011). The low blastocyst formation rates obtained *in vitro* is mainly because of insufficient cytoplasmic maturation in the oocyte or inability to maintain embryo development beyond the transition of embryonic genome activation (Sirard *et al.*, 2006).

Oocyte growth and maturation are strictly dependent on establishing functional communications with the surrounding cumulus cells through gap junctions and reciprocal interactions mediated by paracrine and endocrine signals. These processes are accurately regulated by numerous growth factors and hormones. Moreover, during mammalian development, some cytokines, including activin-A, play a functional role in the process of cellular proliferation, differentiation and morphogenesis. In early pre-implantation embryos, cytokines produced by both the female genital tract and the embryo itself act on embryonic cells as paracrine/autocrine factors (reviewed by Heyner *et al.*, 1993).

[†] E-mail: roser.morato@udg.edu

Activin is an important member of the transforming growth factor β (TGF β) superfamily, and its roles in gonadal functions have been extensively studied in mammals (Mather *et al.*, 1997). Activins are homodimers or heterodimers of the β A or β B subunits of inhibin linked to one another by a single disulfide bond (Hata *et al.*, 1988). Dimerization of these β -subunits gives rise to three forms of activin referred to as activin-A (β A β A), activin-AB (β A β B) and activin-B (β B β B) (Ying, 1988). The expression of protein and mRNA for activin-A and activin receptors in the ovary has been localized in both oocyte and granulosa cells of follicles at various developmental stages (primates: Sidis *et al.*, 1998; swine: Van den Hurk and Van de Pavert, 2001; bovine: Hulshof *et al.*, 1997, Izadyar *et al.*, 1998; ovine: Thomas *et al.*, 2003; and caprine: Silva *et al.*, 2004). In addition to being a local regulator of folliculogenesis, activin-A is able to directly stimulate FSH synthesis and secretion, and to promote the release of the gonadotrophin-releasing hormone (GnRH) (Childs and Unabia, 1997). Activin-A can also stimulate the increase of FSH and LH receptors in granulosa cells, and it plays a role in progesterone production (Alak *et al.*, 1998; Tsuchiya *et al.*, 1999). Thus, the granulosa cells are likely to be the main source of paracrine factors and are crucial for oocyte maturation.

Activin-A and activin receptors have been detected in mouse (Lu *et al.*, 1993) and bovine (Yoshioka *et al.*, 1998b) embryos from zygote to the morula stage, suggesting that the protein might play a role in embryogenesis. As activin-A is produced by oviduct epithelial cells (Gandolfi *et al.*, 1995), the addition of activin-A to embryo culture *in vitro* may reproduce the environment of the cleavage-stage embryos in the oviduct *in vivo*. However, the effect of activin-A on *in vitro* embryo culture (IVC) is still controversial. Positive effects of activin-A on embryo development were shown when the protein was added at the earliest stages of bovine (Yoshioka and Kamomae, 1996; Yoshioka *et al.*, 1998a, Lee *et al.*, 2009) embryo culture, whereas no effects on blastocyst formation rates, depending on the timing of its addition to the culture medium, were reported in the literature either (Park *et al.*, 2010; Trigal *et al.*, 2011).

Summarizing, although prepubertal oocytes give rise to similar rates of fertilization and cleavage to those achieved using adult oocytes, the capacity of prepubertal oocytes to develop in the blastocyst stage is relatively lower and it would seem that embryos from prepubertal oocytes are less capable of establishing pregnancies (reviewed in Armstrong, 2001). Therefore, to improve the developmental competence of prepubertal oocytes after their recovery from the follicle, two approaches have been described. The first has been the addition of putative growth promoting substances to the culture medium (gonadotrophins, steroids, growth factors). The second approach has attempted to mimic the intrafollicular conditions of the oocytes. However, none of these approaches have managed to improve developmental competence from prepubertal goat oocytes. Keeping in mind that protein and mRNA for activin-A and activin receptors have been localized in both oocyte and granulosa cells of follicles

at various developmental stages in a variety of species including caprine (Silva *et al.*, 2004 and 2006), and the potential role of activin-A in the regulation of oocyte maturation has been demonstrated to promote *in vitro* oocyte maturation (IVM) in the rat (Itoh *et al.*, 1990), cow (Stock *et al.*, 1997), rhesus monkey (Alak *et al.*, 1996) and human (Alak *et al.*, 1998), we hypothesized that the addition of activin-A to the maturation and/or embryo culture medium could improve the *in vitro* potential of prepubertal goat oocytes for nuclear and cytoplasmic maturation and embryo development. Such a system could potentially provide an unlimited source of competent oocytes for biomedical application.

To the best of our knowledge, there is no study that reports the effects of activin-A on *in vitro* developmental potential of prepubertal goat oocytes. Therefore, the present study was conducted to evaluate the effect of adding activin-A at the IVM and IVC media on meiotic maturation, embryo development and embryo quality of *in vitro*-produced blastocysts.

Material and methods

Reagents

Unless specified, all chemicals, reagents and hormones were purchased from Sigma (St Louis, MO, USA). Activin-A was purchased from R&D Systems (Minneapolis, MN, USA), and it was reconstituted at 50 μ g/ml in sterile phosphate-buffered saline (PBS) containing at least 0.1% bovine serum albumin (BSA) and stored in aliquots at -20°C until use.

Oocyte collection and *in vitro* maturation (IVM)

Prepubertal goat (1 to 2 months old) ovaries were obtained from a slaughterhouse and transported to the laboratory in PBS at 38°C within 1 h of collection. Cumulus–oocyte complexes (COCs) were recovered by the slicing technique. Only COCs with a compact cumulus and homogeneous cytoplasm were selected for IVM. The COCs were washed three times in the IVM medium (TCM199 supplemented with 275 μ g/ml sodium pyruvate, 146 μ g/ml L-glutamine, 50 μ g/ml gentamycin, 10% (v/v) donor bovine serum (CanSera, Ontario, Canada), 10 μ g/ml FSH, 10 μ g/ml LH, 1 μ g/ml 17 β -estradiol and 100 μ M cysteamine. Groups of 25 to 30 COCs were transferred into drops of 100 μ l of the IVM medium, covered with mineral oil, and incubated for 24 h at 38.5°C and 5% CO_2 in humidified air.

In vitro fertilization (IVF)

Fresh semen was collected with an artificial vagina from four Murciano-Granadina bucks of proven fertility and transported to the laboratory at 35°C within 30 min. The motile sperm fraction was selected by the swim-up method (Parrish *et al.*, 1986). Briefly, 90 μ l of ejaculate mixture was deposited at the bottom of a conical tube containing 2 ml mDM (defined medium modified by Younis *et al.*, 1991). After 1 h of incubation at 38.5°C , the upper layer of the tubes was recovered in a tube of 15 ml and centrifuged at $160 \times g$

for 3 min. The sperm pellet was resuspended in a 1 : 1 proportion (v/v) with mDM containing heparin sodium salt (final concentration of 50 µg/ml), and it was incubated for 45 min at 38.5°C and 5% CO₂ in humidified air. After maturation, oocytes were washed three times in the fertilization medium (Tyrode's medium (TALP) supplemented with 1 µg/ml hypotaurine and 0.3 mg/ml glutathione) and groups of 25 to 30 oocytes were transferred to microdrops of 100 µl IVF medium, under mineral oil and co-cultured with sperm in a final concentration of 4 × 10⁶ motile sperm/ml at 38.5°C and 5% CO₂ in humidified air.

In vitro culture (IVC)

At 24-h post-insemination (hpi), presumptive zygotes were denuded of surrounding cumulus cells and attached spermatozoa, washed three times in synthetic oviductal fluid medium (SOF (Takahashi and First, 1992)), and groups of 20 presumptive embryos were placed into droplets of 20 µl of SOF medium supplemented with 10% fetal calf serum (FCS) under mineral oil and cultured in a humidified atmosphere with 5% CO₂, 5% O₂ and 90% N₂ for 8 days. The cleavage rate was evaluated at 48 hpi and blastocyst formation rates were recorded at Days 7, 8 and 9 post-insemination (pi).

Evaluation of nuclear maturation stage

After the *in vitro* maturation, oocytes were freed from their surrounding cumulus cells by vortexing and then washed three times in PBS. To evaluate meiotic progression, the oocytes were fixed in ethanol : acetic acid (3 : 1) overnight. The oocytes were mounted on microscope slides with vaseline, covered with a glass coverslip and stained with 1% Iacmoid in 45% acetic acid solution. The stage of nuclear maturation was assessed by phase-contrast microscopy (magnification, ×40). The criteria used to determine oocyte maturation stage have been described elsewhere (Rodríguez-Gonzalez *et al.*, 2002). Oocytes were scored as: germinal vesicle (GV) stage, when their chromatin, either filamentous or slightly condensed, was enclosed in a nuclear membrane; germinal vesicle breakdown (GVBD) stage, when there was no visible nuclear membrane and chromatin was condensed; metaphase I (MI) stage, when the chromatin was distributed at the equator of a large spindle; anaphase I stage, when homologous chromosomes could be seen to move away from each other and there was a visible spindle; telophase I stage, when homologous chromosomes were apart, and metaphase II (MII) stage, when the chromatin was located at the spindle and the first polar body extrusion was detected.

Evaluation of blastocyst quality

Hoechst staining. At the end of IVC, blastocysts were stained with 1 mg/ml Hoechst 33342 (H3570; Invitrogen, Life Technologies, Burlington, Ontario, Canada) for 5 min at 38°C. Cell count was carried out under a fluorescence microscope (Olympus BX53; Olympus, Germany).

Blastocyst differential staining. The cellular composition of Day 8 pi blastocysts was assessed by differential staining

of inner cell mass (ICM) and trophoctoderm (TE) cells, as described by Thouas *et al.* (2001), with some modifications. Briefly, embryos that reached the blastocyst stage were first incubated in PBS with 1% (v/v) Triton X-100 and 100 µg/ml propidium iodide for up to 15 s. Blastocysts were then immediately transferred into a fixative solution of 100% ethanol supplemented with 25 µg/ml Hoechst 33342 for 2 h. Fixed and stained whole blastocysts were mounted on a slide in a drop of 3 µl of glycerol and flattened with a coverslip. Cells were counted under UV illumination (Olympus BX53) and a 460 nm excitation filter. The images showed ICM and TE nuclei as blue and red/pink, respectively.

Experimental design

Two experiments were conducted in this study. Each experiment was replicated at least five times.

Experiment 1. *Effect of activin-A during IVM on the nuclear stage, embryo development and blastocyst quality of prepubertal goat oocytes.* The objective was to assess the effects of recombinant human activin-A at concentrations of 10 ng/ml and 100 ng/ml added at the IVM of prepubertal goat oocytes on nuclear maturation, cleavage and blastocyst formation rates, and embryo quality as compared with the control group, without activin-A. COCs were randomly assigned into one of the three IVM treatment groups: (1) A0: IVM medium without activin-A (control group); (2) A10: supplemented with 10 ng/ml activin-A and; (3) A100: supplemented with 100 ng/ml activin-A. After IVM, COCs were fertilized and the cleavage rate and blastocyst yield were assessed at Day 2 pi and Days 7, 8 and 9 pi, respectively. At the end of the culture period (Day 9 pi), and to determine the effects on embryo quality, *in vitro*-produced blastocysts were fixed and stained to determine the total cell number. In each repetition, a sample of COCs was obtained after IVM, denuded, washed in PBS, fixed overnight, Iacmoid-stained and observed under a phase-contrast microscope to determine their nuclear status.

Experiment 2. *In vitro embryo development in the presence of 10 ng/ml activin-A during IVM and IVC of prepubertal goat oocytes.* The aim was to evaluate the effect of recombinant human activin-A added to the IVM and IVC media on the cleavage rate, embryo development and blastocyst quality of oocytes from prepubertal goat. COCs were randomly allocated into four groups: (1) A0A0: COCs matured in the IVM medium and cultured in the SOF medium; (2) A0A10: COCs matured in the IVM medium and cultured in the SOF medium supplemented with 10 ng/ml activin-A; (3) A10A0: COCs matured in the IVM medium plus 10 ng/ml activin-A and cultured in the SOF medium; and (4) A10A10: COCs matured in the IVM medium plus 10 ng/ml activin-A and cultured in the SOF medium plus 10 ng/ml activin-A. The cleavage rate was recorded at 48 hpi and blastocysts yield at Days 7, 8 and 9 pi, respectively. To assess the quality of blastocysts obtained from the treatment groups, cell number and the cellular composition of Day 8 pi blastocysts were evaluated by differential staining of ICM and TE.

Statistical analysis

Data were analyzed using the Statistical Analysis Systems package (SAS, v9.1; Cary, NC, USA). At least five replicates were performed in all experimental groups. χ^2 analysis was used to determine which concentration groups differed in the nuclear stage after *in vitro* maturation. Cleavage rates, blastocyst yield and blastocyst cell number were analyzed using ANOVA. All data were checked for normality and homogeneity of variances using the Levene and Kolmogorov–Smirnov tests. The most suitable activin-A concentration for conducting the experiments depicted in Table 1 was determined after evaluating the effects of supplementing IVM with three different concentrations of activin-A (0, 10 and 100) on cleavage and blastocyst development rates by running a one-way ANOVA (factor = activin-A concentration in IVM) and a post-hoc Sidak test for pairwise comparisons). The effects of supplementing IVM and IVC media with 10 ng/ml activin-A on cleavage rate, number of cells per blastocyst, and accumulated number and percentages of blastocysts at Day 9 were evaluated through a two-way ANOVA where the factors were the composition of IVM (A0 v. A10) and IVC (A0 v. A10). On the other hand, the effects of 10 ng/ml activin-A

on the blastocyst development parameters, evaluated as numbers and percentages of blastocysts at Days 7, 8 and 9, were tested by a three-way ANOVA for repeated measures, where the two inter-subjects factors were IVM and IVC, and the intra-subject factor was the culturing time (Days 7, 8 and 9). A post-hoc Sidak’s test was also used for multiple comparisons. In all the cases, the interaction between factors within the model was also evaluated, and the significance level was set a $P < 0.05$.

Results

Experiment 1. Effect of activin-A during IVM on the nuclear stage, embryo development and blastocyst quality of prepubertal goat oocytes

At the end of the IVM period, nuclear maturation stages were established and results are shown in Figure 1. Oocytes cultured in the presence of activin-A showed similar rates of oocytes at GV, MI, ATI and MII stage to those of the control group (A0). Indeed, the percentages of MII oocytes recorded after maturation for the 10 ng/ml and 100 ng/ml concentrations were not significantly different from those observed for the control oocytes ($P > 0.05$). No differences according to activin-A concentration were observed ($P > 0.05$).

Table 1 shows the embryo development rates of oocytes cultured in the maturation medium alone or in the medium supplemented with activin-A (10 and 100 ng/ml). No differences in cleavage rates were detected among the different experimental groups ($P > 0.05$). Similarly, no significant differences ($P > 0.05$) were observed in the percentage of embryos developing to the blastocyst stage on Days 7, 8 and 9 pi. No interaction ($P > 0.05$) between activin-A concentration and culture period were either observed.

An amount of 54 blastocysts were stained to assess the embryo quality. The total cell number of Day 8 blastocysts obtained from oocytes cultured in the maturation medium alone or in the medium supplemented with activin-A is




Figure 1 Meiotic status of prepubertal goat oocytes matured in TCM 199 supplemented with various concentrations of activin-A during IVM. Treatments A0, A10 and A100: oocytes matured in TCM199 supplemented with 0, 10 and 100 ng/ml of activin-A for 24 h. GV=germinal vesicle; MI=metaphase I; A–T=anaphase–telophase; MII=metaphase II; DEG=degenerated oocytes.

Table 1 Effects of supplementing IVM medium with three different concentrations of activin-A (0, 10 and 100 ng/ml) on cleavage and blastocyst development rates from prepubertal goat oocytes

Items	Activin-A treatments ^a			RMSE	Contrasts (P-values)		
	A0	A10	A100		A0 v. A10	A0 v. A100	A10 v. A100
No. of oocytes	398	406	406				
% of cleaved oocytes	41.67	46.47	45.90	7.88	0.50	0.58	0.99
% Blastocysts (Day 7)*	11.70	16.95	12.50	4.50	0.10	0.94	0.18
% Blastocysts (Day 8)*	16.69	21.97	18.52	5.65	0.21	0.82	0.50
% Blastocysts (Day 9)*	21.74	24.93	21.89	5.65	0.55	0.99	0.58
% Blastocysts (Day 9)**	9.03	11.43	9.85	2.62	0.23	0.83	0.51

RMSE = root mean squared error.

Data are given as means. Treatments A0, A10 and A100 were: oocytes matured in TCM199 supplemented with 0, 10 and 100 ng/ml of activin-A, respectively.

^aOverall treatment effect not significant for any of the measured parameters ($P > 0.05$). Data analyzed by ANOVA with activin-A concentration in *in vitro* maturation medium as a treatment factor.

*Rate of blastocyst is calculated from cleaved embryos.

**Rate of blastocyst is calculated from *in vitro* matured oocytes.

Table 2 Effect of supplementing IVM medium with three different concentrations of activin-A (0, 10 and 100 ng/ml) on mean cell number from prepubertal goat blastocysts

Items	Activin-A treatments ^a			RMSE	Contrasts (P-values)		
	A0	A10	A100		A0 v. A10	A0 v. A100	A10 v. A100
No. of blastocysts	18	18	18				
Blastocyst mean cell number	176.72	192.83	187.94	75.77	0.80	0.90	0.98

RMSE = root mean squared error.

Data are given as means. Treatments A0, A10 and A100 were: oocytes matured in TCM199 supplemented with 0, 10 and 100 ng/ml of activin-A, respectively.

^aOverall treatment effect not significant for any of the measured parameters ($P > 0.05$). Data analyzed by ANOVA with activin-A concentration in *in vitro* maturation medium as a treatment factor.

Table 3 Effects of supplementing IVM and IVC media with 10 ng/ml activin-A on cleavage and blastocyst development rates from prepubertal goat oocytes

Items	Activin-A supplementation				RMSE	P-value
	A0A0	A10A0	A0A10	A10A10 ^c		
No. of oocytes	296	267	283	287		
% of cleaved oocytes	50.87	50.79	54.96	51.68	10.76	0.89
% Blastocysts (Day 7)*	7.32	8.14	11.27	8.91	3.39	0.24
% Blastocysts (Day 8)*	10.51	13.54	14.64	11.25	3.95	0.26
% Blastocysts (Day 9)*	13.13 ^a	15.77 ^{ab}	19.51 ^b	13.69 ^{ab}	4.81	0.12
% Blastocysts (Day 9)**	6.23 ^a	7.74 ^a	10.52 ^b	6.91 ^a	2.09	0.01

IVM = *in vitro* maturation; IVC = *in vitro* embryo culture; RMSE = root mean squared error.

Data are given as means. Treatment A0A0 and A0A10 were: oocytes matured in TCM199 and cultured in SOF medium supplemented with 0 and 10 ng/ml of activin-A, respectively. Treatments A10A0 and A10A10 were: oocytes matured in TCM199 supplemented with 10 ng/ml and cultured in SOF medium with 0 and 10 ng/ml of activin-A, respectively.

^{a,b}Values in the same row with different superscript letters differ significantly ($P < 0.05$).

^cInteraction effects were observed only between addition of activin-A at IVM medium and IVC medium ($P = 0.001$). Data analyzed by ANOVA where the factors were the composition of IVM (A0 v. A10) and IVC (A0 v. A10).

*Rate of blastocyst is calculated from cleaved embryos.

**Rate of blastocyst is calculated from *in vitro* matured oocytes.

summarized in Table 2. There were no differences among groups, irrespective of the maturation medium ($P > 0.05$). All blastocysts showed a similar number of cells.

Experiment 2. *In vitro* embryo development in the presence of 10 ng/ml activin-A during IVM and IVC of prepubertal goat oocytes

There were no significant differences in the cleavage rate among treatments. No significant differences were either observed when activin-A was added at the *in vitro* maturation medium (A10A0; A10A10) ($P > 0.05$). However, embryo development, in terms of total number of blastocysts obtained at Day 9, improved significantly ($P < 0.05$) with the addition of activin-A at the *in vitro* culture medium (A0A10). The highest blastocyst formation rate (19.5%) was obtained when activin-A was added at the *in vitro* culture medium (A0A10) but not at the *in vitro* maturation medium. Moreover, interaction effects between addition of activin-A at IVM medium and IVC medium were observed ($P = 0.001$) (Table 3).

Blastocyst quality was evaluated in terms of total number of cells per blastocyst and the proportion of ICM cells and TE cells assessed by the differential staining (Table 4).

No differences in any of these variables were detected among groups ($P > 0.05$).

Discussion

The present study was designed to evaluate the effect of activin-A during the IVM and IVC on maturation rates, embryo development and embryo quality of prepubertal goat oocytes. In general, prepubertal oocytes show reduced developmental competence, when compared with results obtained with adult oocytes (Armstrong, 2001). Indeed, the defective capacity of these oocytes to develop up to blastocyst at a rate equivalent to that of adult oocytes has been reported by several other studies in cattle (Presicce *et al.*, 1997), in sheep (Ledda *et al.*, 1997), in pig (Marchal *et al.*, 2001) and more recently in goats stimulated hormonally (Leoni *et al.*, 2009).

In the present work, the presence of activin-A during IVM of cumulus-enclosed prepubertal goat oocytes did not affect nuclear maturation rates. Thus, the results obtained in Experiment 1 indicate that activin-A added to maturation media at concentrations of 10 and 100 ng/ml resulted in

Table 4 Number of TE and ICM cells in blastocysts developed from goat oocytes after IVM and IVC with or without 10 ng/ml activin-A

Items	Activin-A supplementation ^a				RMSE	P-value
	A0A0	A10A0	A0A10	A10A10		
Total blastocysts	9	8	14	10		
Total number of cells	72.44	76.75	79.00	73.80	14.12	0.69
TE cells	50.78	54.62	56.50	52.00	10.30	0.56
ICM cells	21.67	22.12	22.50	21.80	4.72	0.97
% ICM : TE	0.43	0.41	0.40	0.42	0.06	0.70

TE = trophectoderm; ICM = inner cell mass; IVM = *in vitro* maturation; IVC = *in vitro* embryo culture; RMSE = root mean squared error.

Data are given as means. Treatment A0A0 and A0A10 were: oocytes matured in TCM199 and cultured in presence of 0 and 10 ng/ml of activin-A, respectively. Treatments A10A0 and A10A10 were: oocytes matured in TCM199 supplemented with 10 ng/ml and cultured SOF medium supplemented with 0 and 10 ng/ml of activin-A, respectively.

^aOverall treatment effect not significant for any of the measured parameters ($P > 0.05$). Data analyzed by ANOVA where the factors were the composition of IVM (A0 v. A10) and IVC (A0 v. A10).

similar rates of oocytes in the MII stage, as compared with those oocytes that matured without activin-A. Similar results have already been reported in bovine (Vantol *et al.*, 1994; Izadyar *et al.*, 1996) and porcine (Coskun and Lin, 1994) oocytes, suggesting no effect of activin-A on oocyte maturation. Considering that activin-A is able to directly stimulate FSH synthesis and secretion, promote the release of the GnRH, and also stimulate the increase of FSH and LH receptors in the granulosa cells (Alak *et al.*, 1998; Norwitz *et al.*, 2002), the ineffectiveness of exogenous activin-A may be either because of activin-A production by the COCs and/or the presence of FSH and LH in our culture medium. However, this does not exclude the possibility that activin-A, secreted by the granulosa cells and present in the follicular fluid, has an effect on the cytoplasmic maturation of goat oocytes, which would affect subsequent pre-implantation embryo development. Nevertheless, our results demonstrate that activin-A added to IVM has no effect on the percentage of blastocysts obtained after IVF. This observation is consistent with other reports in bovine (Vantol *et al.*, 1994; Izadyar *et al.*, 1996) and pig (Coskun and Lin, 1994), which showed that adding activin-A during IVM did not result in an improvement of blastocyst development.

Regarding the effect of activin-A during the whole embryo culture period, our data clearly show that activin-A promotes embryonic development, showing an improving effect on the blastocyst yield of prepubertal goat oocytes. Given that activin-A is produced by the oviduct epithelial cells (Gandolfi *et al.*, 1995), addition of activin-A to embryo culture *in vitro* could help reproduce the environment of the cleavage-stage embryos in the oviduct *in vivo*. Controversial results have been reported about the effects of activin-A on embryo development depending on the timing of its addition to the culture medium. Thus, Yoshioka *et al.* (1998a) observed that activin-A had an effect on blastocyst yield when it was added to IVC medium of bovine embryos before the 9- to 16-cell stage. However, a recent study on bovine (Trigal *et al.*, 2011) has demonstrated a better effect of activin-A when it is added later in culture media. The development-enhancing effect of activin-A may be associated with the timing of the

activation of appropriate genes within the embryonic genome. Studies in mouse (Lu *et al.*, 1993) and bovine (Yoshioka *et al.*, 1998b) showed that activin-A and activin receptors are detected in embryos from the zygote to the morula stage, suggesting that the protein might play a role in embryogenesis during the early stage of embryo development. Our data show an evident improving effect on Day 9 blastocyst yield, suggesting an effect during the late stages of embryo development in prepubertal goats. Otherwise, we should also consider the possibility of an accumulative effect of activin-A added during *in vitro* maturation and culture medium. Consequently, greater effects are observed as the *in vitro* culture is longer. Thus, further studies will be required to determine the functions of activin-A during these stages of development.

Finally, the quality of blastocysts, as determined by the evaluation of their cell number, seemed to be similar among all of the groups. The number of cells of the embryo may be informative of embryonic viability (VanSoom *et al.*, 1997). The lower quality of embryos produced *in vitro* could be attributed to a lower number of cells. Cell counts were unaffected by the presence of activin-A, similarly to other authors showing no differences in total cell counts between bovine blastocysts produced in the presence or absence of activin-A (Yoshioka *et al.*, 1998a, Trigal *et al.*, 2010). In addition, in our study, blastocyst cell counts were similar to those reported in adult and prepubertal goats (Koeman *et al.*, 2003; Romaguera *et al.*, 2010).

Several studies have demonstrated that the expression of protein and mRNA for activin-A and activin receptors in the ovary is present in both oocytes and granulosa cells of follicles at various stages of development (pigs: (Van den Hurk and Van de Pavert, 2001); and ruminants: (Hulshof *et al.*, 1997; Izadyar *et al.*, 1998)), suggesting a physiological role of activin in the growth and development of follicles and/or oocytes. Although there is convincing evidence that activin signaling is important for ovarian function, information on its localization and function is mainly obtained from rodents and cows. Few studies on goats (Silva *et al.*, 2004 and 2006) have demonstrated that activin-A, its binding protein

follicle-stimulating hormone and its receptors are formed in all types of goat follicles, in all compartments of antral follicles and in ovarian surface epithelium. The widespread distribution pattern of the follistatin–activin–activin receptor system in goat ovaries points to a crucial role of this system in various reproductive processes, including follicle growth and differentiation, and luteal activity. However, there are limited data on goats and on prepubertal animals (Brawtal, 1994; Zhao *et al.*, 2001; Patel *et al.*, 2007) about the expression of protein and mRNA for activin-A and activin receptors. Therefore, it is necessary to determine whether activin-A and its receptors are present in the embryos from the zygote to the morula stage to improve the results on *in vitro* embryo development of prepubertal goat oocytes.

In conclusion, the addition of 10 ng/ml of activin-A throughout the whole embryo culture improves blastocyst yield. This is the first study that provides data of the effects of activin-A on *in vitro* developmental potential of prepubertal goat oocytes. Nevertheless, to the best of our knowledge, there is no study reporting the gene expression and protein localization for activin-A and activin receptors in prepubertal goats. A detailed profiling of the activin-A effects on embryonic development of prepubertal goat oocytes and determination of gene expression and protein synthesis for activin-A require further experiments.

Acknowledgments

This study was financially supported by the Spanish Ministry of Science and Innovation (Project Number: code: AGL2011-23784) and a predoctoral grant awarded to S.H. from the Spanish Ministry of Foreign Affairs and Cooperation. The authors are grateful to Dr Chuck Simmons, an English instructor of our University, for his English assistance.

References

Alak BM, Smith GD, Woodruff TK, Stouffer RL and Wolf DP 1996. Enhancement of primate oocyte maturation and fertilization *in vitro* by inhibin A and activin A. *Fertility and Sterility* 55, 646–653.

Alak BM, Coskun S, Friedman CI, Kennard EA, Kim HM and Seifer DB 1998. Activin A stimulates meiotic maturation of human oocytes and modulates granulosa cell steroidogenesis *in vitro*. *Fertility and Sterility* 70, 1126–1130.

Armstrong DT 2001. Effects of maternal age on oocyte developmental competence. *Theriogenology* 55, 1303–1322.

Brawtal R 1994. Expression of messenger-RNA for follistatin and inhibin/activin subunits during follicular-growth and atresia. *Journal of Molecular Endocrinology* 13, 253–264.

Childs G and Unabia G 1997. Cytochemical studies of the effects of activin on gonadotropin-releasing hormone (GnRH) binding by pituitary gonadotropes and growth hormone cells. *Journal of Histochemistry and Cytochemistry* 45, 1603–1610.

Coskun S and Lin Y 1994. Effects of transforming growth-factors and activin A on *in-vitro* porcine oocyte maturation. *Molecular Reproduction and Development* 38, 153–159.

Gandolfi F, Modina S, Brevini T, Passoni L, Artini P, Petraglia F and Lauria A 1995. Activin beta(A) subunit is expressed in bovine oviduct. *Molecular Reproduction and Development* 40, 286–291.

Hata K, Kilkuchi K, Tada K and Tsuiki S 1988. Analysis of glucose-6-phosphate translocase and hexose-6-phosphatase phosphohydrolase, the 2 obligatory components of microsomal glucose-6-phosphatase system, in rat-liver. *Tohoku Journal of Experimental Medicine* 155, 173–181.

Heyner S, Shan N, Smith R, Watson A and Schultz G 1993. The role of growth-factors in embryo production. *Theriogenology* 39, 151–161.

Hulshof S, Figueiredo J, Beckers J, Bevers M, Vanderstichele H and Van den Hurk R 1997. Bovine preantral follicles and activin: Immunohistochemistry for activin and activin receptor and the effect of bovine activin A *in vitro*. *Theriogenology* 48, 133–142.

Itoh M, Igarashi M, Yamada K, Hasegawa Y, Seki M, Eto Y and Shibai H 1990. Activin A stimulates meiotic maturation of the rat oocyte *in vitro*. *Biochemical and Biophysical Research Communications* 166, 1479–1484.

Izadyar F, Zeinstra E, Colenbrander B, Vanderstichele H and Bevers M 1996. *In vitro* maturation of bovine oocytes in the presence of bovine activin A does not affect the number of embryos. *Animal Reproduction Science* 45, 37–45.

Izadyar F, Dijkstra G, Van Tol H, van de Eljden-Van Raaij A, Van den Hurk R, Colenbrander B and Bevers M 1998. Immunohistochemical localization and mRNA expression of activin, inhibin, follistatin, and activin receptor in bovine cumulus–oocyte complexes during *in vitro* maturation. *Molecular Reproduction and Development* 49, 186–195.

Khatir H, Lonergan P and Mermillod P 1998. Kinetics of nuclear maturation and protein profiles of oocytes from prepubertal and adult cattle during *in vitro* maturation. *Theriogenology* 50, 917–929.

Koeman J, Keefer C, Baldassarre H and Downey B 2003. Developmental competence of prepubertal and adult goat oocytes cultured in semi-defined media following laparoscopic recovery. *Theriogenology* 60, 879–889.

Ledda S, Bogliolo L, Calvia P, Leoni G and Naitana S 1997. Meiotic progression and developmental competence of oocytes collected from juvenile and adult ewes. *Journal of Reproduction and Fertility* 109, 73–78.

Lee K, Bettogowda A, Wee G, Ireland JJ and Smith GW 2009. Molecular determinants of oocyte competence: potential functional role for maternal (oocyte-derived) follistatin in promoting bovine early embryogenesis. *Endocrinology* 150, 2463–2471.

Leoni GG, Succu S, Satta V, Paolo M, Bogliolo L, Bebbere D, Spezzigu A, Madeddu M, Berlinguer F, Ledda S and Naitana S 2009. *In vitro* production and cryotolerance of prepubertal and adult goat blastocysts obtained from oocytes collected by laparoscopic oocyte-pick-up (LOPU) after FSH treatment. *Reproduction Fertility and Development* 21, 901–908.

Lu R, Matsuyama S, Nishihara M and Takahashi M 1993. Developmental expression of activin/inhibin beta A, beta B, and alpha subunits, and activin receptor-IB genes in preimplantation mouse embryos. *Biology of Reproduction* 49, 1163–1169.

Marchal R, Feugang J, Perreau C, Venturi E, Terqui M and Mermillod P 2001. Meiotic and developmental competence of prepubertal and adult swine oocytes. *Theriogenology* 56, 17–29.

Mather JP, Moore A and Li RH 1997. Activins, inhibins, and follistatins: further thoughts on a growing family of regulators. *Proceedings of the Society for Experimental Biology and Medicine* 215, 209–222.

Norwitz E, Xu S, Jeong K, Bedecarrats G, Winebrenner L, Chin W and Kaiser U 2002. Activin A augments GnRH-mediated transcriptional activation of the mouse GnRH receptor gene. *Endocrinology* 143, 985–997.

Park JE, Oh HJ, Hong SG, Jang G, Kim MK and Lee BC 2010. Effects of activin A on the *in vitro* development and mRNA expression of bovine embryos cultured in chemically-defined two-step culture medium. *Reproduction in Domestic Animals* 45, 585–593.

Parrish J, Suskoparrish J, Leibfriedrutledge M, Cristser E, Eyestone W and First N 1986. Bovine *in vitro* fertilization with frozen-thawed semen. *Theriogenology* 25, 591–600.

Patel O, Bettogowda A, Ireland J, Coussens P, Lonergan P and Smith G 2007. Functional genomics studies of oocyte competence: evidence that reduced transcript abundance for follistatin is associated with poor developmental competence of bovine oocytes. *Reproduction* 133, 95–106.

Presicce G, Jiang S, Simkin M, Zhang L, Looney C, Godke R and Yang X 1997. Age and hormonal dependence of acquisition of oocyte competence for embryogenesis in prepubertal calves. *Biology of Reproduction* 56, 386–392.

Rodriguez-Gonzalez E, Lopez-Bejar M, Velilla E and Paramio M 2002. Selection of prepubertal goat oocytes using the brilliant cresyl blue test. *Theriogenology* 57, 1397–1409.

Romaguera R, Casanovas A, Morato R, Izquierdo D, Catala M, Jimenez-Macedo AR, Mogas T and Paramio MT 2010. Effect of follicle diameter on oocyte apoptosis, embryo development and chromosomal ploidy in prepubertal goats. *Theriogenology* 74, 364–373.

- Romaguera R, Moll X, Morato R, Roura M, Palomo MJ, Catala MG, Jimenez-Macedo AR, Hammami S, Izquierdo D, Mogas T and Paramio MT 2011. Prepubertal goat oocytes from large follicles result in similar blastocyst production and embryo ploidy than those from adult goats. *Theriogenology* 76, 1–11.
- Sidis Y, Fujiwara T, Leykin L, Isaacson K, Toth T and Schneyer A 1998. Characterization of inhibin/activin subunit, activin receptor, and follistatin messenger ribonucleic acid in human and mouse oocytes: evidence for activin's paracrine signaling from granulosa cells to oocytes. *Biology of Reproduction* 59, 807–812.
- Silva J, van den Hurk R, van Tol H, Roelen B and Figueiredo J 2004. Gene expression and protein localisation for activin-A, follistatin and activin receptors in goat ovaries. *Journal of Endocrinology* 183, 405–415.
- Silva J, Tharasanit T, Taverne M, Van der Weijden G, Santos R, Figueiredo J and Van den Hurk R 2006. The activin-follistatin system and in vitro early follicle development in goats. *Journal of Endocrinology* 189, 113–125.
- Sirard MA, Richard F, Blondin P and Robert C 2006. Contribution of the oocyte to embryo quality. *Theriogenology* 65, 126–136.
- Stock A, Woodruff T and Smith L 1997. Effects of inhibin A and activin A during in vitro maturation of bovine oocytes in hormone- and serum-free medium. *Biology of Reproduction* 56, 1559–1564.
- Takahashi Y and First NL 1992. In vitro development of bovine one cell embryos influence of glucose, lactate, pyruvate, amino acids and vitamins. *Theriogenology* 37, 963–978.
- Thomas FH, Armstrong DG and Telfer EE 2003. Activin promotes oocyte development in ovine preantral follicles in vitro. *Reproductive Biology and Endocrinology* 1, 76.
- Thouas GA, Korfiatis NA, French AJ, Jones GM and Trounson AO 2001. Simplified technique for differential staining of inner cell mass and trophoctoderm cells of mouse and bovine blastocysts. *Reproductive BioMedicine Online* 3, 25–29.
- Trigal B, Gomez E, Diez C, Caamano JN, Martin D, Carrocera S and Munoz M 2011. In vitro development of bovine embryos cultured with activin A. *Theriogenology* 75, 584–588.
- Trigal B, Gomez E, Diez C, Caamano JN, Molina I, Carrocera S, Martin D and Munoz M 2010. In vitro development of bovine morulae produced and/or cultured with activin. *Reproduction Fertility and Development* 22, 236.
- Tsuchiya M, Minegishi T, Kishi H, Tano M, Kameda T, Hirakawa T, Ibuki Y, Mizutani T and Miyamoto K 1999. Control of the expression of luteinizing hormone receptor by local factors in rat granulosa cells. *Archives of Biochemistry and Biophysics* 367, 185–192.
- Van den Hurk R and Van de Pavert S 2001. Localization of an activin/activin receptor system in the porcine ovary. *Molecular Reproduction and Development* 60, 463–471.
- VanSoom A, Ysebaert M and DeKruif A 1997. Relationship between timing of development, morula morphology, and cell allocation to inner cell mass and trophoctoderm in in vitro-produced bovine embryos. *Molecular Reproduction and Development* 47, 47–56.
- Vantol H, Deloos F, Vanderstichele H and Bevers M 1994. Bovine activin A does not affect the in vitro maturation of bovine oocytes. *Theriogenology* 41, 673–679.
- Ying S 1988. Inhibins, activins, and follistatins – gonadal proteins modulating the secretion of follicle-stimulating-hormone. *Endocrine Reviews* 9, 267–293.
- Yoshioka K and Kamomae H 1996. Recombinant human activin a stimulates development of bovine one-cell embryos matured and fertilized in vitro. *Molecular Reproduction and Development* 45, 151–156.
- Yoshioka K, Suzuki C and Iwamura S 1998a. Activin A and follistatin regulate developmental competence of in vitro-produced bovine embryos. *Biology of Reproduction* 59, 1017–1022.
- Yoshioka K, Takata M, Taniguchi T, Yamanaka H and Sekikawa K 1998b. Differential expression of activin subunits, activin receptors and follistatin genes in bovine oocytes and preimplantation embryos. *Reproduction Fertility and Development* 10, 293–298.
- Younis A, Zuelke K, Happer K, Oliveira M and Brackett B 1991. In vitro fertilization of goat oocytes. *Biology of Reproduction* 44, 1177–1182.
- Zhao J, Taverne M, van der Weijden G, Bevers M and Van den Hurk R 2001. Effect of activin A on in vitro development of rat preantral follicles and localization of activin A and activin receptor II. *Biology of Reproduction* 65, 967–977.

G_{ENERAL} **D**_{ISCUSSION}

General Discussion

This thesis is focused on improvement of the *in vitro* embryo production from oocytes of prepubertal goats. For several years, *in vitro* embryo production from prepubertal goats has been considered a promising option for providing an excellent source of embryos for basic research on development biology and for commercial applications of transgenic and cloning technologies. However, despite successful results and reports of numerous attempts, the effectiveness of *in vitro* embryo production protocols for prepubertal goat oocytes has never reached a satisfactory level. Oocyte quality is the main factor for embryos reaching blastocyst stage from IVM/IVF/IVC oocytes. Until now, attention has been focused on selection of criteria to assess the most competent oocytes to develop up to blastocyst stage. In our laboratory, several studies have been reported describing the best criteria to select the larger and more competent oocytes for *in vitro* blastocyst production: Brilliant Cresyl Blue (BCB) stain (Rodriguez-Gonzalez *et al.*, 2002; Catalá *et al.*, 2011), oocyte (Jiménez-Macedo *et al.*, 2006; Anguita *et al.*, 2007) and follicle (Romaguera *et al.*, 2010a; 2011) diameters.

People working in this field has to combine the most adequate oocyte selection criteria and the *in vitro* culture protocol to became high rates of blastocyst development. For this reason, the two studies included in the present thesis work were designed to optimize *in vitro* culture system of prepubertal goat oocytes collected from ovaries of slaughtered goat females, using supplements added during IVM and/or IVC, in order to increase both rates of development and quality of *in vitro*-produced blastocysts at a rate equivalent to that of adult oocytes.

The first study was conducted to examine the effect of supplementing *in vitro* maturation medium with low level of hormones and/or insulin-transferrin-selenium plus ascorbic acid in the developmental competence and embryo quality of small oocytes from prepubertal goats.

Previous studies in our laboratory using prepubertal goat oocytes showed that not only the age of the oocyte donor is crucial for embryo production, but also the size of oocyte plays an important role in embryo production. Firstly, we hypothesized that supplementing the IVM medium of small oocytes with low hormone levels, ITS and L-

ascorbic acid called growth medium, could significantly improve maturation along with developmental competency compared to the conventional goat oocyte maturation system. However, our results have showed that the culture of small prepubertal goat oocytes in the growth medium during IVM did not improve the blastocyst percentage when compared to those matured in conventional IVM medium (Hammami *et al.*, 2013). Similarly, in ovine, Catalá *et al.*, (2013) also observed that growth medium failed to increase embryo production of lamb oocytes. These discrepancies with the findings of Wu *et al.*, (2006) on gilt oocytes and Córdova *et al.*, (2010) on calf oocytes may be related to variability among biological requirements of immature oocytes depending on species used.

In these initial studies it was possible to duplicate the performance of blastocyst production with the use of oocyte diameter as selection criteria. The rate of blastocysts production from prepubertal goat oocytes $\geq 125 \mu\text{m}$ was 20.65% compared to 10.40 % from oocytes $< 125 \mu\text{m}$ diameter. Thus, the capacity of those oocytes, once fertilized, to develop to the blastocyst stage *in vitro* seems to be more related to oocyte diameter than to the conditions of *in vitro* maturation system. Several studies concluded that oocyte diameter is correlated to follicle diameter and as both increase the developmental capability of the oocytes improves (revised by Gandolfi *et al.*, 2005). It is noteworthy that COCs coming from a wide range of follicle diameters and selected based on their morphology alone produced higher cleavage and blastocyst rates than oocytes from small follicles. One possibility could be, as Romaguera *et al.*, (2010a) discussed, that oocytes from follicles of $\geq 3 \text{ mm}$ could have enough time to be submitted to prematuration processes and to acquire a complete developmental competence to blastocyst stage. Anguita *et al.*, (2007) confirmed this hypothesis previously in our laboratory showing that prepubertal goat oocyte diameter and, therefore, oocyte developmental competence seems to be related to the amount of cyclin B, cdc2 kinase protein and MPF activity. Maturation promoting factor (MPF) is a heterodimer composed of a p34cdc2 catalytic subunit, with serine-threonine kinase activity, and a cyclin B1 regulatory subunit. MPF activity appears just before germinal vesicle break down (GVBD) increasing until metaphase I; its activity is decreased in anaphase-telophase while its maximum level is reached at metaphase II (MII). In prepubertal goats, Anguita *et al.*, (2007) showed a higher MPF activity in oocytes with a diameter larger than $135 \mu\text{m}$ and, after IVM, p34cdc2 protein expression increased with the

oocyte diameter. Possibly, the p34cdc2 RNA accumulation in the oocyte takes place during the last growth stage and the translation begins when the oocyte reaches its maximum size. MPF activity seems to confer higher developmental competence to oocytes through promotion of cytoplasmic maturation. Hence, enough time to be submitted to prematuration processes and to acquire a complete oocyte competence is important for successful embryo development.

When analysing the effects of different growth medium components combinations (low level of hormones, ITS and ascorbic acid with the conventional medium during IVM), on the cleavage and developmental capacity of prepubertal goat oocytes, our findings indicated that the supplementation of conventional medium with ascorbic acid and ITS during 12h and 24h in presence of normal level of hormones provided a significant decrease in the percentage of blastocysts obtained from <125 μm oocytes diameter, compared to the conventional IVM medium (7.65% and 7.05% vs 13.12%, respectively). According to our data, in calf oocytes, Córdova *et al.*, (2010) observed that the presence of ascorbic acid significantly reduced the proportion of embryos developing to the blastocyst stage. These data are in disagreement with observations in pig and buffalo oocytes in that the presence of ITS during *in vitro* maturation had a beneficial effect on the nuclear and cytoplasmic maturation (Jeong *et al.*, 2008) and on the yield of blastocysts (Raghu *et al.*, 2002a). On the other hand, in absence of supplementation with ITS and ascorbic acid, no differences were observed on cleavage and blastocyst rates using low or normal level of hormones during the first 12h of IVM. The growth of mouse oocytes in a medium without FSH has been demonstrated, and these oocytes acquire competence to undergo maturation, fertilization and embryonic development (Peluso, 1988; Eppig and Schroeder, 1989; Wu *et al.*, 2000; Murray *et al.*, 2008). Also, when mouse follicles have been cultured in presence of LH, their growth was not affected (Murray *et al.*, 2008). These results suggest that gonadotrophins may not be required for the acquisition of full development competence by growing oocytes.

Aside from adding antioxidants, vitamins and/or different hormone levels during IVM, in the second study we investigated whether activin-A supplementation to the IVM medium of prepubertal goat oocytes, could affect their nuclear maturation and competence to develop up to blastocyst stage. Results of this study (Hammami *et al.*, 2014) showed no effect of activin-A on oocyte maturation and subsequent embryo

development, similarly to results reported in bovine (Van Tol *et al.*, 1994; Izadyar *et al.*, 1996) and porcine (Coskun and Lin, 1994) oocytes. Considering that activin-A is able to directly stimulate FSH synthesis and secretion, promote the release of the gonadotrophin-releasing hormone (GnRH), and also stimulate the increase of FSH and LH receptors in granulosa cells (Alak *et al.*, 1998; Norwitz *et al.*, 2002), the ineffectiveness of exogenous activin-A may be either due to activin-A production by the COCs and/or the presence of FSH and LH in our culture medium.

The two studies suggest that in our conditions no effectiveness on the subsequent development to the blastocyst stage of prepubertal goat oocytes is obtained, neither when different maturation media are combined (ascorbic acid, insulin–transferrin–selenium, and a low level of hormones) with the conventional IVM medium using small oocytes diameter, nor when activin-A is added to the conventional IVM medium.

Given that activin-A is produced by oviduct epithelial cells (Gandolfi *et al.*, 1995), addition of activin-A to *in vitro* embryo culture could help to reproduce the environment of the *in vivo* embryonic development in the oviduct. In bovine, activin-A was reported a benefit effect depending on the timing of its addition to the culture medium (Yoshioka *et al.*, 1998a; Trigal *et al.*, 2011). Our study also sought to determine the suitability of a culture medium supplemented with activin-A for the *in vitro* culture of *in vitro* matured and fertilized prepubertal goat oocytes. The main finding of this study was that the addition of 10 ng/mL of activin-A throughout the whole embryo culture promotes embryonic development, improving the blastocyst yield (Hammami *et al.*, 2014). Otherwise, we should also consider the possibility of an accumulative effect of activin-A added during *in vitro* maturation and culture medium. Consequently, greater effects are observed as the *in vitro* culture is longer. Thus, further studies will be required to determine the functions of activin-A during these stages of development. The development-enhancing effect of activin-A may be associated with the timing of the activation of appropriate genes within the embryonic genome. Studies in mouse (Lu *et al.*, 1993) and bovine (Yoshioka *et al.*, 1998b) showed that activin-A and activin receptors are detected in embryos from the zygote to the morula stage, suggesting that the protein might play a role in embryogenesis during the early stage of embryo development. As far as we know, no studies on detection of the activin receptors in goat oocytes and embryos have been previously reported, though studies using other species confirm the presence of them in different stages of development. A detailed profiling of

the activin-A effects on embryonic development of prepubertal goat oocytes and determination of gene expression and protein synthesis for activin-A require further experiments.

Finally, the effect of growth medium during IVM on blastocyst quality was measured by post-warming survival following vitrification method and blastocyst cell count. The ability of blastocysts derived from big oocytes to survive the vitrification/warming process at 3h was lower than that of blastocysts obtained from small oocytes. According to previous data in bovine (Morató *et al.*, 2010), these differences in survival rates of *in vitro*-produced blastocysts depend on their developmental stage. So, the decrease in the cryosurvival rate may be explained by the developmental stage of blastocysts derived from large oocytes (hatching and hatched) as compared to the stage of development of the blastocysts obtained from small oocytes (early and expanded). Otherwise, at 72h the re-expansion rate was significantly higher in the blastocysts obtained from small oocytes matured in growth medium than in those blastocysts from oocytes matured in conventional medium, reflecting a higher oocyte quality. Previous studies in bovine showed that the culture medium supplementation (Rizos *et al.*, 2001; Gómez *et al.*, 2008) affects intrinsic embryo quality in terms of cryosurvival. Thus, a higher oocyte quality was showed after the oocyte culture in growth medium. However, our results showed that the use of growth medium did not improve the blastocyst cell counts. Similarly, cell number was unaffected by the presence of activin-A in the IVM media.

So, in summary, we can conclude that the addition of ascorbic acid, insulin-transferrin-selenium and a low level of hormones during the first 12h of IVM of prepubertal goat oocytes would be useful to comply with the requirements of the intrinsic quality of blastocysts produced *in vitro*. Notwithstanding, the capacity of those oocytes to develop to the blastocyst stage *in vitro* seems to be more related to oocyte diameter than to the conditions of the IVM system. Finally, according to the results obtained in the second study, the addition of 10 ng/mL of activin-A throughout the whole embryo culture improves blastocyst yield.

GENERAL CONCLUSIONS

General Conclusions

The results of this PhD thesis let conclude that:

1. The addition of ascorbic acid, insulin-transferrin-selenium and a low level of hormones during the first 12h of IVM of prepubertal goat oocytes would be useful to comply with the requirements of the intrinsic quality of blastocysts produced *in vitro*.
2. The capacity of prepubertal goat oocytes to develop to the blastocyst stage *in vitro* seems to be more related to oocyte diameter than to the conditions of the IVM system.
3. Activin-A added to IVM did not result in an improvement of blastocyst development.
4. The addition of 10 ng/mL of activin-A throughout the whole embryo culture period promotes embryonic development, showing an improving effect on the blastocyst yield of prepubertal goat oocytes.

REFERENCES

References

Abdoon A, Kandil O, Otoi T, Suzuki T, 2001: Influence of oocyte quality, culture media and gonadotropins on cleavage rate and development of in vitro fertilized buffalo embryos. *Animal Reproduction Science* 65 215-223.

Abe H, Yamashita S, Itoh T, Satoh T, Hoshi H, 1999: Ultrastructure of bovine embryos developed from in vitro-matured and -fertilized oocytes: comparative morphological evaluation of embryos cultured either in serum-free medium or in serum-supplemented medium. *Molecular Reproduction and Development* 53 325-35.

Agarwal A, Said T, Bedaiwy M, Banerjee J, Alvarez J, 2006: Oxidative stress in an assisted reproductive techniques setting. *Fertility and Sterility* 86 503-512.

Alak B, Smith G, Woodruff T, Stouffer R, Wolf D, 1996: Enhancement of primate oocyte maturation and fertilization in vitro by inhibin A and activin A. *Fertility and Sterility* 66 646-653.

Alak B, Coskun S, Friedman C, Kennard E, Kim M, Seifer D, 1998: Activin A stimulates meiotic maturation of human oocytes and modulates granulosa cell steroidogenesis in vitro. *Fertility and Sterility* 70 1126-1130.

Albano RM, Groome N, Smith JC, 1993: Activins are expressed in preimplantation mouse embryos and in ES and EC cells and are regulated on their differentiation. *Development* 117 711-23.

Ali A, Bilodeau J, Sirard M, 2003: Antioxidant requirements for bovine oocytes varies during in vitro maturation, fertilization and development. *Theriogenology* 59 939-949.

Ali A, Coenen K, Bousquet D, Sirard MA, 2004: Origin of bovine follicular fluid and its effect during in vitro maturation on the developmental competence of bovine oocytes. *Theriogenology* 62 1596-1606.

Alvarez G, Dalvit G, Achi M, Miguez M, Cetica P, 2009: Immature oocyte quality and maturational competence of porcine cumulus-oocyte complexes subpopulations. *Biocell* 33 167-177.

Anguita B, Jiménez-Macedo A, Izquierdo D, Mogas T, Paramio M, 2007: Effect of oocyte diameter on meiotic competence, embryo development, p34 (cdc2) expression and MPF activity in prepubertal goat oocytes. *Theriogenology* 67 526-536.

Anguita B, Paramio M, Jiménez-Macedo A, Morató R, Mogas T, Izquierdo D, 2008: Total RNA and protein content, Cyclin B 1 expression and developmental competence of prepubertal goat oocytes. *Animal Reproduction Science* 103 290-303.

Armstrong D, Kotaras P, Earl C, 1997: Advances in production of embryos in vitro from juvenile and prepubertal oocytes from the calf and lamb. *Reproduction Fertility and Development* 9 333-339.

References

- Bai J, Hou J, Guan H, Yan F, Cui X, Liu L, Wang S, An X, 2008: Effect of 2-mercaptoethanol and cysteine supplementation during in vitro maturation on the developmental competence of oocytes from hormone-stimulated lambs. *Theriogenology* 70 758-764.
- Baldassarre H, Wang B, Kafidi N, Keefer C, Lazaris A, Karatzas C, 2002: Advances in the production and propagation of transgenic goats using laparoscopic ovum pick-up and in vitro embryo production technologies. *Theriogenology* 57 275-284.
- Baldassarre H, Karatzas C, 2004: Advanced assisted reproduction technologies (ART) in goats. *Animal Reproduction Science* 82-83 255-266.
- Bavister BD, 1995. Culture of preimplantation embryos: facts and artifacts. *Human Reproduction* 1 91-148.
- Beilby KH, de Graaf SP, Evans G, Maxwell WMC, Wilkening S, Wrenzycki C, Grupen CG, 2011: Quantitative mRNA expression in ovine blastocysts produced from X- and Y-chromosome bearing sperm, both in vitro and in vivo. *Theriogenology* 76 471-481.
- Beker-van Woudenberg A, van Tol H, Roelen B, Colenbrander B, Bevers M, 2004: Estradiol and its membrane-impermeable conjugate (estradiol-bovine serum albumin) during in vitro maturation of bovine oocytes: Effects on nuclear and cytoplasmic maturation, cytoskeleton, and embryo quality. *Biology of Reproduction* 70 1465-1474.
- Betts DH, King WA, 2001: Genetic regulation of embryo death and senescence. *Theriogenology* 55 171-191
- Bevers M, Dieleman S, vandenHurk R, Izadyar F, 1997: Regulation and modulation of oocyte maturation in the bovine. *Theriogenology* 47 13-22.
- Bilezikjian L, Blount A, Donaldson C, Vale W, 2006: Pituitary actions of ligands of the TGF-beta family: activins and inhibins. *Reproduction* 132 207-215.
- Bing Y, Nagai T, Rodriguez-Martinez H, 2001: Effects of cysteamine, FSH and estradiol-17 beta on in vitro maturation of porcine oocytes. *Theriogenology* 55 867-876.
- Blondin P, Sirard MA, 1995: Oocyte and follicular morphology as determining characteristics for developmental competence in bovine oocytes. *Molecular Reproduction and Development* 41 54-62.
- Booth PJ, Holm P, Callesen H, 2005: The effect of oxygen tension on porcine embryonic development is dependent on embryo type. *Theriogenology* 63 2040-2052.
- Bormann C, Onger E, Krisher R, 2003: The effect of vitamins during maturation of caprine oocytes on subsequent developmental potential in vitro. *Theriogenology* 59 1373-1380.

References

- Bowles C, Lishman A, 1998: Attempts to improve the yield of bovine blastocysts by incorporating insulin, selenium and transferrin in the in vitro system. *South African Journal of Animal Science* 28 30-37.
- Brevini T, Cillo F, Antonini S, Tosetti V, Gandolfi F, 2007: Temporal and spatial control of gene expression in early embryos of farm animals. *Reproduction Fertility and Development* 19 35-42.
- Brison D, Schultz R, 1997: Apoptosis during mouse blastocyst formation: Evidence for a role for survival factors including transforming growth factor alpha. *Biology of Reproduction* 56 1088-1096.
- Buettner G, 1993: The pecking order of free-radicals and antioxidants: lipid-peroxidation, alpha-tocopherol, and ascorbate. *Archives of Biochemistry and Biophysics* 300 535-543.
- Byskov A, Andersen C, Hossaini A, Xia G, 1997: Cumulus cells of oocyte-cumulus complexes secrete a meiosis-activating substance when stimulated with FSH. *Molecular Reproduction and Development* 46 296-305.
- Catalá M, Izquierdo D, Uzbekova S, Morató R, Roura M, Romaguera R, Papillier P, Paramio M, 2011: Brilliant cresyl blue stain selects largest oocytes with highest mitochondrial activity, maturation-promoting factor activity and embryo developmental competence in prepubertal sheep. *Reproduction* 142 517-527.
- Catalá M, Montserrat R, Izquierdo D, Morató R, Hammami S, Paramio MT, 2013: Blastocyst development, MPF activity and ATP content of lamb oocytes supplemented with insulin-transferrin-selenium (ITS) and ascorbic acid at IVM. *Small Ruminant Research* 112 103-107.
- Cebrian-Serrano A, Salvador I, Raga E, Dinnyes A, Silvestre MA, 2013: Beneficial effect of melatonin on blastocyst in vitro production from heat-stressed bovine oocytes. *Reproduction in Domestic Animals* 48 738-746.
- Cetica P, Pintos L, Dalvit G, Beconi M, 2001: Antioxidant enzyme activity and oxidative stress in bovine oocyte in vitro maturation. *IUBMB Life* 51 57-64.
- Chamnongpol S, Dodson W, Cromie MJ, Harris ZL, Groisman EA, 2002: Fe(III)-mediated cellular toxicity. *Molecular Microbiology* 45 711-719.
- Choe C, Shin YW, Kim EJ, Cho SR, Kim HJ, Choi SH, Han MH, Han J, Son DS, Kang D, 2010: Synergistic effects of glutathione and β -mercaptoethanol treatment during in vitro maturation of porcine oocytes on early embryonic development in a culture system supplemented with L-cysteine. *Journal of Reproduction and Development* 56 575-582.
- Choi W, Banerjee J, Falcone T, Bena J, Agarwal A, Sharma R, 2007: Oxidative stress and tumor necrosis factor-alpha-induced alterations in metaphase II mouse oocyte spindle structure. *Fertility and Sterility* 88 1220-1231.

References

- Cognié Y, Benoit F, Poulin N, Khatir H, Driancourt MA, 1998: Effect of follicle size and of the Fec (B) Booroola gene on oocyte function in sheep. *Journals of Reproduction and Fertility* *112* 379-386.
- Cognié Y, 1999: State of the art in sheep-goat embryo transfer. *Theriogenology* *51* 105-116.
- Cognié Y, Baril G, Poulin N, Mermillod P, 2003: Current status of embryo technologies in sheep and goat. *Theriogenology* *59* 171-188.
- Cognié Y, Poulin N, Locatelli Y, Mermillod P, 2004: State-of-the-art production, conservation and transfer of in-vitro-produced embryos in small ruminants. *Reproduction Fertility and Development* *16* 437-445.
- Córdova B, Morató R, Izquierdo D, Paramio T, Mogas T, 2010: Effect of the addition of insulin-transferrin-selenium and/or L-ascorbic acid to the in vitro maturation of prepubertal bovine oocytes on cytoplasmic maturation and embryo development. *Theriogenology* *74* 1341-1348.
- Córdova B, Morató R, De Frutos C, Bermejo-Alvarez P, Paramio T, Gutierrez-Adan A, Mogas T, 2011: Effect of leptin during in vitro maturation of prepubertal calf oocytes: Embryonic development and relative mRNA abundances of genes involved in apoptosis and oocyte competence. *Theriogenology* *76* 1706-1715.
- Coskun S, Lin Y, 1994: Effects of Transforming Growth-Factors and activin A on in-vitro porcine oocyte maturation. *Molecular Reproduction and Development* *38* 153-159.
- Cox J, Avila J, Saravia F, Maria A, 1994: Assessment of fertilizing ability of goat spermatozoa by in-vitro fertilization of cattle and sheep intact oocytes. *Theriogenology* *41* 1621-1629.
- Crozet N, Ahmedali M, Dubos MP, 1995: Developmental competence of goat oocytes from follicles of different size categories following maturation, fertilization and culture in vitro. *Journals of Reproduction and Fertility* *103* 293-298.
- Damiani P, Fissore R, Cibelli J, Long C, Balise J, Robl J, DUBY R, 1996: Evaluation of developmental competence, nuclear and ooplasmic maturation of calf oocytes. *Molecular Reproduction and Development* *45* 521-534.
- Dang-Nguyen T, Kaneda M, Somfai T, Haraguchi S, Matsukawa K, Akagi S, Kikuchi K, Nakai M, Nguyen B, Tajima A, Kanai Y, Nagai T, 2011: Development of single blastomeres derived from two-cell embryos produced in vitro in pigs. *Theriogenology* *76* 88-96.
- De la Fuente R, King W, 1997: Use of a chemically defined system for the direct comparison of inner cell mass and trophoctoderm distribution in murine, porcine and bovine embryos. *Zygote* *5* 309-320.

References

- De Matos DG, Furnus CC, Moses DF, Baldassarre H, 1995: Effect of cysteamine on glutathione level and developmental capacity of bovine oocyte matured in vitro. *Molecular Reproduction and Development* 42 432-436.
- De Matos DG, Herrera C, Cortvrindt R, Smits J, Van Soom A, Nogueira D, Pasqualini RS, 2002: Cysteamine supplementation during in vitro maturation and embryo culture: a useful tool for increasing the efficiency of bovine in vitro embryo production. *Molecular Reproduction and Development* 62 203-209.
- De Smedt V, Crozet N, Ahmedali M, Martino A, Cognié Y, 1992: In vitro maturation and fertilization of goat oocytes. *Theriogenology* 37 1049-1060.
- De Smedt V, Crozet N, Gall L, 1994: Morphological and functional-changes accompanying the acquisition of meiotic competence in ovarian goat oocyte. *Journal of Experimental Zoology* 269 128-139.
- De Souza J, Duffard N, Bertoldo M, Locatelli Y, Corbin E, Fatet A, Freitas V, Vicente J., Mermillod P, 2013: Influence of heparin or the presence of cumulus cells during fertilization on the in vitro production of goat embryos. *Animal Reproduction Science* 138 82-89.
- Earl C, Irvine B, Armstrong D, 1994: Development of techniques for the production of viable embryos from 6 to 7 week old lambs. *Animal Production in Australia* 20 428.
- Ebert RM, Ulmer S, Zeck J, Meissner-Weigl D, Schneider H, Stopper N, Schupp M, Kassem F, 2006: Selenium supplementation restores the antioxidative capacity and prevents cell damage in bone marrow stromal cells in vitro. *Stem Cells* 24 1226 –1235.
- Edwards R, 1965: Maturation in vitro of mouse, sheep, cow, pig, rhesus monkey and human ovarian oocytes. *Nature* 208 349-351.
- Endo M, Kawahara-Miki R, Cao F, Kimura K, Kuwayama T, Monji Y, Iwata H, 2013: Estradiol supports in vitro development of bovine early antral follicles. *Reproduction* 145 85-96.
- Eppig J, Schroeder A, 1989: Capacity of mouse oocytes from preantral follicles to undergo embryogenesis and development to live young after growth, maturation, and fertilization in vitro. *Biology of Reproduction* 41 268-276.
- Eppig J, 1996: Coordination of nuclear and cytoplasmic oocyte maturation in eutherian mammals. *Reproduction, Fertility and Development* 8 485-489.
- Eppig J, Hosoe M, O'Brien M, Pendola F, Requena A, Watanabe S, 2000: Conditions that affect acquisition of developmental competence by mouse oocytes in vitro: FSH, insulin, glucose and ascorbic acid. *Molecular and Cellular Endocrinology* 163 109-116.
- Eramaa M, Hilden K, Tuuri T, Ritvos O, 1995: Regulation of inhibin activin subunit messenger ribonucleic-acids (messenger-rnas) by Activin-a and expression of Activin

References

receptor messenger-rnas in cultured human granulosa-luteal cells. *Endocrinology* *136* 4382-4389.

Fatehi A, Roelen B, Colenbrander B, Schoevers E, Gadella B, Bevers M, van den Hurk R, 2005: Presence of cumulus cells during in vitro fertilization protects the bovine oocyte against oxidative stress and improves first cleavage but does not affect further development. *Zygote* *13* 177-185.

Fischer B, Bavister BD, 1993: Oxygen tension in the oviduct and uterus of rhesus monkeys, hamsters and rabbits. *Journal of Reproduction and Fertility* *99* 673-679.

Fouladi-Nashta A, Alberio R, Kafi M, Nicholas B, Campbell K, Webb R, 2005: Differential staining combined with TUNEL labelling to detect apoptosis in preimplantation bovine embryos. *Reproductive BioMedicine Online* *10* 497-502.

Fukaya T, Yamanaka T, Terada Y, Murakami T, Yajima A, 1998: Growth hormone improves mouse embryo development in vitro, and the effect is neutralized by growth hormone receptor antibody. *The Tohoku Journal of Experimental Medicine* *184* 113-122.

Funahashi H, Day B, 1993: Effects of the duration of exposure to hormone supplements on cytoplasmic maturation of pig oocytes in vitro. *Journals of Reproduction and Fertility* *98* 179-185.

Gall L, Chene N, Dahirel M, Ruffini S, Boulesteix C, 2004: Expression of epidermal growth factor receptor in the goat cumulus-oocyte complex. *Molecular Reproduction and Development* *67* 439-445.

Gandolfi F, Modina S, Brevini T, Passoni L, Artini P, Petraglia F, Lauria A, 1995: Activin beta (A) subunit is expressed in bovine oviduct. *Molecular Reproduction and Development* *40* 286-291.

Gandolfi F, Milanesi E, Pocar P, Luciano A, Brevini T, Acocella F, Lauria A, Armstrong D, 1998: Comparative analysis of calf and cow oocytes during in vitro maturation. *Molecular Reproduction and Development* *49* 168-175.

Gandolfi F, Brevini TA, Cillo F, Antonini S, 2005: Cellular and molecular mechanisms regulating oocyte quality and the relevance for farm animal reproductive efficiency. *Revue scientifique et technique* *24* 413-423.

Garcia-Garcia RM, Ward F, Fair S, O'Meara CM, Wad M, Duffy P, Lonergan P, 2007: Development and quality of sheep embryos cultured in commercial G1.3/G2.3 sequential media. *Animal Reproduction Science* *98* 233-240.

Gardner DK, Lane M, Spitzer A, Batt PA, 1994: Enhanced rates of cleavage and development for sheep zygotes cultured to the blastocyst stage in vitro in the absence of serum and somatic cells: amino acids, vitamins, and culturing embryos in groups stimulate development. *Biology of Reproduction* *50* 390-400.

References

- Głabowski W, Kurzawa R, Wiszniewska B, Baczkowski T, Marchlewicz M, Brelik P, 2005: Growth factors effects on preimplantation development of mouse embryos exposed to tumor necrosis factor alpha. *Reproductive Biology* 5 83-99.
- Gómez E, Rodriguez A, Munoz M, Caamano JN, Hidalgo CO, Moran E, Facal N, Diez C, 2008: Serum free embryo culture medium improves in vitro survival of bovine blastocysts to vitrification. *Theriogenology* 69 1013-1021.
- Grazul-Bilska AT, Choi JT, Bilski JJ, Weigl RM, Kirsch JD, Kraft KC, Reynolds LP, Redmer DA, 2003: Effects of epidermal growth factor on early embryonic development after in vitro fertilization of oocytes collected from ewes treated with follicle stimulating hormone. *Theriogenology* 59 1449-1457.
- Gruppen CG, Nagashima H, Nottle MB, 1997: Role of epidermal growth factor and insulin-like growth factor-I on porcine oocyte maturation and embryonic development in vitro. *Reproduction Fertility and Development* 9 571-575.
- Guler A, Poulin N, Mermillod P, Terqui M, Cognié Y, 2000: Effect of growth factors, EGF and IGF-I, and estradiol on in vitro maturation of sheep oocytes. *Theriogenology* 54 209-218.
- Gupta, PSP, Ravindra, JP, Girish Kumar V, Raghu HM, Nandi S, 2005: Stimulation of in vitro oocyte maturation with a novel peptide isolated from follicular fluid of the buffalo (*Bubalus bubalis*). *Small Ruminant Research* 59 33-40.
- Gutiérrez-Adán A, Lonergan P, Rizos D, Ward F, Boland M, Pintado B, de la Fuente J, 2001: Effect of the in vitro culture system on the kinetics of blastocyst development and sex ratio of bovine embryos. *Theriogenology* 55 1117-1126.
- Halliwell B, Gutteridge JMC, 1989: *Free Radicals in Biology and Medicine*, Ed 2. Clarendon Press, Oxford.
- Hammami S, Morató R, Romaguera R, Roura M, Catalá M, Paramio M, Mogas T, Izquierdo D, 2013: Developmental competence and embryo quality of small oocytes from pre-pubertal goats cultured in IVM medium supplemented with low level of hormones, insulin–transferrin selenium and ascorbic acid. *Reproduction in Domestic Animals* 48 339-344.
- Hammami S, Izquierdo D, Catalá M, Paramio M, Morató R, 2014: In vitro developmental competence of prepubertal goat oocytes cultured with recombinant activin-A. *Animal* 8 94-101.
- Hegde A, Behr B, 2012: Media composition: growth factors. *Methods in Molecular Biology* 912 177-198.
- Heldin C, Miyazono K, tenDijke P, 1997: TGF-beta signalling from cell membrane to nucleus through SMAD proteins. *Nature* 390 465-471.

References

- Herrick JR, Behboodi E, Memili E, Blash S, Echelard Y, Krisher RL, 2004: Effect of macromolecule supplementation during in vitro maturation of goat oocytes on developmental potential. *Molecular Reproduction and Development* 69 338-346.
- Homa ST, Brown CA, 1992: Changes in linoleic acid during follicular development and inhibition of spontaneous breakdown of germinal vesicles in cumulus-free bovine oocytes. *Journal of Reproduction and Fertility* 94 153-160.
- Hosseini MS, Hashem MA, Jeong YW, Lee MS, Kim S, Kim JH, Koo OJ, Park SM, Lee EG, Park SW, Kang SK, Lee BC, Hwang WS, 2007: Temporal effects of alpha-tocopherol and L-ascorbic acid on in vitro fertilized porcine embryo development. *Animal Reproduction Science* 100 107-117.
- Hosseini SM, Forouzanfar M, Hajian M, Asgari V, Abedi P, Hosseini L, Ostadhosseini S, Moulavi F, Safahani Langrroodi M, Sadeghi H, Bahramian H, Eghbalsaied Sh, Nasr-Esfahani MH, 2009: Antioxidant supplementation of culture medium during embryo development and/or after vitrification-warming; which is the most important? *Journal of Assisted Reproduction and Genetics* 26 355-364.
- Hu J, Ma X, Bao J, Li W, Cheng D, Gao Z, Lei A, Yang C, Wang H, 2011: Insulin-transferrin-selenium (ITS) improves maturation of porcine oocytes in vitro. *Zygote* 19 191-197.
- Huang Y, Tang X, Xie W, Zhou Y, Li D, Zhou Y, Zhu J, Yuan T, Lai L, Pang D, Ouyang H, 2011: Vitamin C enhances in vitro and in vivo development of porcine somatic cell nuclear transfer embryos. *Biochem Biophys Res Commun* 411 397-401.
- Hulshof S, Figueiredo J, Beckers J, Bevers M, Vanderstichele H, VandenHurk R, 1997: Bovine preantral follicles and activin: Immunohistochemistry for activin and activin receptor and the effect of bovine activin A in vitro. *Theriogenology* 48 133-142.
- Itoh M, Igarashi M, Yamada K, Hasegawa Y, Seki M, Eto Y, Shibai H, 1990: Activin A stimulates meiotic maturation of the rat oocyte in vitro. *Biochemical and Biophysical Research Communications* 166 1479-1484.
- Iwasaki S, Yoshida N, Ushijima H, Watanabe S, Nakahara T, 1990: Morphology and proportion of inner cell mass of bovine blastocysts fertilized in vitro and in vivo. *Journals of Reproduction and Fertility* 90 279-284.
- Izadyar F, Zeinstra E, Colenbrander B, Vanderstichele H, Bevers M, 1996: In vitro maturation of bovine oocytes in the presence of bovine activin A does not affect the number of embryos. *Animal Reproduction Science* 45 37-45.
- Izadyar F, Dijkstra G, Van Tol H, van de Eljnden-Van Raaij A, Van den Hurk R, Colenbrander B, Bevers M, 1998: Immunohistochemical localization and mRNA expression of activin, inhibin, follistatin, and activin receptor in bovine cumulus-oocyte

References

complexes during in vitro maturation. *Molecular Reproduction and Development* 49 186-195.

Izquierdo D, Villamediana P, Palomo M, Mogas T, Paramio M, 1998: Effect of sperm capacitation and fertilization media on IVF and early embryo development of prepubertal goat oocytes. *Theriogenology* 49 1501-1513.

Izquierdo D, Villamediana P, Lopez-Bejar M, Paramio M, 2002: Effect of in vitro and in vivo culture on embryo development from prepubertal goat IVM-IVF oocytes. *Theriogenology* 57 1431-1441.

Jaakma U, Zhang B, Larsson B, Niwa K, RodriguezMartinez H, 1997: Effects of sperm treatments on the in vitro development of bovine oocytes in semidefined and defined media. *Theriogenology* 48 711-720.

Jeong Y, Hossein M, Bhandari D, Kim YW, Kim J, Park S, Lee E, Park S, Jeong Y, Lee J, Kim S, Hwang W, 2008: Effects of insulin-transferrin-selenium in defined and porcine follicular fluid supplemented IVM media on porcine IVF and SCNT embryo production. *Animal Reproduction Science* 106 13-24.

Jiménez-Macedo A, Izquierdo D, Anguita B, Paramio M, 2005: Comparison between intracytoplasmic sperm injection and in vitro fertilisation employing oocytes derived from prepubertal goats. *Theriogenology* 64 1249-1262.

Jiménez-Macedo A, Anguita B, Izquierdo D, Mogas T, Paramio M, 2006: Embryo development of prepubertal goat oocytes fertilised by intracytoplasmic sperm injection (ICSI) according to oocyte diameter. *Theriogenology* 66 1065-1072.

Jiménez-Macedo AR, Paramio MT, Anguita B, Morató R, Romaguera R, Mogas T, Izquierdo D, 2007: Effect of ICSI and embryo biopsy on embryo development and apoptosis according to oocyte diameter in prepubertal goats. *Theriogenology* 67 1399-1408.

Karja NW, Wongsrikeao P, Murakami M, Agung B, Fahrudin M, Nagai T, Otoi T, 2004: Effects of oxygen tension on the development and quality of porcine in vitro fertilized embryos. *Theriogenology* 62 1585-1595.

Katska L, Kania G, Rynska B, Gajda B, Smorag Z, 2002: Offspring after transferring goat embryos originating from in vitro matured and fertilised oocytes. *Med.Weter.* 58 462-463.

Katska-Ksiazkiewicz L, Rynska B, Gajda B, Smorag Z, 2004: Effect of donor stimulation, frozen semen and heparin treatment on the efficiency of in vitro embryo production in goats. *Theriogenology* 62 576-586.

Katska-Ksiazkiewicz L, Opiela J, Rynska B, 2007: Effects of oocyte quality, semen donor and embryo co-culture system on the efficiency of blastocyst production in goats. *Theriogenology* 68 736-744.

References

- Kauffold J, Amer HAH, Bergfeld U, Weber W, Sobiraj A, 2005: The in vitro developmental competence of oocytes from juvenile calves is related to follicular diameter. *Journal of Reproduction and Development* 51 325-332.
- Keskintepe L, Darwish G, Younis A, Brackett B, 1994: In vitro development of morulae from immature caprine oocytes. *Zygote* 2 97-102.
- Keskintepe L, Simplicio A, Brackett B, 1998: Caprine blastocyst development after in vitro fertilization with spermatozoa frozen in different extenders. *Theriogenology* 49 1265-1274.
- Khatir H, Lonergan P, Mermillod P, 1998: Kinetics of nuclear maturation and protein profiles of oocytes from prepubertal and adult cattle during in vitro maturation. *Theriogenology* 50 917-929.
- Khatun M, Bhuiyan M, Ahmed J, Haque A, Rahman M, Shamsuddin M, 2011: In vitro maturation and fertilization of prepubertal and pubertal black Bengal goat oocytes. *Journal of Veterinary Science* 12 75-82.
- Kitagawa Y, Suzuki K, Yoneda A, Watanabe T, 2004: Effects of oxygen concentration and antioxidants on the in vitro developmental ability, production of reactive oxygen species (ROS), and DNA fragmentation in porcine embryos. *Theriogenology* 62 1186-1197.
- Knight P, Glister C, 2003: Local roles of TGF-beta superfamily members in the control of ovarian follicle development. *Animal Reproduction Science* 78 165-183.
- Kniijn H, Gjorret J, Vos P, Hendriksen P, Van der Weijden B, Maddox-Hyttel P, Dieleman S, 2003: Consequences of in vivo development and subsequent culture on apoptosis, cell number, and blastocyst formation in bovine embryos. *Biology of Reproduction* 69 1371-1378.
- Koeman J, Keefer C, Baldassarre H, Downey B, 2003: Developmental competence of prepubertal and adult goat oocytes cultured in semi-defined media following laparoscopic recovery. *Theriogenology* 60 879-889.
- Korhonen K, Julkunen H, Kananen K, Bredbacka P, Tiirikka T, Rätty M, Vartia K, Kaimio I, Kontinen A, Halmekytö M, Vilkki J, Peippo J, Lindeberg H, 2012: The effect of ascorbic acid during biopsy and cryopreservation on viability of bovine embryos produced in vivo. *Theriogenology* 77 201-205.
- Krisher RL, 2004: The effect of oocyte quality on development. *Journal of Animal Science* 82 14-23.
- Lane M, Maybach JM, Gardner DK, 2002: Addition of ascorbate during cryopreservation stimulates subsequent embryo development. *Human Reproduction* 17 2686-2693.

References

Ledda S, Bogliolo L, Calvia P, Leoni G, Naitana S, 1997: Meiotic progression and developmental competence of oocytes collected from juvenile and adult ewes. *Journals of Reproduction and Fertility* 109 73-78.

Ledda S, Bogliolo L, Leoni G, Naitana S, 1999a: Production and lambing rate of blastocysts derived from in vitro matured oocytes after gonadotropin treatment of prepubertal ewes. *Journal of Animal Science* 77 2234-2239.

Ledda S, Bogliolo L, Leoni G, Naitana S, 1999b: Follicular size affects the meiotic competence of in vitro matured prepubertal and adult oocytes in sheep. *Reproduction Nutrition Development* 39 503-508.

Lee K, Bettgowda A, Wee G, Ireland J, Smith G, 2009: Molecular determinants of oocyte competence: Potential functional role for maternal (Oocyte-Derived) follistatin in promoting bovine early embryogenesis. *Endocrinology* 150 2463-2471.

Lee KS, Kim EY, Jeon K, Cho SG, Han YJ, Yang BC, Lee SS, Ko MS, Riu KJ, Lee HT, Park SP, 2011: 3,4-Dihydroxyflavone acts as an antioxidant and antiapoptotic agent to support bovine embryo development in vitro. *Journal of Reproduction and Development* 57 127-134.

Leibo SP, Loskutoff N.M, 1993: Cryobiology of in vitro-derived bovine embryos. *Theriogenology* 39 81-94

Leoni GG, Rosati I, Succu S, Bogliolo L, Bebbere D, Berlinguer F, Ledda S, Naitana S, 2007: A low oxygen atmosphere during IVF accelerates the kinetic of formation of in vitro produced ovine blastocysts. *Reproduction in Domestic Animals* 42 299-304.

Leoni GG, Succu S, Satta V, Paolo M, Bogliolo L, Bebbere D, Spezzigu A, Madeddu M, Berlinguer F, Ledda S, Naitana S, 2009: In vitro production and cryotolerance of prepubertal and adult goat blastocysts obtained from oocytes collected by laparoscopic oocyte-pick-up (LOPU) after FSH treatment. *Reproduction Fertility and Development* 21 901-908.

Liu S, Feng H, Marchesi D, Chen Z, Hershlag A, 2011: Dose-dependent effects of gonadotropin on oocyte developmental competence and apoptosis. *Reproduction Fertility and Development* 23 990-996.

Liu Y, Sakaeda T, Takara K, Nakamura T, Ohmoto N, Komoto C, Kobayashi H, Yagami T, Okamura N, Okumura K, 2003: Effects of reactive oxygen species on cell proliferation and death in HeLa cells and its MDR1-overexpressing derivative cell line. *Biological and Pharmaceutical Bulletin*. 26 278-281.

Lonergan P, Monaghan P, Rizos D, Boland M, Gordon I, 1994: Effect of follicle size on bovine oocyte quality and developmental competence following maturation, fertilization, and culture in-vitro. *Molecular Reproduction and Development* 37 48-53.

References

- Lonergan P, O'Kearney-Flynn M, Boland M, 1999: Effect of protein supplementation and presence of an antioxidant on the development of bovine zygotes in synthetic oviduct fluid medium under high or low oxygen tension. *Theriogenology* 51 1565-1576.
- Lonergan P, Rizos D, Gutierrez-Adan A, Fair T, Boland M, 2003: Oocyte and embryo quality: Effect of origin, culture conditions and gene expression patterns. *Reproduction in Domestic Animals* 38 259-267.
- Lu R, Matsuyama S, Nishihara M, Takahashi M, 1993: Developmental expression of activin/inhibin beta A, beta B, and alpha subunits, and activin receptor-IIB genes in preimplantation mouse embryos. *Biology of Reproduction* 49 1163-1169.
- Lv L, Yue W, Liu W, Ren Y, Li F, Lee K, Smith G, 2010: Effect of oocyte selection, estradiol and antioxidant treatment on in vitro maturation of oocytes collected from prepubertal Boer goats. *Italian Journal of Animal Science* 9 50-54.
- Magalhães-Padilha DM, Duarte AB, Araújo VR, Saraiva MV, Almeida AP, Rodrigues GQ, Matos MH, Campello CC, Silva JR, Gastal MO, Gastal EL, Figueiredo JR, 2012: Steady-state level of insulin-like growth factor-I (IGF-I) receptor mRNA and the effect of IGF-I on the in vitro culture of caprine preantral follicles. *Theriogenology* 77 206-213.
- Makarevich AV, Markkula M, 2002: Apoptosis and cell proliferation potential of bovine embryos stimulated with insulin-like growth factor I during in vitro maturation and culture. *Biology of Reproduction* 66 386-392.
- Malik R, Lohan I, 1999: Peritoneal fluid from rabbits or goats as media for in vitro maturation, fertilization and initial culture of caprine oocytes. *Animal Reproduction Science* 54 195-201.
- Mamo S, Rizos D, Lonergan P, 2012: Transcriptomic changes in the bovine conceptus between the blastocyst stage and initiation of implantation. *Animal Reproduction Science* 134 56-63.
- Manjunatha BM, Devaraj M, Gupta PS, Ravindra JP, Nandi S, 2009: Effect of taurine and melatonin in the culture medium on buffalo in vitro embryo development. *Reproduction in Domestic Animals* 44 12-16.
- Marchal R, Feugang J, Perreau C, Venturi E, Terqui M, Mermillod P, 2001: Meiotic and developmental competence of prepubertal and adult swine oocytes. *Theriogenology* 56 17-29.
- Marchal R, Vigneron C, Perreau C, Bali-Papp A, Mermillod P, 2002: Effect of follicular size on meiotic and developmental competence of porcine oocytes. *Theriogenology* 57 1523-1532.
- Marei WF, Wathes DC, Fouladi-Nashta AA, 2009: The effect of linolenic acid on bovine oocyte maturation and development. *Biology osuring follicular development and*

References

inhibition of spontaneous breakdown of germinal vesicles in cumulus-free bovine oocytes. *Journal of Reproduction and Fertility* 94 153– 160.

Marei WF, Wathes DC, Fouladi-Nashta AA, 2010. Impact of linoleic acid on bovine oocyte maturation and embryo development. *Biology of Reproduction*,139 979-988.

Martino A, Palomo M, Mogas T, Paramio M, 1994: Influence of the collection technique of prepubertal goat oocytes on in vitro maturation and fertilization. *Theriogenology* 42 859-873.

Martino A, Mogas T, Palomo M, Paramio M, 1995: In vitro maturation and fertilization of prepubertal goat oocytes. *Theriogenology* 43 473-485.

Mather J, Moore A, Li R, 1997: Activins, inhibins, and follistatins: Further thoughts on a growing family of regulators. *Proceedings of the Society for Experimental Biology and Medicine* 215.

Matsui M, Takahashi Y, Hishinuma M and Kanagawa H 1995: Insulin and insulin-like growth factor-I (IGF-I) stimulate the development of bovine embryos fertilized in vitro. *Journal of Veterinary Medical Science* 57 1109–1111.

Mattioli M, Bacci M, Galeati G, Seren E, 1991: Effects of LH and FSH on the maturation of pig oocytes In vitro. *Theriogenology* 36 95-105.

Men H, Monson R, Rutledge J, 2002: Effect of meiotic stages and maturation protocols on bovine oocyte's resistance to cryopreservation. *Theriogenology* 57 1095-1103.

Mermillod P, Oussaid B, Cognié Y, 1999: Aspects of follicular and oocyte maturation that affect the developmental potential of embryos. *Journal of Reproduction and Fertility*.Supplement 54 449-460.

Mermillod P, Locatelli Y, Dalbiès-Tran R, Uzbekova S, Baril G, Guignot F, Perreau C, Poulin N, Touzé JL, Pennetier S, Schmaltz B, Cognié Y, 2006: In vitro production of ruminant embryos: results, limits and perspectives. *Symposium COA/INRA Scientific Cooperation in Agriculture* 59-78.

Mertens M, Lopez-Bejar M, Paramio M, 2005: Intracytoplasmic glutathione levels in heifer oocytes cultured in different maturation media and its effect on embryo development. *Reproduction in Domestic Animals* 40 126-130.

Miclea I, Pacală N, Hettig A, Zăhan M, Miclea V, 2012: Alpha-tocopherol and Ascorbic Acid Combinations Influence the Maturation of Sheep Oocytes. *Scientific Papers: Animal Science and Biotechnologies* 45 310-313.

Mizushima S, Fukui Y, 2001: Fertilizability and developmental capacity of bovine oocytes cultured individually in a chemically defined maturation medium. *Theriogenology* 55 1431-1445.

References

- Mogas T, Palomo M, Izquierdo M, Paramio M, 1997a: Developmental capacity of in vitro matured and fertilized oocytes from prepubertal and adult goats. *Theriogenology* 47 1189-1203.
- Mogas T, Palomo M, Izquierdo M, Paramio M, 1997b: Morphological events during in vitro fertilization of prepubertal goat oocytes matured in vitro. *Theriogenology* 48 815-829.
- Moor R, Trounson A, 1977: Hormonal and follicular factors affecting maturation of sheep oocytes in vitro and their subsequent developmental capacity. *Journal of Reproduction and Fertility* 49 101-109.
- Moore K, Rodríguez-Sallaberry CJ, Kramer JM, Johnson S, Wroclawska E, Goicoa S, Niasari-Naslaji A, 2007: In vitro production of bovine embryos in medium supplemented with a serum replacer: effects on blastocyst development, cryotolerance and survival to term. *Theriogenology* 68 1316-1325.
- Morató R, Izquierdo D, Paramio MT, Mogas T, 2010: Survival and apoptosis rates after vitrification in cryotop devices of in vitro-produced calf and cow blastocysts at different developmental stages. *Fertility and Development* 22 1141-1147.
- Morton KM, Maxwell WMC, Evans G, 2008: Effect of aspiration pressure during oocyte harvesting on oocyte recovery and in vitro development of ovine oocytes. *Reproduction in Domestic Animals* 43 106-110.
- Moslemi M, Tavanbakhsh S, 2011: Selenium-vitamin E supplementation in infertile men: effects on semen parameters and pregnancy rate. *International Journal of General Medicine* 4 99-104.
- Motlik J, Crozet N, Fulka J, 1984: Meiotic competence in vitro of pig oocytes isolated from early antral follicles. *Journal of Reproduction and Fertility* 72 323-328.
- Mtango N, Varisanga M, Dong Y, Rajamahendran R, Suzuki T, 2003: Growth factors and growth hormone enhance in vitro embryo production and post-thaw survival of vitrified bovine blastocysts. *Theriogenology* 59 1393-1402.
- Murray A, Swales A, Smith R, Molinek M, Hillier S, Spears N, 2008: Follicular growth and oocyte competence in the in vitro cultured mouse follicle: effects of gonadotrophins and steroids. *Molecular Human Reproduction* 14 75-83.
- Nath A, Sharma V, Dubey P, Pratheesh M, Gade N, Saikumar G, Sharma G, 2013: Impact of gonadotropin supplementation on the expression of germ cell marker genes (MATER, ZAR1, GDF9, and BMP15) during in vitro maturation of buffalo (*Bubalus bubalis*) oocyte. *In vitro Cellular and Developmental Biology-Animal* 49 34-41.
- Neuber E, Luetjens CM, Chan AW, Schatten GP, 2002: Analysis of DNA fragmentation of in vitro cultured bovine blastocysts using TUNEL. *Theriogenology* 57 2193-2202.

References

- Norwitz E, Xu S, Jeong K, Bedecarrats G, Winebrenner L, Chin W, Kaiser U, 2002: Activin A augments GnRH-mediated transcriptional activation of the mouse GnRH receptor gene. *Endocrinology* *143* 985-997.
- O'Brien J, Beck N, Maxwell W, Evans G, 1997a: Effect of hormone pre-treatment of prepubertal sheep on the production and developmental capacity of oocytes in vitro and in vivo. *Reproduction Fertility and Development* *9* 625-631.
- O'Brien J, Catt S, Ireland K, Maxwell W, Evans G, 1997b: In vitro and in vivo developmental capacity of oocytes from prepubertal and adult sheep. *Theriogenology* *47* 1433-1443.
- Olson SE, Seidel GE Jr, 2000: Culture of in vitro-produced bovine embryos with vitamin E improves development in vitro and after transfer to recipients. *Biology of Reproduction* *62* 248-252.
- Ongeri EM, Bormann CL, Butler RE, Melican D, Gavin WG, Echelard Y, Krisher RL, Behboodi E, 2001: Development of goat embryos after in vitro fertilization and parthenogenetic activation by different methods. *Theriogenology* *55* 1933-1945
- Otoi T, Yamamoto K, Koyama N, Tachikawa S, Suzuki T, 1997: Bovine oocyte diameter in relation to developmental competence. *Theriogenology* *48* 769-774.
- Palma GA, Müller M, Brem G. 1997: Effect of insulin-like growth factor I (IGF-I) at high concentrations on blastocyst development of bovine embryos produced in vitro. *Journal of Reproduction and Fertility* *110* 347-353.
- Palomo M, Izquierdo D, Mogas T, Paramio M, 1999: Effect of semen preparation on IVF of prepubertal goat oocytes. *Theriogenology* *51* 927-940.
- Paramio MT, 2010: In vivo and in vitro embryo production in goats. *Small Ruminant Research* *89* 144-148.
- Park J, Oh H, Hong S, Jang G, Kim M, Lee B, 2010: Effects of Activin A on the in vitro development and mRNA expression of bovine embryos cultured in chemically-defined two-step culture medium. *Reproduction in Domestic Animals* *45* 585-593.
- Parrish JJ, Suskoparrish JL, Leibfriedrutledge ML, Critser ES, Eyestone WH, First NL, 1986: Bovine in vitro fertilization with frozen-thawed semen. *Theriogenology* *25* 591-600.
- Pawshe C, Palanisamy A, Taneja M, Jain S, Totey S, 1996: Comparison of various maturation treatments on in vitro maturation of goat oocytes and their early embryonic development and cell numbers. *Theriogenology* *46* 971-982.
- Peluso J, 1988: Effect of the peripubertal pattern of LH and FSH secretion on in vitro oestradiol-17-Beta secretion and follicular growth within juvenile rat ovaries. *Journals of Reproduction and Fertility* *84* 239-245.

References

- Peng C, Mukai S, 2000: Activins and their receptors in female reproduction. *Biochemistry and Cell Biology-Biochimie Et Biologie Cellulaire* 78 261-279.
- Pereira R, Tuli R, Wallenhorst S, Holtz W, 2000: The effect of heparin, caffeine and calcium ionophore A 23187 on in vitro induction of the acrosome reaction in frozen-thawed bovine and caprine spermatozoa. *Theriogenology* 54 185-192.
- Phillips D, 2005: Activins, inhibins and follistatins in the large domestic species. *Domestic Animal Endocrinology* 28 1-16.
- Presicce G, Jiang S, Simkin M, Zhang L, Looney C, Godke R, Yang X, 1997: Age and hormonal dependence of acquisition of oocyte competence for embryogenesis in prepubertal calves. *Biology of Reproduction* 56 386-392.
- Raghu H, Nandi S, Reddy S, 2002a: Follicle size and oocyte diameter in relation to developmental competence of buffalo oocytes in vitro. *Reproduction Fertility and Development* 14 55-61.
- Raghu H, Nandi S, Reddy S, 2002b: Effect of insulin, transferrin and selenium and epidermal growth factor on development of buffalo oocytes to the blastocyst stage in vitro in serum-free, semidefined media. *Veterinary Record* 151 260-265.
- Rajesh N, Shankar MB, Deecaraman M, 2010: Effect of vitamin A supplementation at different gaseous environments on in vitro development of pre-implantation sheep embryos to the blastocyst stage. *Animal* 4 1884-1890.
- Rajikin M, Yusoff M, Abdullah R, 1994: Ultrastructural studies of developing goat oocytes in-vitro. *Theriogenology* 42 1003-1016.
- Revel F, Mermillod P, Peynot N, Renard J, Heyman Y, 1995: Low Developmental Capacity of In-Vitro Matured and Fertilized Oocytes from Calves Compared with that of Cows. *Journal of Reproduction and Fertility* 103 115-120.
- Rho G, Hahnel A, Betteridge K, 2001: Comparisons of oocyte maturation times and of three methods of sperm preparation for their effects on the production of goat embryos in vitro. *Theriogenology* 56 503-516.
- Rizos D, Ward F, Boland MP, Lonergan P, 2001: Effect of culture system on the yield and quality of bovine blastocysts as assessed by survival after vitrification. *Theriogenology* 56 1-16.
- Rizos D, Ward F, Duffy P, Boland M, Lonergan P, 2002a: Consequences of bovine oocyte maturation, fertilization or early embryo development in vitro versus in vivo: Implications for blastocyst yield and blastocyst quality. *Molecular Reproduction and Development* 61 234-248.
- Rizos D, Lonergan P, Boland MP, Arroyo-Garcia R, Pintado B, de la Fuente J, Gutierrez-Adan A, 2002b: Analysis of differential messenger RNA expression between

References

bovine blastocysts produced in different culture systems: implications for blastocyst quality. *Biology of Reproduction* 66 589-595

Rizos D, Clemente M, Bermejo-Alvarez P, de La Fuente J, Lonergan P, Gutierrez-Adan A, 2008: Consequences of in vitro culture conditions on embryo development and quality. *Reproduction in Domestic Animals* 43 44-50.

Roberts R, Iatropoulou A, Ciantar D, Stark J, Becker D, Franks S, Hardy K, 2005: Follicle-stimulating hormone affects metaphase I chromosome alignment and increases aneuploidy in mouse oocytes matured in vitro. *Biology of Reproduction* 72 107-118.

Roberts V, Barth S, Elroey A, Yen S, 1993: Expression of Inhibin/activin subunits and follistatin messenger ribonucleic-acids and proteins in ovarian follicles and the corpus-luteum during the human menstrual-cycle. *Journal of Clinical Endocrinology & Metabolism* 77 1402-1410.

Rodriguez-Gonzalez E, Lopez-Bejar M, Velilla E, Paramio M, 2002: Selection of prepubertal goat oocytes using the brilliant cresyl blue test. *Theriogenology* 57 1397-1409.

Rodriguez-Gonzalez E, Lopez-Bejar M, Mertens M, Paramio M, 2003: Effects on in vitro embryo development and intracellular glutathione content of the presence of thiol compounds during maturation of prepubertal goat oocytes. *Molecular Reproduction and Development* 65 446-453

Romaguera R, Casanovas A, Morató R, Izquierdo D, Catalá M, Jiménez-Macedo A, Mogas T, Paramio M, 2010a: Effect of follicle diameter on oocyte apoptosis, embryo development and chromosomal ploidy in prepubertal goats. *Theriogenology* 74 364-373.

Romaguera R, Morató R, Jiménez-Macedo A, Catalá M, Roura M, Paramio M, Palomo M, Mogas T, Izquierdo D, 2010b: Oocyte secreted factors improve embryo developmental competence of COCs from small follicles in prepubertal goats. *Theriogenology* 74 1050-1059.

Romaguera R, Moll X, Morató R, Roura M, Palomo MJ, Catalá MG, Jiménez-Macedo A, Hammami S, Izquierdo D, Mogas T, Paramio M, 2011: Prepubertal goat oocytes from large follicles result in similar blastocyst production and embryo ploidy than those from adult goats. *Theriogenology* 76 1-11.

Rooke JA, Watt RG, Ashworth CJ, McEvoy TG, 2012: Inclusion of bovine lipoproteins and the vitamin E analogue, Trolox, during in vitro culture of bovine embryos changes both embryo and fetal development. *Journal of Reproduction and Development* 24 309-316.

Rose RC, Bode AM, 1993: Biology of free radical scavengers: an evaluation of ascorbate. *The FASEB Journal* 7 1135-1142.

References

- Sadatsuki M, Tsutsumi O, Yamama R, Muramatsu M, Taketani Y, 1993: Local regulatory effects of activin-A and follistatin on meiotic maturation of rat oocytes. *Biochem.Biophys.Res.Comm* 196 388-395.
- Sakagami N, Umeki H, Nishino O, Uchiyama H, Ichikawa K, Takeshita K, Kaneko E, Akiyama K, Kobayashi S, Tamada H, 2012: Normal calves produced after transfer of embryos cultured in a chemically defined medium supplemented with epidermal growth factor and insulin-like growth factor I following ovum pick up and in vitro fertilization in Japanese black cows. *Journal of Reproduction and Development* 58 140-146.
- Samake S, Amoah E, Mobini S, Gazal O, Gelaye S, 2000: In vitro fertilization of goat oocytes during the non-breeding season. *Small Ruminant Research* 35 49-54.
- Sanbuissho A, Threlfall W, 1990: The influence of serum and gonadotropins on in vitro maturation and fertilization of bovine oocytes. *Theriogenology* 34 341-348.
- Schoevers E, Kidson A, Verheijden J, Bevers M, 2003: Effect of follicle-stimulating hormone on nuclear and cytoplasmic maturation of sow oocytes in vitro. *Theriogenology* 59 2017-2028.
- Selokar N, Saha A, Saini M, Muzaffar M, Chauhan M, Manik R, Palta P, Singla S, 2012: A protocol for differential staining of inner cell mass and trophectoderm of embryos for evaluation of health status. *Curr.Sci* 102 1256-1257.
- Shabankareh HK, Akhondi M, 2012: Effect of glucose levels and MEM vitamins during the first step of IVC in synthetic oviduct fluid medium on in vitro development of ovine zygotes. *Small Ruminant Research* 106 54-58.
- Shehab-El-Deen MA, Leroy JL, Maes D, Van Soom A, 2009: Cryotolerance of bovine blastocysts is affected by oocyte maturation in media containing palmitic or stearic acid. *Reproduction in Domestic Animal* 44(1) 140-142.
- Shirazi A, Sadeghi N, 2007: The effect of ovine oocyte diameter on nuclear maturation. *Small Ruminant Research* 69 103-107.
- Shirazi A, Shams-Esfandabadi N, Ahmadi E, Heidari B, 2010: Effects of growth hormone on nuclear maturation of ovine oocytes and subsequent embryo development. *Reproduction in Domestic Animals* 45 530-536.
- Shirazi A, Ardali MA, Ahmadi E, Nazari H, Mamuee M, Heidari B, 2012: The effect of macromolecule source and type of media during in vitro maturation of sheep oocytes on subsequent embryo development. *Journal of Reproduction and Infertility* 13 13-19.
- Sidis Y, Fujiwara T, Leykin L, Isaacson K, Toth T, Schneyer A, 1998: Characterization of inhibin/activin subunit, activin receptor, and follistatin messenger ribonucleic acid in human and mouse oocytes: Evidence for activin's paracrine signaling from granulosa cells to oocytes. *Biology of Reproduction* 59 807-812.

References

- Silva C, Knight P, 1998: Modulatory actions of activin-A and follistatin on the developmental competence of in vitro-matured bovine oocytes. *Biology of Reproduction* 58 558-565.
- Silva J, van den Hurk R, van Tol H, Roelen B, Figueiredo J, 2004: Gene expression and protein localisation for activin-A, follistatin and activin receptors in goat ovaries. *Journal of Endocrinology* 183 405-415.
- Silva J, Tharasanit T, Taverne M, Van der Weijden G, Santos R, Figueiredo J, Van den Hurk R, 2006: The activin-follistatin system and in vitro early follicle development in goats. *Journal of Endocrinology* 189 113-125.
- Singh B, Barbe G, Armstrong D, 1993: Factors influencing resumption of meiotic maturation and cumulus expansion of porcine oocyte-cumulus cell complexes in-vitro. *Molecular Reproduction and Development* 36 113-119.
- Sirard MA, 2001: Resumption of meiosis: Mechanism involved in meiotic progression and its relation with developmental competence. *Theriogenology* 55 1241-1254.
- Sirard MA, Richard F, Blondin P, Robert C, 2006: Contribution of the oocyte to embryo quality. *Theriogenology* 65 126-136.
- Sirotkin A, 1992: Involvement of steroid-hormones in bovine oocytes maturation in vitro. *The Journal of Steroid Biochemistry and Molecular Biology* 41 855-858.
- Slavik T, Libik M, Wierzchos E, Fulka J, 2005: An attempt to reduce polyspermic penetration in lamb oocytes. *Folia Biologica* 51 34-39.
- Sorensen R, Wassarman P, 1976: Relationship between growth and meiotic maturation of mouse oocyte. *Developmental Biology* 50 531-536.
- Spicer L, Echtenkamp S, 1995: The ovarian insulin and insulin-like growth-factor system with an emphasis on domestic-animals. *Domestic Animal Endocrinology* 12 223-245.
- Stock A, Woodruff T, Smith L, 1997: Effects of inhibin A and activin A during in vitro maturation of bovine oocytes in hormone- and serum-free medium. *Biology of Reproduction* 56 1559-1564.
- Tajik P, Esfandabadi N, 2003: In vitro maturation of caprine oocytes in different culture media. *Small Ruminant Research* 47 155-158.
- Takahashi Y, First NL, 1992: In vitro development of bovine one cell embryos influence of glucose, lactate, pyruvate, amino acids and vitamins. *Theriogenology* 37 963-978.
- Takahashi M, Nagai T, Okamura N, Takahashi H, Okano A, 2002: Promoting effect of beta-mercaptoethanol on in vitro development under oxidative stress and cystine uptake of bovine embryos. *Biology of Reproduction* 66 562-567.

References

- Tao Y, Zhou B, Xia G, Wang F, Wu Z, Fu M, 2004: Exposure to L-ascorbic acid or alpha-tocopherol facilitates the development of porcine denuded oocytes from metaphase I to metaphase II and prevents cumulus cells from fragmentation. *Reproduction in Domestic Animals* 39 52-57.
- Tao Y, Chen H, Tian NN, Huo DT, Li G, Zhang YH, Liu Y, Fang FG, Ding JP, Zhang XR, 2010: Effects of L-ascorbic acid, alpha-tocopherol and co-culture on in vitro developmental potential of porcine cumulus cells free oocytes. *Reproduction in Domestic Animal* 45 19-25.
- Tareq KM, Akter QS, Khandoker MA, Tsujii H, 2012: Selenium and vitamin E improve the in vitro maturation, fertilization and culture to blastocyst of porcine oocytes. *Journal of Reproduction and Development* 58 621-628.
- Tatemoto H, Sakurai N, Muto N, 2000: Protection of porcine oocytes against apoptotic cell death caused by oxidative stress during in vitro maturation: Role of cumulus cells. *Biology of Reproduction* 63 805-810.
- Tatemoto H, Ootaki K, Shigeta K, Muto N, 2001: Enhancement of developmental competence after in vitro fertilization of porcine oocytes by treatment with ascorbic acid 2-O-alpha-glucoside during in vitro maturation. *Biology of Reproduction* 65 1800-1806.
- Tatemoto H, Muto N, Sunagawa I, Shinjo A, Nakada T, 2004: Protection of porcine oocytes against cell damage caused by oxidative stress during in vitro maturation: role of superoxide dismutase activity in porcine follicular fluid. *Biology of Reproduction* 71 1150-1157.
- Tena-Sempere M, 2008: Ghrelin as a pleiotrophic modulator of gonadal function and reproduction. *Nature Reviews Endocrinology* 4 666-674.
- Teotia A, Sharma G, Majumdar A, 2001: Fertilization and development of caprine oocytes matured over granulosa cell monolayers. *Small Ruminant Research* 40 165-177.
- Tervit HR, Whittingham DG, Rowson LE, 1972: Successful culture in vitro of sheep and cattle ova. *Journal of Reproduction and Fertility* 30 493-497.
- Tervit HR, Rowson LE, 1974: Birth of lambs after culture of sheep ova in vitro for up to 6 days. *Journal of Reproduction and Fertility* 38 177-179.
- Thibault C, Gerard M, 1973: Cytoplasmic and nuclear maturation of rabbit oocytes in vitro. *Annales De Biologie Animale Biochimie Biophysique* 13 145-156.
- Thibault C, Gerard M, Menezo Y, 1975: Preovulatory and ovulatory mechanisms in oocyte maturation. *Journals of Reproduction and Fertility* 45 605-610.
- Thomas F, Armstrong D, Telfer E, 2003: Activin promotes oocyte development in ovine preantral follicles in vitro. *Reproductive Biology and Endocrinology* 1 76.

References

- Thouas G, Korfiatis N, French A, Jones G, Trounson A, 2001: Simplified technique for differential staining of inner cell mass and trophectoderm cells of mouse and bovine blastocysts. *Reproductive BioMedicine Online* 3 25-29.
- Tibary A, Anouassi A, Khatir H, 2005: Update on reproductive biotechnologies in small ruminants and camelids. *Theriogenology* 64 618-638.
- Tisdall D, Hudson N, Smith P, McNatt K, 1994: Localization of Ovine Follistatin and Alpha-Inhibin and Beta(a)-Inhibin Messenger-Rna in the Sheep Ovary during the Estrous-Cycle. *Journal of Molecular Endocrinology* 12 181-193.
- Tremoleda J, Tharasanit T, Van Tol H, Stout T, Colenbrander B, Bevers M, 2003: Effects of follicular cells and FSH on the resumption of meiosis in equine oocytes matured in vitro. *Reproduction* 125 565-577.
- Trigal B, Gómez E, Diez C, Caamano J, Martin D, Carrocera S, Munoz M, 2011: In vitro development of bovine embryos cultured with activin A. *Theriogenology* 75 584-588.
- Urdaneta A, Jimenez A, Paramio M, Izquierdo D, 2004: Cysteamine, glutathione and ionomycin treatments improve in vitro fertilization of prepubertal goat oocytes. *Zygote* 12 277-284.
- Van den Hurk R, Van de Pavert S, 2001: Localization of an activin/activin receptor system in the porcine ovary. *Molecular Reproduction and Development* 60 463-471.
- Van der Valk J, Brunner D, De Smet K, Svenningsen AF, Honegger P, Knudsen L, Lindl T, Noraberg J, Price A, Scarino ML, Gstraunthaler G, 2010: Optimization of chemically defined cell culture media - Replacing fetal bovine serum in mammalian in vitro methods. *Toxicology in Vitro* 24 1053-1063.
- Van Soom A, Vanvlaenderen I, Mahmoudzadeh A, Deluyker H, Dekruif A, 1992: Compaction rate of in vitro fertilized bovine embryos related to the interval from insemination to 1st cleavage. *Theriogenology* 38 905-919.
- Van Soom A, Boerjan M, Bols P, Vanroose G, Lein A, Coryn M, deKruif A, 1997: Timing of compaction and inner cell allocation in bovine embryos produced in vivo after superovulation. *Biology of Reproduction* 57 1041-1049.
- Van Soom A, Mateusen B, Leroy J, De Kruif A, 2003: Assessment of mammalian embryo quality: what can we learn from embryo morphology? *Reproductive Biomedicine Online* 7 664-670.
- Van Tol H, Deloos F, Vanderstichele H, Bevers M, 1994: Bovine activin A does not affect the in vitro maturation of bovine oocytes. *Theriogenology* 41 673-679.
- Velilla E, Izquierdo D, Rodriguez-Gonzalez E, Lopez-Bejar M, Vidal F, Paramio M, 2004: Distribution of prepubertal and adult goat oocyte cortical granules during meiotic

References

maturation and fertilisation: Ultrastructural and cytochemical study. *Molecular Reproduction and Development* 68 507-514.

Velilla E, Rodriguez-Gonzalez E, Vidal F, Paramio M, 2005: Microtubule and microfilament organization in immature, in vitro matured and in vitro fertilized prepubertal goat oocytes. *Zygote* 13 155-165.

Velilla E, Rodriguez-Gonzalez E, Vidal F, Izquierdo D, Paramio M, 2006: Mitochondrial organization in prepubertal goat oocytes during in vitro maturation and fertilization. *Molecular Reproduction and Development* 73 617-626.

Villamediana P, Vidal F, Paramio M, 2001: Cytogenetic analysis of caprine 2-to 4-cell embryos produced in vitro. *Zygote* 9 193-199.

Wang B, Baldassarre H, Tao T, Gauthier M, Neveu N, Zhou J, Leduc M, Duguay F, Bilodeau A, Lazaris A, Keefer C, Karatzas C, 2002a: Transgenic goats produced by DNA pronuclear microinjection of in vitro derived zygotes. *Molecular Reproduction and Development* 63 437-443.

Wang F, Tian X, Zhang L, Tan D, Reiter RJ, Liu G. 2013a: Melatonin promotes the in vitro development of pronuclear embryos and increases the efficiency of blastocyst implantation in murine. *Journal of Pineal Research* 55 267-274.

Wang X, Falcone T, Attaran M, Goldberg JM, Agarwal A, Sharma RK, 2002b: Vitamin C and vitamin E supplementation reduce oxidative stress-induced embryo toxicity and improve the blastocyst development rate. *Fertility and Sterility* 78 1272-1277.

Wang Z, Xu Z, Yu S, 2007: Effects of oocyte collection techniques and maturation media on in vitro maturation and subsequent embryo development in Boer goat. *Czech Journal of Animal Science* 52 21-25.

Wang Z, Lin P, Yu S, 2013b: Effects of ghrelin on developmental competence and gene expression of in vitro fertilized ovine embryos. *Theriogenology* 79 695-701.

Wani AR, Khan MZ, Sofi KA, Lone FA, Malik AA, Bhat FA, 2012: Effect of cysteamine and epidermal growth factor (EGF) supplementation in maturation medium on in vitro maturation, fertilization and culturing of embryos in sheep Small Ruminant Research 106 160-164.

Watson AJ, 1992: The cell biology of blastocyst development. *Molecular Reproduction and Development* 33 492-504.

Watson AJ, 2006: Oocyte cytoplasmic maturation: A key mediator of both oocyte and embryo developmental competence. *Journal of Animal Science* 84 1.

Wongsrikeao P, Nagai T, Agung B, Taniguchi M, Kunishi M, Suto S, Otoi T, 2007: Improvement of transgenic cloning efficiencies by culturing recipient oocytes and donor

References

cells with antioxidant vitamins in cattle. *Molecular Reproduction and Development* 74 694-702.

Wu D, Cheung QCK, Wen L, Li J, 2006: A growth-maturation system that enhances the meiotic and developmental competence of porcine oocytes isolated from small follicles. *Biology of Reproduction* 75 547-554.

Wu J, Nayudu PL, Kiesel PS, Michelmann HW, 2000: Luteinizing hormone has a stage-limited effect on preantral follicle development in vitro. *Biology of Reproduction* 63 320-327.

Yoshida M, Ishizaki Y, Kawagishi H, Bamba K, Kojima Y, 1992: Effects of pig follicular fluid on maturation of pig oocytes in vitro and on their subsequent fertilizing and developmental capacity in vitro. *Journal of Reproduction and Fertility* 95 481-488.

Yoshioka K, Kamomae H, 1996: Recombinant human activin-A stimulates development of bovine one-cell embryos matured and fertilized in vitro. *Molecular Reproduction and Development* 45 151-156.

Yoshioka K, Suzuki C, Iwamura S, 1998a: Activin A and follistatin regulate developmental competence of in vitro-produced bovine embryos. *Biology of Reproduction* 59 1017-1022.

Yoshioka K, Takata M, Taniguchi T, Yamanaka H, Sekikawa K, 1998b: Differential expression of activin subunits, activin receptors and follistatin genes in bovine oocytes and preimplantation embryos. *Reproduction Fertility and Development* 10 293-298.

Younis A, Zuelke K, Happer K, Oliveira M, Brackett B, 1991: In vitro fertilization of goat oocytes. *Biology of Reproduction* 44 1177-1182.

Zhang X, Armstrong D, 1990: Presence of amino-acids and insulin in a chemically defined medium improves development of 8-cell rat embryos in vitro and subsequent implantation in vivo. *Biology of Reproduction* 42 662-668.

Zhao J, Taverne M, van der Weijden G, Bevers M, van den Hurk R, 2001: Effect of activin A on in vitro development of rat preantral follicles and localization of activin A and activin receptor II. *Biology of Reproduction* 65 967-977.

Zhou P, Wu YG, Li Q, Lan GC, Wang G, Gao D, Tan JH, 2008: The interactions between cysteamine, cystine and cumulus cells increase the intracellular glutathione level and developmental capacity of goat cumulus-denuded oocytes. *Reproduction*. 135 605-611.

Agradecimientos

Sin lugar a duda, esta tesis no podría haberse realizado sin las aportaciones de un conjunto de personas e instituciones a quien quiero manifestar mis agradecimientos, entre ellas:

A mis directoras de tesis, Dra. Dolors Izquierdo i Tugas y Dra. Roser Morató Molet por sus consejos, seguimiento, supervisión continua a lo largo de estos años y sobre todo las revisiones del presente trabajo. Siempre con un gran criterio científico, pero también por todo el apoyo recibido y sus cualidades humanas. A ambas les agradezco sinceramente, no sólo sus aportaciones, sino también su paciencia.

A la Dra María Teresa Paramio, por aceptarme en su grupo de investigación y por el afecto demostrado durante todos estos años, este trabajo no hubiese sido posible sin su apoyo durante toda la investigación.

Agradezco de manera muy especial a mis compañeras del laboratorio María Catalá, Montse Roura y Roser Romaguera por su colaboración en los trabajos de esta tesis, por estar siempre dispuestas a solucionar cualquier problema que me pudiera surgir, por todas las horas que compartimos juntas, sobre todo por la amistad que me brindaron y los ánimos recibidos.

A la Dra María Rodríguez Prado por haber compartido conmigo sus conocimientos estadísticos, su colaboración en el análisis de los resultados y sobre todo su amistad.

A todas ellas, por haberme permitido aprender tantas cosas, por su ejemplo de profesionalidad y por haber confiado en mí, por todo muchas gracias!!

Mi agradecimiento a todo el personal del departamento de ciència animal i dels aliments: investigadores, profesores, también a todo el personal administrativo y técnico, tanto del departamento como de la granja experimental de la facultat, y en especial a Júlia, Carmen, Blas y Cristóbal. Todos ellos siempre han puesto a mi disposición los medios necesarios para llevar a cabo cualquier asunto y por el buen trato que me han dispensado durante el transcurso de esta tesis.

Quisiera dar las gracias a los amigos y compañeros de la facultat que han estado en cada uno de los instantes vividos en esta etapa: Dra Pipi, Furius, María R, Roser R, Cavrini, Abi, Rosa, Núria A, Ghizlène, Cristina, Manel, Leidy, Ridha, Adel, Arturo y Carlos. Con ellos he compartido momentos inolvidables, diversión, viajes, conversaciones, cumpleaños, platos típicos de nuestros países, fiestas (como la boda sin sofá), risas, libro de excusas,..... Sin olvidarme de los amigos que hice en Barcelona, Fedwa, Tugba, Núria F, Roger, Joan C, Miguel, Najet, Mahrez... A todos gracias por haberme acompañado, por hacerme reír, por brindarme su ayuda y amistad desde el primer momento y siempre.

Al personal del matadero Castellbisbal por facilitarnos la obtención de los ovarios, también a Xinxeta (DEP) y los que hicieron posible todos estos experimentos: Panxona y los machos cabríos de la granja.

Aknowledgements

A la Agencia Española de Cooperación Internacional para el Desarrollo, por la beca entregada durante gran parte del desarrollo de la presente investigación.

¡¡Gracias a todos!!

Mes remerciements les plus sincères vont également à ma famille: ma mère, mon frère et mes sœurs qui m'ont toujours soutenus et encouragés sans faille, pour leur générosité y compréhension. Qu'ils trouvent dans ce travail le résultat de leur confiance et bienveillance!!!

À mes chers petits neveux et nièces, aucune dédicace ne saurait exprimer tout l'amour que j'ai pour vous, votre joie et votre gaieté me comblent de bonheur, que Dieu, vous protège et vous garde!! De même je remercie ma grand-mère, mes oncles, tantes, cousins et toute ma famille.

J'adresse également mes remerciements à tous mes professeurs en Tunisie, spécialement Dr. Fethi Mensi, pour ses précieux conseils au cours de ces années.

Ces remerciements ne seraient pas complets sans une pensée pour mes ami(e)s de longue date, Hajer, Wafa, Nadhem, Mohamed, Chokri.... qui n'ont jamais cessé de me témoigner leur sincère amitié, confiance et encouragements.

Un immense merci à ma chère et meilleure amie Esmâ, qui a su m'encourager et me soutenir tout au long de ces années, merci de m'avoir montré que l'on peut toujours rire face aux difficultés, de m'avoir soutenue dans les moments où j'avais vraiment besoin de bons conseils.

Mes derniers remerciements vont à la personne qui me manque terriblement aujourd'hui mon très cher et adoré grand-père «Ahmed», pour son soutien, sa tendresse, son amour, sa générosité.....et la compréhension dont il a toujours fait preuve à notre égard!!!

Merci à tous!!