



Control biológico de moscas blancas en cultivo de tomate: interacciones entre sus enemigos naturales

Rafael Moreno Ripoll

ADVERTIMENT. La consulta d'aquesta tesi queda condicionada a l'acceptació de les següents condicions d'ús: La difusió d'aquesta tesi per mitjà del servei TDX (www.tdx.cat) ha estat autoritzada pels titulars dels drets de propietat intel·lectual únicament per a usos privats emmarcats en activitats d'investigació i docència. No s'autoritza la seva reproducció amb finalitats de lucre ni la seva difusió i posada a disposició des d'un lloc aliè al servei TDX. No s'autoritza la presentació del seu contingut en una finestra o marc aliè a TDX (framing). Aquesta reserva de drets afecta tant al resum de presentació de la tesi com als seus continguts. En la utilització o cita de parts de la tesi és obligat indicar el nom de la persona autora.

ADVERTENCIA. La consulta de esta tesis queda condicionada a la aceptación de las siguientes condiciones de uso: La difusión de esta tesis por medio del servicio TDR (www.tdx.cat) ha sido autorizada por los titulares de los derechos de propiedad intelectual únicamente para usos privados enmarcados en actividades de investigación y docencia. No se autoriza su reproducción con finalidades de lucro ni su difusión y puesta a disposición desde un sitio ajeno al servicio TDR. No se autoriza la presentación de su contenido en una ventana o marco ajeno a TDR (framing). Esta reserva de derechos afecta tanto al resumen de presentación de la tesis como a sus contenidos. En la utilización o cita de partes de la tesis es obligado indicar el nombre de la persona autora.

WARNING. On having consulted this thesis you're accepting the following use conditions: Spreading this thesis by the TDX (www.tdx.cat) service has been authorized by the titular of the intellectual property rights only for private uses placed in investigation and teaching activities. Reproduction with lucrative aims is not authorized neither its spreading and availability from a site foreign to the TDX service. Introducing its content in a window or frame foreign to the TDX service is not authorized (framing). This rights affect to the presentation summary of the thesis as well as to its contents. In the using or citation of parts of the thesis it's obliged to indicate the name of the author.



DEPARTAMENTO DE BIOLOGÍA ANIMAL

FACULTAD DE BIOLOGÍA
UNIVERSIDAD DE BARCELONA

Programa de doctorado de Biología Animal
Bienio 2005-2007

Control biológico de moscas blancas en cultivo de tomate: interacciones entre sus enemigos naturales

Memoria presentada por **Rafael Moreno Ripoll** para
optar al título de Doctor por la Universidad de Barcelona

Tesis realizada en Entomología, Institut de Recerca i Tecnologia Agroalimentàries
(IRTA, Cabrils)

Directoras

Dra. Rosa Gabarra Ambert
Investigadora de Entomología
IRTA – Centro de Cabrils

Dra. Núria Agustí Abella
Investigadora de Entomología
IRTA – Centro de Cabrils

Tutora

Dra. Marta Goula Goula
Investigadora titular Dto. de Biología
Animal
Facultad de Biología
Universidad de Barcelona

Bibliografia

Agarwala, B.K., y Dixon, A.F.G. (1993) Kin recognition: Egg and larval cannibalism in *Adalia bipunctata* (Coleoptera: Coccinellidae). *European Journal of Entomology* 90, 45-50.

Agarwala, B.K., Yasuda, H. y Kajita, Y. (2003) Effects of conspecific and heterospecific feces and oviposition of two predatory ladybirds: role of fecal cues in predator avoidance. *Journal of Chemical Ecology* 29(2), 357-376.

Agarwala, B.K., Yasuda, H. y Sato, S. (2008) Life history response of a predatory ladybird, *Harmonia axyridis* (Pallas) (Coleoptera: Coccinellidae), to food stress. *Applied Entomology and Zoology* 43(2), 183-189.

Agustí, N. (1998) Biología de *Dicyphus tamaninii* Wagner (Heteroptera: Miridae) i identificació molecular de les preses ingerides. Tesis doctoral, Universidad de Barcelona.

Agustí, N., de Vicente, M.C. y Gabarra, R. (1999) Development of sequence amplified characterized region (SCAR) markers of *Helicoverpa armigera*: a new polymerase chain reaction-based technique for predator gut analysis. *Molecular Ecology* 8, 1467-1474.

Agustí, N., de Vicente, M.C. y Gabarra, R. (2000) Developing SCAR markers to study predation on *Trialeurodes vaporariorum*. *Insect Molecular Biology* 9, 263-268.

Agustí, N., Shayler, S., Harwood, J.D., Vaughan, I.P., Sunderland, K.D. y Symondson, W.O.C. (2003) Collembola as alternative prey sustaining spiders in arable ecosystems: prey detection within predators using molecular markers. *Molecular Ecology* 12, 3467-3475.

Agustí, N., Bourguet, D., Spataro, T., Delos, M., Eychenne, N., Folcher, L y Ardití, R. (2005) Detection, identification and geographical distribution of European corn borer larval parasitoids using molecular markers. *Molecular Ecology* 14, 3267-3274.

Agustí, N. y Gabarra, R (2006) Utilización de métodos moleculares en programas de Control Integrado: aplicación a estudios de depredación y parasitismo. *PHYTOMA España* 180, 46-51.

Agustí, N. y Gabarra, R. (2009a) Puesta a punto de una cría masiva del depredador polífago *Dicyphus tamaninii* Wagner (Heteroptera: Miridae). *Boletín Sanidad Vegetal: Plagas*, 35, 205-218.

Agustí, N. y Gabarra, R. (2009b) Effect of adult age and insect density of *Dicyphus tamaninii* Wagner (Heteroptera: Miridae) on progeny. *Journal of Pest Science* 82(3), 241-246.

Albajes, R. y Alomar, O. (1999) Current and potential use of polyphagous predators. Pp. 265-275. En Albajes, R., Gullino, M.L., van Lenteren, J.C. y Elad, Y. (Eds). *Integrated Pest and Disease Management in Greenhouse Crops*. Kluwer Academic Publishers, Dordrecht, Netherlands.

Albajes, R., Sarasúa, M.J., Avilla, J., Arnó, J. y Gabarra, R. (2003) Integrated pest management in the mediterranean region: the case of Catalonia, Spain. Pp. 341-355. En Maredia, K.M., Dakouo, D. y Mota-Sanchez, D. (Eds). *Integrated Pest Management in the Global Arena*. CABI Publishing, Wallingford.

Alomar, O. (1994) Els mirids depredadors (Heteroptera: Miridae) en el control integrat de plagues en conreus de tomàquet. Tesis doctoral, Universidad de Barcelona.

Alomar, O., Goula, M. y Albajes, R. (2002) Colonisation of tomato fields by predatory mirid bugs (Hemiptera: Heteroptera) in northern Spain. *Agriculture, Ecosystems and Environment* 89, 105-115.

Alomar, O., Riudavets, J. y Castañé, C. (2006) *Macrolophus caliginosus* in the biological control of *Bemisia tabaci* on greenhouse melons. *Biological Control* 36 (2), 154-162.

Amornsak, W., Gordh, G. y Graham, G. (1998) Detecting parasitized eggs with polymerase chain reaction and DNA sequence of *Trichogramma australicum* Girault. *Australian Journal of Entomology*, 37, 174-179.

Arim, M. y Marquet, P.A. (2004) Intraguild predation: a widespread interaction related to species biology. *Ecology Letters* 7, 557-564.

Arnó, J., Matas, M., Martí, M., Ariño, J., Roig, J. y Gabarra, R. (2005) Coexistence between *Trialeurodes vaporariorum* and *Bemisia tabaci* and impact of natural enemies in tomato crops under Mediterranean conditions. *IOBC/WPRS Bulletin* 28, 1-4.

Arnó J., Albajes R. y Gabarra R. (2006a) Within-plant distribution and sampling of single and mixed infestations of *Bemisia tabaci* and *Trialeurodes vaporariorum* (Homoptera: Aleyrodidae) in winter tomato crops. *Journal of Economic Entomology* 99, 331-340.

Arnó, J., Castañé, C., Riudavets, J., Roig, J. y Gabarra, R. (2006b) Characterization of damage to tomato plants produced by the zoophytophagous predator *Nesidiocoris tenuis*. *IOBC/WPRS Bulletin* 29, 249-254.

Arnó, J., Sorribas, R., Prat, M., Matas, M., Pozo, C., Rodríguez, D., Garreta, A., Gómez, A. y Gabarra, R. (2009) *Tuta absoluta*, a new pest in IPM tomatoes in the northeast of Spain. *IOBC/WPRS Bulletin* 49, 203-208.

Arnó, J., Gabarra, R., Liu, T.X., Simmons, A.M. y Gerling, D. (2010a) Natural enemies of *Bemisia tabaci*: predators and parasitoids. Pp. 385-421. En Stansly, P. A., Naranjo, S.E. (Eds). *Bemisia: Bionomics and Management of a Global Pest*. Springer, Dordrecht, the Netherlands.

Arnó J., Castañé C., Riudavets J. y Gabarra R. (2010b) Risk of damage to tomato crops by the generalist zoophytophagous predator *Nesidiocoris tenuis* (Reuter) (Hemiptera: Miridae). *Buletin of Entomological Research* 100, 105-115.

Ashfaq, M., Shah, G.S., Noor, A.R., Ansari, S.P y Mansoor, S. (2010) Report of a parasitic wasp (Hymenoptera: Encyrtidae) parasitizing cotton mealybug (Hemiptera: Pseudococcidae) in Pakistan and use of PCR for estimating parasitism levels. *Biocontrol Science and Technology* 20(6), 625-630.

Avilla J., Albajes R., Alomar O., Castañé C. y Gabarra R. (2004) Biological control of whiteflies in protected vegetable crops. Pp. 171-184. En Parrella, M.P. y Heinz, K.M. (Eds). *Biological Control of Arthropods Pests in Protected culture*. Batavia, Ball Publishing.

Barnadas, I., Gabarra, R. y Albajes, R. (1998) Predatory capacity of two mirid bugs preying on *Bemisia tabaci*. *Entomologia Experimentalis et Applicata* 86, 215-219.

Bográn, C.E., y Heinz, K.M. (2002) Host selection by the heteronomous hyperparasitoid *Encarsia pergandiella*: multiple-choice tests using *Bemisia argentifolii* as primary host. *Entomologia Experimentalis et Applicata* 103,11-21.

Bográn, C.E. y Heinz, K.M. (2006) Time delay and initial population density affect interactions between *Encarsia pergandiella* Howard and *Eretmocerus mundus* Mercet (Hymenoptera: Aphelinidae). *Environmental Entomology* 35(3), 661-669.

Boisclair J., Brueren G.J. y van Lenteren J.C. (1990) Can *Bemisia tabaci* be controlled with *Encarsia formosa*? *IOBC/WPRS Bulletin* 13, 32-35.

Bonato, O., Couton, L. y Fargues, J. (2006) Feeding preference of *Macrolophus caliginosus* (Heteroptera: Miridae) on *Bemisia tabaci* and *Trialeurodes vaporariorum* (Homoptera: Aleyrodidae). *Journal of Economic Entomology* 99, 1143-1151.

Brodeur, J. y Rosenheim, J.A. (2000) Intraguild interactions in aphid parasitoids. *Entomologia Experimentalis et Applicata* 97, 93-108.

Buddle, C.M., Walker, S.E. y Rypstra, A.L. (2003) Cannibalism and density-dependent mortality in the wolf spider *Pardosa milvina* (Araneae: Lycosidae). *Canadian Journal of Zoology* 81, 1293-1297.

Calvo, J., Bolckmans, K., Stansly, P.A. y Urbaneja, A. (2009) Predation by *Nesidiocoris tenuis* on *Bemisia tabaci* and injury to tomato. *Biocontrol* 54, 237-246.

Castañé, C., Alomar, O., Goula, M. y Gabarra, R. (2004) Colonization of tomato greenhouses by the predatory mirid bugs *Macrolophus caliginosus* and *Dicyphus tamaninii*. *Biological Control* 30, 591-597.

Castañé, C., Arnó, J., Beitia, F. y Gabarra, R. (2008) Control biológico de moscas blancas. Pp. 239-253. En Jacas, J.A., Urbaneja, A. (Eds). Control Biológico de Plagas Agrícolas. Phytoma España, S.L.

Castañé, C., Arnó, J., Gabarra, R. y Alomar, O. (2011) Plant damage to vegetable crops by zoophytophagous mirid predators. *Biological Control*. doi: 10.1016/j.biocontrol.2011.03.007.

Caterino M.S., Cho, S. y Sperling, A.H. (2000) The current state of insect molecular systematics: a thriving tower of Babel. *Annual Review of Entomology*, 45, 1-54.

Chacón, J.M., Landis. D.A. y Heimpel, G.E. (2008) Potential for biotic interference of a classical biological control agent of the soybean aphid. *Biological Control* 46, 216-225.

Choh, Y., Masayoshi, U. y Takabayashi, J. (2010) Predation-related odours reduce oviposition in a herbivorous mite. *Experimental and Applied Acarology* 50, 1-8.

Colfer, R.G. y Rosenheim, J.A. (2001) Predation on immature parasitoids and its impact on aphid suppression. *Oecologia* 126, 292-304.

Cottrell, T.E. (2005) Predation and cannibalism of lady beetle eggs by adult lady beetles. *Biological Control* 34,159-164.

Crumrine, P.W. (2010) Size-structured cannibalism between top predators promotes the survival of intermediate predators in an intraguild predation system. *Journal of the North American Benthological Society* 29(2), 636-646.

De Barro, P. J. y Hart P. (2001) Antibiotic curing of parthenogenesis in *Eretmocerus mundus* Mercet (Australian parthenogenteic form) (Hymenoptera, Aphelinidae). *Entomologia Experimentalis et Applicata* 99, 225-230.

De Barro, P.J., Trueman, J.W.H. y Frohlich, D.R. (2005) *Bemisia argentifolii* is a race of *B. tabaci* (Hemiptera: Aleyrodidae): the molecular genetic differentiation of *B. tabaci* populations around the world. *Bulletin of Entomological Research* 95, 193-203.

DeBach, P. (1964) Biological control of insect pest and weeds. Chapman & Hall. LTD. London.

Dixon, A. G. F. (2000) Intraguild Predation. Pp. 190-216. In *Intraguild Predation in Insect Predator-Prey Dynamics: Ladybird Beetles and Biological Control*. Cambridge Press.

Eilenberg, J. Hajek, A. y Lomer, C. (2001) Suggestions for unifying the terminology in biological control. *BioControl* 46, 387-400.

Eitam, A. y Blaustein, L (2004) Oviposition habitat selection by mosquitoes in response to predator (*Notonecta maculata*) density. *Physiological Entomology* 29, 188-191.

FAOSTAT (FOOD AND AGRICULTURE ORGANIZATION OF THE UNITED NATIONS). <http://faostat.fao.org/site/339/default.aspx>.

Faria, M. y Wraight, S.P. (2001) Biological control of *Bemisia tabaci* with fungi. *Crop Protection* 20, 767-778.

Fazal, S. y Xiang, R.S. (2004) Interaction of *Serangium japonicum* (Coleoptera: Coccinellidae), an obligate predator of whitefly with immature stages of *Eretmocerus sp.* (Hymenoptera: Aphelinidae) within whitefly host (Homoptera: Aleyrodidae). *Asian Journal of Plant Science* 3, 243-246.

Félix, S. y Soares, A. O. (2004) Intraguild predation between the aphidophagous ladybird beetles *Harmonia axyridis* and *Coccinella undecimpunctata* (Coleoptera: Coccinellidae): the role of body weight. *European Journal of Entomology* 101, 237-242.

Ferris, G. y Rudolf, W.H.W. (2007) Responses of larval dragonflies to conspecific and heterospecific predator cues. *Ecological Entomology* 32, 283-288.

Finke, D.L. y Denno, R.F. (2003) Intraguild predation relaxes natural enemy impacts on herbivore populations. *Ecological Entomology* 28, 67-73.

Folmer, O., Black, M., Hoeh, W., Lutz, R. y Vrijenhoek, R. (1994) DNA primers for the amplification of mitochondrial cytochrome c oxidase subunit I from diverse metazoan invertebrates. *Molecular Marine Biology and Biotechnology* 3(5), 294-299.

Foltan, F., Sheppard, S., Konvicka, M. y Symondson, W.O.C. (2005) The significance of facultative scavenging in generalist predator nutrition. Detecting decayed prey in the guts of predators using PCR. *Molecular Ecology* 14, 4147-4158.

Fox, L.R. (1975) Cannibalism in natural populations. *Annual Review of Ecology and Systematics* 6, 87-106.

Fréchette, B., Rojo, S., Alomar, O. y Lucas, E. (2007) Intraguild predation between syrphids and mirids: who is the prey? Who is the Predator? *Biocontrol* 52, 175-191.

Gabarra, R., Arnó, J., Alomar, O. y Albajes, R. (1999) Naturally occurring populations of *Encarsia pergandiella* (Hymenoptera: Aphelinidae) in tomato greenhouses. *IOBC/WPRS Bulletin* 22, 85-88.

Gabarra, R. y Besri, M. (1999) Tomatoes. Pp. 420-434. En Albajes, R., Gullino, M.L., van Lenteren, J.C. y Elad, Y. (Eds). *Integrated Pest and Disease Management in Greenhouse Crops*. Kluwer Academic Publishers, Dordrecht.

Gabarra R., Alomar, O., Castañé, C., Goula, M. y Albajes, R. (2004) Movement of greenhouses whitefly and its predators between in- and outside of Mediterranean greenhouses. *Agriculture, Ecosystems Environment*. 102, 341-348.

Gabarra, R., Zapata, R., Castañé, C., Riudavets, J. y Arnó, J. (2006) Releases of *Eretmocerus mundus* and *Macrolophus caliginosus* for controlling *Bemisia tabaci* on spring and autumn greenhouse tomato crops. *IOBC/WPRS Bulletin* 29, 71-76.

Gabarra, R., Arnó, J. y Riudavets, J. (2008) Tomate. Pp. 410-422. En Jacas, J.A., Urbaneja, A. (Eds). *Control Biológico de Plagas Agrícolas*. Phytoma España, S.L.

Gariepy, T.D, Kuhlmann, U., Haye, T., Gillott, C. y Erlandson, M. (2005) A single-step multiplex PCR assay for the detection of European *Peristenus* spp., parasitoids of *Lygus* spp. *Biocontrol Science and Technology* 15(5), 481-495.

Gariepy, T.D., Kuhlmann, U., Gillott, C. y Erlandson, M. (2007) Parasitoids, predators and PCR: the use of diagnostic molecular markers in biological control of Arthropods. *Journal of Applied Entomology* 131, 225-240.

Gariepy, T., Kuhlmann, U., Gillott, C. y Erlandson, M. (2008) A large-scale comparison of conventional and molecular methods for the evaluation of host-parasitoid associations in non-target risk-assessment studies. *Journal of Applied Ecology* 45, 708-715.

Gerling, D. (1990) Natural enemies of whiteflies: predators and parasitoids. Pp. 147-186. En Gerling, D. (Ed). *Whiteflies: their Bionomics, Pest Status and Management*. Wimborne, UK, Intercept Ltd.

Gerling, D. y Fried, R. (2000) Biological studies with *Eretmocerus mundus* Mercet (Hymenoptera: Aphelinidae) in Israel. *IOBC/WPRS Bulletin* 23(1), 117-123.

Gerling, D., Alomar, O. y Arnó, J. (2001) Biological control of *Bemisia tabaci* using predators and parasitoids. *Crop Protection* 20, 779-799.

Gill, R.J. (1990) The morphology of whiteflies. Pp. 1346. En: Gerling P (ed) *Whiteflies: their Bionomics, Pest Status and Management*. Intercept Ltd, Andover.

Goula, M. y Alomar, O. (1994) Míridos (Heteroptera Miridae) de interés en el control integrado de plagas en el tomate. Guía para su identificación. *Boletín de Sanidad Vegetal Plagas* 20, 131-143.

Greenstone, M.H. (2006) Molecular methods for assessing insect parasitism. *Bulletin of Entomological Research* 96, 1-13.

Harper, G.L., King, R.A., Dodd, C.S., Harwood, J.D., Glen, D.M., Bruford, M.W. y Symondson, W.O.C. (2005) Rapid screening of invertebrate predators for multiple prey DNA targets. *Molecular Ecology* 14, 819-827.

Harwood, J.D., Sunderland, K.D. y Symondson, W.O.C. (2004) Prey selection by linyphiid spiders: molecular tracking of the effects of alternative prey on rates of aphid consumption in the field. *Molecular Ecology* 13, 3549-3560.

Harwood, J.D., Desneux, N., Yoo, H.Y.S., Rowley, D.L., Greenstone, M.H., Obrycki, J.J. y O'Neil, R.J. (2007) Tracking the role of alternative prey in soybean aphid predation by *Orius insidiosus*: a molecular approach. *Molecular Ecology* 16, 4390-4400.

Hoddle, M.S., Van Driesche, R.G. y Sanderson, J.P. (1998) Biology and use of the whitefly parasitoid *Encarsia formosa*. *Annual review of entomology* 43, 645-669.

Hoelmer, K.A., Osborne, L.S. y Yokomi, R.K. (1994) Interactions of the whitefly predator *Delphastus pusillus* (Coleoptera: Coccinellidae) with parasitized sweetpotato whitefly (Homoptera: Aleyrodidae). *Environmental Entomology* 23, 136-139.

Horowitz A.R., Kontsedalov, S., Khasdan, V. y Ishaaya, I. (2005) Biotypes B and Q of *Bemisia tabaci* and their relevance to neonicotinoid and pyriproxyfen resistance. *Archives of Insect Biochemical and Physiology* 58, 216-25.

Hunter, M.S. (1989) Suitability of stages of female *Encarsia pergandiella* [Hymenoptera: Aphelinidae] for development of conspecific male hyperparasites. *Entomophaga* 34, 265-74.

Hurtado, M., Agustí, N. y Sabater-Muñoz, B. (2008) Aplicación de técnicas moleculares al control biológico de plagas. Pp. 453-465. En Jacas, J.A., Urbaneja, A. (Eds). Control Biológico de Plagas Agrícolas. Phytoma España, S.L.

Ingels, B. y De Clercq, P. (2011) Effect of size, extraguild prey and habitat complexity on intraguild interactions: a case study with the invasive ladybird *Harmonia axyridis* and the hoverfly *Episyrphus balteatus*. *BioControl*. doi 10.1007/s10526-011-9363-9.

Jacas, J. y Urbaneja, A. (2008a) Origen de las plagas e historia del control biológico. Pp. 3-14. En Jacas, J.A., Urbaneja, A. (Eds). Control Biológico de Plagas Agrícolas. Phytoma España, S.L.

Jacas, J. y Urbaneja, A. (2008b) Situación actual y retos del control Biológico de Plagas. Pp, 485-494. En Jacas, J.A., Urbaneja, A. (Eds). Control Biológico de Plagas Agrícolas. Phytoma España, S.L.

Jakobsen, L., Enkegaard, A. y Brødsgaard, H.F. (2004) Interactions between two polyphagous predators, *Orius majusculus* (Hemiptera: Anthocoridae) and *Macrolophus caliginosus* (Heteroptera: Miridae). *Biocontrol Science and Technology* 14, 17-24.

Jarman, S.N. (2004) Amplicon: software for designing PCR primers on aligned DNA sequences. *Bioinformatics* 20, 1644-1645.

Jervis, M. y Kidd, N. (1986) Host-feeding strategies in hymenopteran parasitoids. *Biological Reviews* 61, 395-343.

Johansson, F. (1993) Intraguild predation and cannibalism in odonate larvae: effects of foraging behaviour and zooplankton availability. *Oikos* 66, 80-87.

Juen, A. y Traugott, M. (2005) Detecting predation and scavenging by DNA gut-content analysis a case study using a soil insect predator-prey system. *Oecologia* 142, 344-352.

Juen, A. y Traugott, M. (2007) Revealing species-specific trophic links in soil food webs: molecular identification of scarab predators. *Molecular Ecology* 16, 1545-1557.

Kajita, Y. y Evans, E.W. (2009) Ovarian dynamics and oosorption in two species of predatory lady beetles (Coleoptera: Coccinellidae). *Physiological Entomology* 34, 185-194.

King, R.A., Read, D.S., Traugott, M. y Symondson, W.O.C. (2008) Molecular analysis of predation: a review of best practice for DNA-based approaches. *Molecular Ecology* 17, 947-963.

King, R.A., Vaughan, I.P., Bell, J.R., Bohan, D.A. y Symondson, W.O.C. (2010) Prey choice by carabid beetles feeding on an earthworm community analysed using species- and lineage-specific PCR primers. *Molecular Ecology* 19, 1721-1732.

King, R.A., Moreno-Ripoll, R., Agustí, N., Shayler, S.P., Bell, J.R., Bohan, D.A. y Symondson, W.O.C. (2011) Multiplex reactions for the molecular detection of predation on pest and non-pest invertebrates in agroecosystems. *Molecular Ecology Resources* 11, 370-373.

Kuusk, A-K, y Agustí, N. (2008) Group-specific primers for DNA-based detection of springtails (Hexapoda: Collembola) within predator gut contents. *Molecular Ecology Resources* 8, 678-681.

Kuusk, A-K., Cassel-Lundhagen, A., Kvarnheden, A. y Ekbom, B. (2008) Tracking aphid predation by lycosid spiders in spring-sown cereals using PCR-based gut-content analysis. *Basic and Applied Ecology* 9, 718-725.

Laycock, A., Camm, E., Van Laerhoven, S., y Gillespie, D. (2006) Cannibalism in a zoophytophagous omnivore is mediated by prey availability and plant substrate. *Journal of Insect Behavior* 19, 219-229.

Lima, S. L. (1998) Nonlethal effects in the ecology of predator-prey interactions. What are the ecological effects of anti-predator decision-making? *BioScience* 48, 25-34.

Lipa, J.J. y Smits, P.H. (1999) Microbial control of pests in greenhouses. Pp. 295-309. En Albajes, R., Gullino, M.L., van Lenteren, J.C. y Elad, Y. (Eds). *Integrated Pest and Disease Management in Greenhouse Crops*, Kluwer Academic Publishers, Dordrecht.

Liu, T.X. y Stansly, P.A. (1996) Oviposition, development, and survivorship of *Encarsia pergandiella* (Hymenoptera: Aphelinidae) in four instars of *Bemisia argentifolii* (Homoptera: Aleyrodidae). *Annals of the Entomological Society of America* 89, 96-102.

Losey, J.E. y Denno, R.F. (1998) Positive predator-predator interactions: enhanced predation rates and synergistic suppression of aphid populations. *Ecology* 79, 2143-2152.

Lucas, E., Coderre, D. y Brodeur, J. (1997) Instar-specific defense of *Coleomegilla maculata lengi* (Coccinellidae): influence on attack success of the intraguild predator *Chrysoperla rufilabris* (Chrysopidae). *Entomophaga* 42, 3-12.

Lucas, E., Coderre, D. y Brodeur, J. (1998) Intraguild predation among aphid predators: characterization and influence of extraguild prey density. *Ecology* 79, 1084-1092.

Lucas, E. y Alomar, O. (2002) Impact of the presence of *Dicyphus tamaninii* Wagner (Heteroptera: Miridae) on whitefly (Homoptera: Aleyrodidae) predation by *Macrolophus caliginosus* (Wagner) (Heteroptera: Miridae). *Biological Control* 25, 123-128.

Lucas, E., Fréchette, B. y Alomar, O. (2009) Resource quality, resource availability, and intraguild predation among omnivorous mirids. *Biocontrol Science and Technology* 19(6), 555-572.

Magalhães, S., Tudorache, C., Montserrat, M., van Maanen, R., Sabelis, M.W. y Janssen, A. (2004) Diet of intraguild predators affects anti predator behavior in intraguild prey. *Behavioral Ecology* 16, 364-370.

Malo, S. (2009) Control biològic de *Bemisia tabaci* (Homoptera: Aleyrodidae): *Eretmocerus mundus* (Hymenoptera: Aphelinidae), *Macrolophus pygmaeus* (Heteroptera: Miridae) i la seva interacció. Tesis doctoral, Universidad de Lleida.

Malo, S., Riudavets, J. y Gabarra, R. (2009) Insecticide tolerance of *Eretmocerus mundus* populations collected in commercial vegetable greenhouses. *IOBC/WPRS Bulletin* 49, 107112.

MAPA. Anuario de estadística, 2009. Ministerio del Medio Ambiente y Medio Rural y Marino. <http://www.mapa.es/es/estadistica/pags/anuario/2009/indice.asp>.

Martinez-Cascales, J.I., Cenis, J.L., Cassis, G. y Sanchez, J.A. (2006) Species identity of *Macrolophus melanotoma* (Costa 1853) and *Macrolophus pygmaeus* (Rambur 1839) (Insecta: Heteroptera: Miridae) based on morphological and molecular data and bionomic implications. *Insect Systematics & Evolution* 37, 385-404.

Martinou A.F., Milonas, P.G. y Wright, D.J. (2009) Patch residence decisions made by *Aphidius colemani* in the presence of a facultative predator. *Biological control* 49(3), 234-238.

McPherson, M. J. y Møller, S. G. (2000) PCR. Springer, New York.

Montserrat, M., Albajes, R. y Castañé, C. (2000a) Comparative behaviour of three predators used in biological control in greenhouse crops. *IOBC/WPRS Bulletin* 23, 267-272.

Montserrat, M., Albajes, R. y Castañé, C. (2000b) Functional response of four heteropteran predators preying on greenhouse whitefly (Homoptera: Aleyrodidae) and western flower thrips (Thysanoptera: Thripidae). *Environmental Entomology* 29, 1075-1082.

Montserrat, M., Bas, C., Magalhães, S., Sabelis, M.W., de Roos, A.M. y Janssen, A. (2007) Predators induce egg retention in prey. *Oecologia* 150, 699-705.

Moreno-Ripoll, R., Gabarra, R. y Agustí, N. (2009) Tracking predation of *Macrolophus pygmaeus* and *Nesidiocoris tenuis* (Heteroptera: Miridae) in tomato protected crops by molecular methods. *IOBC/WPRS Bulletin* 49, 307-312.

Moretti, R. y Calvitti, M (2008) Intrinsic competition between the parasitoids *Eretmocerus mundus* and *Encarsia Formosa* in *Bemisia tabaci*. *Entomologia Experimentalis et Applicata* 129, 44-53.

Moriones, E. y Luis-Arteaga, M. (1999) Viral diseases. Pp. 16-33. En Albajes, R., Gullino, M.L., van Lenteren, J.C. y Elad, Y. (Eds). *Integrated Pest and Disease Management in Greenhouse Crops*, Kluwer Academic Publishers, Dordrecht.

Moriones, E. y Navas-Castillo, J. (2010) Tomato yellow leaf curl disease epidemics. Pp. 259-282. En Stansly, P.A. y Naranjo, S.E. (Eds). *Bemisia: Bionomics and Management of a Global Pest*. Dordrecht, the Netherlands, Springer.

Mound, L.A. y Halsey, S.H., (1978) Whiteflies of the world. Pp. 340. British Museum of Natural History and Wiley, New York, NY..

Moya, A., Guirao, P., Cifuentes, D., Beitia, F. y Cenis, J.L. (2001) Genetic diversity of Iberian populations of *Bemisia tabaci* (Hemiptera: Aleyrodidae) based on random amplified polymorphic DNA-polymerase chain reaction. *Molecular Ecology* 10, 891-897.

Naranjo, S.E. y Gibson, R.L. (1996) Phytophagy in predaceous Heteroptera: effects on life history and population dynamics. Pp. 57-93. En Alomar O. y Wiedenmann R.N. (Eds). Zoophytophagous Heteroptera: Implications for Life History and Integrated Pest Management. Thomas Say Publications in Entomology, Lanham, MD.

Naranjo, S.E., Cañas, L. y Ellsworth, P.C. (2004) Mortalidad de *Bemisia tabaci* en un sistema de cultivos múltiples. *Horticultura Internacional* 43, 14-21.

Naranjo, S.E. (2007) Intraguild predation on *Eretmocerus* sp nr. *emiratus*, a parasitoid of *Bemisia tabaci*, by three generalist predators with implications for estimating the level and impact of parasitism. *Biocontrol Science and Technology* 17, 605-622.

Nóia, M., Borges, I. y Soares, A.O. (2008) Intraguild predation between the aphidophagous ladybird beetles *Harmonia axyridis* and *Coccinella undecimpunctata* (Coleoptera: Coccinellidae): the role of intra and extraguild prey densities. *Biological Control* 46, 140-146.

Nomikou, M., Janssen, A. y Sabelis, M.W. (2003) Herbivore host plant selection: whitefly learns to avoid host plants that harbour predators of her offspring. *Oecologia* 136, 484-488.

Okuyama, T. (2002) The role of antipredator behavior in an experimental community of jumping spiders with intraguild predation. *Population Ecology* 44, 121-125.

Oliveira, M.R.V., Henneberry, T.J. y Anderson, P. (2001) History, current status, and collaborative research projects for *Bemisia tabaci*. *Crop Protection* 20, 709-725.

Palumbo, J.C., Horowitz, A.R. y Prabhaker, N. (2001) Insecticidal control and resistance management for *Bemisia tabaci*. *Crop Protection* 20, 739-765.

Perdikis, D.C., Margaritopoulos, J.T., Stamatis, C., Mamuris, Z., Lykouressis, D. P., Tsitsipis, J. A., y Pekas, A. (2003) Discrimination of the closely related biocontrol agents *Macrolophus melanotoma* (Hemiptera: Miridae) and *M. pygmaeus* using mitochondrial DNA analysis. *Bulletin of Entomological Research* 93, 507-514.

Perdikis, D., Lucas, E., Garantonakis, N., Giatropoulos, A., Kitsis, P., Maselou, D., Panagakis, S., Paraskevopoulos, A., Lykouressis, D. y Fantinou, A. (2009) Intraguild predation between *Macrolophus pygmaeus* and *Nesidiocoris tenuis*. *IOBC/WPRS Bulletin* 49, 301-305.

Perring, T.M., Farrar, C.A. y Royalty, R. N. (1987) Intraplant distribution and sampling of spider mites (Acari: Tetranychidae) on cantaloupe. *Journal of Economic Entomology* 80, 96-101.

Perring, T.M. (2001) The *Bemisia tabaci* species complex. *Crop Protection* 20, 725-737.

Polis, G.A. (1981) The evolution and dynamics of intraspecific predation. *Annual Review of Ecology and Systematics* 12, 225-251.

Polis, G.A., Myers, C.A., y Holt, R.D. (1989) The ecology and evolution of intraguild predation: potential competitors that eat each other. *Annual Review of Ecology and Systematics* 20, 297-330.

Polis, G. A. y Holt, R. D. (1992) Intraguild predation: the dynamics of complex trophic interactions. *Trends in Ecology & Evolution* 7, 151-154.

Polis, G.A., Sears, A.L.W., Huxel, G.R., Strong, D.R. y Maron, J. (2000) When is a trophic cascade a trophic cascade? *Trends in Ecology & Evolution* 15, 473-475.

Pollard, D.G. (1955) Feeding habits of the cotton whitefly, *Bemisia tabaci* Genn. (Homoptera: Aleyrodidae). *Annals of Applied Biology* 43, 664-671.

Pumariño L., Alomar, O. y Agustí, N. (2011) Development of specific ITS markers for plant DNA identification within herbivorous insects. *Bulletin of Entomological Research* 101, 271-276.

Raman, K. y Sanjayan, K.P. (1984) Histology and histopathology of the feeding lesions by *Cyrtopeltis tenuis* Reut. (Hemiptera: Miridae) on *Lycopersicon esculentum* Mill. (Solanaceae). *Proceedings Indian Academy of Science (Animal Science)* 93(6), 543-547.

Rose, M., Zolnerowich, G. y Hunter, M.S. (1995) Systematics, *Eretmocerus*, and biological control,. Pp. 477-497. En Gerling, D. y Mayer, R.T. (Eds). *Bemisia: Taxonomy, Biology, Damage, Control and Management*. Intercept, Andover, UK.

Rosenheim, J.A., Kaya, H.K., Ehler, L.E., Marois, J.J. y Jaffee, B.A. (1995) Intraguild predation among biological-control agents: theory and evidence. *Biological Control* 5, 303-335.

Rosenheim, J.A., Heimpel, G.E. y Mangel, M. (2000) Egg maturation, egg resorption and the costliness of transient egg limitation in insects. *Proceedings of the Royal Society B* 267, 1565-1573.

Rudolf, V.H.W. (2007) The interaction of cannibalism and omnivory: consequences for community dynamics. *Ecology* 88(11), 2697-2705.

Sampson, C. y Jacobson, R.J. (1999) *Macrolophus caliginosus* Wagner (Heteroptera: Miridae): A predator causing damage to UK tomatoes. *IOBC/WPRS Bulletin* 22(1), 213-216.

Sanchez, J.A. (2008) Zoophytophagy in the plant bug *Nesidiocoris tenuis* (Heteroptera: Miridae). *Agricultural and Forest Entomology* 10(2), 75-80.

Schmidt, J.M., Taylor, J.R. y Rosenheim, J.A (1998) Cannibalism and intraguild predation in the predatory Heteroptera. Pp. 131-169. En Coll, M. y Ruberson, J.R. (Eds). *Predatory Heteroptera in Agroecosystems: Their Ecology and Use in Biological Control*. Thomas Say Publication of the Entomological Society of America, Lanham, MD.

Schmitz, O.J., Beckerman, A.P. y O'Brien, K. M. (1997) Behaviourally mediated trophic cascades: effects of predation risk on food web interactions. *Ecology* 78, 1388-1399.

Simon, C., Frati, F., Beckenbach, A., Crespi, B., Liu, H. y Flook, P. (1994) Evolution, weighting, and phylogenetic utility of mitochondrial gene sequences and a compilation of conserved polymerase chain reaction primers. *Annals of the Entomological Society of America* 87, 651-701.

Snyder, W.E. y Ives, A.R. (2001) Generalist predators disrupt biological control by a specialist parasitoid. *Ecology* 82, 705-716.

- Snyder, W.E. y Ives, A.R. (2003)** Interactions between specialist and generalist natural enemies: parasitoids, predators, and pea aphid biocontrol. *Ecology* 84, 91-107.
- Snyder, W.E., Clevenger, G.M. y Eigenbrode, S.D. (2004)** Intraguild predation and successful invasion by introduced ladybird beetles. *Oecologia* 140, 559-565.
- Solomon, M.G., Fitzgerald, J.D. y Murray, R.A. (1996)** Electrophoretic approaches to predator-prey interactions. Pp. 457-468. En Symondson W.O.C. y Liddell, J.E. (Eds). *The Ecology of Agricultural Pests: Biochemical Approaches*. Chapman & Hall, London.
- Sopp, P.I. y Sunderland, K.D. (1989)** Some factors affecting the detection period of aphid remains in predators using ELISA. *Entomologia Experimentalis et Applicata* 51, 11-20.
- Stansly, P.A., Calvo, J. y Urbaneja, A. (2005)** Release rates for control of *Bemisia tabaci* (Homoptera: Aleyrodidae) biotype "Q" with *Eretmocerus mundus* (Hymenoptera: Aphelinidae) in greenhouse tomato and pepper. *Biological Control* 35, 124-133.
- Symondson, W.O.C. (2002)** Molecular identification of prey in predator diets. *Molecular Ecology* 11, 627-641.
- Thompson, J.D., Gibson, T.J., Plewniak, F., Jeanmougin, F. y Higgins, D.G. (1997)** CLUSTAL X Windows interface: flexible strategies for multiple sequence alignment aided by quality analysis tools. *Nucleic Acids research* 25, 4876-4882.
- Tommasini, M.G., Burgio, G., Mazzoni, F. y Maini, S. (2002)** On intra-guild predation and cannibalism in *Orius insidiosus* and *Orius laevigatus* (Rhynchota Anthocoridae): laboratory experiments. *Bulletin of Insectology* 55(1-2), 49-54.

Torreno, H.S. y Magallona, E.D. (1994) Biological relationship of the bug, *Cyrtopeltis tenuis* Reuter (Hemiptera: Miridae) with tobacco. *The Philippine Entomologist* 9(4), 406-425.

Traugott, M., Zangerl, P., Juen, A. Schallhart, N. y Pfiffner, L. (2006) Detecting key parasitoids of lepidopteran pests by multiplex PCR. *Biological Control* 39, 39-46.

Traugott M., Bell J.R., Broad G.R., Powell W., Van Veen J.F., Vollhardt, I.M.G. y Symondson W.O.C. (2008) Endoparasitism in cereal aphids: molecular analysis of a whole parasitoid community. *Molecular ecology* 17, 3928-3938.

Traugott, M. y Symondson, W.O.C. (2008) Molecular analysis of predation on parasitized hosts. *Bulletin of Entomological Research* 98, 223-231.

Trottin-Caudal, Y., Fournier, C., Leyre, L.M., Ctifl, Chabrière, C. y Aprel (2006) La tomate sous serre dans le Sud-Est de la France: Protection contre la punaise *Nesidiocoris tenuis*. *Infos Ctifl* 224, 30-25.

Urbaneja, A., Montón, H. y Mollá, O. (2009) Suitability of the tomato borer *Tuta absoluta* as prey for *Macrolophus pygmaeus* and *Nesidiocoris tenuis*. *Journal of Applied Entomology* 133, 292-296.

van Baalen, M., Krivan, V., van Rijn, P.C.J. y Sabelis, M.W. (2001) Alternative food, switching predators, and the persistence of predator-prey systems. *American Naturalist* 157, 512-524.

Van der Blom, J. (2010) Applied entomology in Spanish greenhouse horticulture. *Proceedings of the Netherlands Entomological Society Meeting* 21, 9-17.

van der Hammen, T., de Roos, A.M., Sabelis, M.W. y Janssen, A. (2010) Order of invasion affects the spatial distribution of a reciprocal intraguild predator. *Oecologia*. 163, 79-89.

van Lenteren, J.C. y Martin, N.A. (1999) Biological control of whiteflies. Pp: 202-216. En Albajes, R., Gullino, M.L., van Lenteren, J.C. y Elad, Y. (Eds). *Integrated Pest and Disease Management in Greenhouse Crops*, Kluwer Academic Publishers, Dordrecht.

Videllet, P., Albajes, R. y Gabarra, R. (1997) Host-feeding activity of *Encarsia pergandiella* Howard on *Bemisia tabaci* (Gennadius). *IOBC/WPRS Bulletin* 20(4), 147-152.

Viggiani, G. y Mazzone, P. (1980) Sull'introduzione in Italia di *Encarsia pergandiella* Howard (Hymenoptera: Aphelinidae), parassita di *Trialeurodes vaporariorum* (Westw.) (Homoptera: Aleyrodidae). *Bolletino del Laboratorio Entomologia Agraria Filippo Silvestri* 37, 39-43.

von Berg, K., Traugott, M., Symondson, W.O.C. y Scheu, S. (2008) The effects of temperature on detection of prey DNA in two species of carabid beetle. *Bulletin of Entomological Research* 98, 263-269.

Walter, G.H. (1983) Divergent male ontogenies' in Aphelinidae (Hymenoptera: Chalcidoidea): a simplified classification and a suggested evolutionary sequence. *Biological Journal of the Linnean Society* 46, 261-282.

Walzer, A. y Schausberger, P. (1999) Cannibalism and interspecific predation in the phytoseiid mites *Phytoseiulus persimilis* and *Neoseiulus californicus*: predation rates and effects on reproduction and juvenile development. *BioControl* 43, 457-468.

Walzer, A., Moder, K. y Schausberger, P. (2009) Spatiotemporal within-plant distribution of the spider mite *Tetranychus urticae* and associated specialist and generalist predators. *Bulletin of Entomological Research* 99, 457-466.

Wheeler Jr., A.G. (2001) Biology of the Plant Bugs (Hemiptera: Miridae) Pests, Predators, Opportunists. Comstock Publishing Associates, Cornell University Press. Ithaca, NY.

Wise , D.H . (2006) Cannibalism, food limitation, intraspecific competition and the regulation of spider populations . *Annual Review of Entomology* 51, 441-465.

Wissinger, S.A., Sparks, G.B., Rouse, G.L., Brown, W.S. y Steltzer, H. (1996) Intraguild predation and cannibalism among larvae of detritivorous caddisflies in subalpine wetlands. *Ecology* 77, 2421-2430.

Zaidi, R.H., Jaal, Z., Hawkes, N.J., Hemingway, J. y Symondson, W.O.C. (1999) Can the detection of prey DNA amongst the gut contents of invertebrate predators provide a new technique for quantifying predation in the field? *Molecular Ecology* 8, 2081-2087.

Zhang, G.F., Lü, Z.C. y Wan, F.H. (2007) Detection of *Bemisia tabaci* remains in predator guts using a sequence-characterized amplified region marker. *Entomologia Experimentalis et Applicata* 123, 81-90.




Ilustración y diseño por Clara Herreros