

## CAPÍTULO 9

### REFERENCIAS BIBLIOGRÁFICAS

- Ábalos, M., Bayona, J.M. and Pawliszyn, J. (2000). Development of a headspace solid-phase microextraction procedure for the determination of free volatile fatty acids in waste waters. *Journal of Chromatography A.*, 873, 107-115.
- Abu-ghararah, Z.H. and Randall, C.W. (1991). The effect of organic compounds on biological phosphorus removal. *Wat. Sci. Tech.*, 23, 585-594.
- Albertson, O.E. (1987). The control of bulking sludges: from the early innovators to current practice. *Journal WPCF*, 59, 4, 172-182.
- Alleman, J.E. and Prakasam, T.B.S. (1983). Reflections on seven decades of activated sludge history. *Journal WPCF*, 55, 5, 436-443.
- Andreasen, K., Petersen, G., Thomsen, H. and Strube, R. (1997). Reduction of nutrient emission by sludge hydrolysis. *Water Sci. Tech.*, 35, 79-85.
- APHA-AWWA-WPCF (1995). *Standard Methods for the Examination of Water and Wastewater*. 19<sup>th</sup> edition, American Public Health Association. Washington, DC.
- Araújo, L.S, Catunda, P.F.C. and van Haandel, A.C. (1998). Biological sludge stabilisation Part 2: Influence of the composition of waste activated sludge on anaerobic stabilisation. *Water SA*, 24, 3, 231-236.
- Aravinthan, V., Mino, T., Takizawa, S., Satoh, H. and Matsuo, T. (2000). Sludge hydrolysate as a carbon source for denitrification. *1<sup>st</sup> World Congress of the International Water Association*, 3, 146-153, Paris 3-7 Julio 2000.
- Ardern, E. and Lockett, W.T. (1914). Experiments on the oxidation of sewage without the aid of filters. *J. Soc. Chem. Ind.*, 33, 523-525.
- Arthur, C.L., Killam, L.M., Motlagh, S., Lim, M., Potter, D.W. and Pawliszyn, J. (1992). Analysis of substituted benzene compounds in groundwater using solid-phase microextraction. *Environ. Sci. Technol.*, 26, 5, 979-983.
- Baeza, J., Gabriel, D. and Lafuente, F.J. (1996). Monitorización y control de una planta piloto de eliminación de nutrientes. 21 Curso de Ingeniería Ambiental: Eliminación Biológica de Nutrientes en Aguas Residuales. Lérida, 28-30 octubre, 187-224.
- Baeza, J., Gabriel, D. and Lafuente, F.J. (2002). In-line fase OUR (oxygen uptake rate) measurements for monitoring and control of WWTP. *Wat. Sci. Tech.*, 45, 4-5, 19-28.
- Banister, S.S. (1996). Optimisation of Primary Sludge Acidogenic Fermentation for Biological Nutrient Removal. M.Sc. Dissertation, Univ. of Pretoria, South Africa.
- Banister, S.S., Pitman, A.R. and Pretorius, W.A. (1998). The Solubilisation of N and P during Primary Sludge Fermentation and Precipitation of the Resultant P. *Water SA*, 24, 4, 337-342.
- Banister, S.S. and Pretorius, W.A. (1998). Optimisation of primary sludge acidogenic fermentation for biological nutrient removal. *Water S.A.* 24, 1, 35-41.
- Barajas, M.G., Knobelsdorf, J., Escalas, A. and Mujeriego, R. (2000). Hydrolysis and Fermentation in a Modified VFA-potential Test. Póster presentado en *Paris 2000 1<sup>st</sup> World Congress of the International Water Association*. Paris 3-7 Julio 2000. [CD-ROM] ISBN 2-9515416-0-0.
- Barajas, M.G. (2002). *Eliminación Biológica de nutrientes en un reactor biológico secuencial. Caracterización y estimulación de las fuentes de carbono del agua residual urbana*. Tesis Doctoral, Universidad Politécnica de Cataluña, España.

- Barajas, M.G., Knobelsdorf, J., Escalas, A. and Mujeriego, R. (2003). Solubilization and Fermentation in a Modified VFA-potential Method. *Environ. Eng. Sci.*, 20, 4, 329-336.
- Barnard, J.L. (1976). A review of biological phosphorus removal in the activated sludge process. *Water SA*, 2, 136-144.
- Barnard, J.L. (1983). Design Consideration Regarding Phosphate Removal in Activated Sludge Plants. *Water Sci. Technol.* 15, 319-328
- Barnard, J.L. (1984). Activated Primary Tanks for Phosphate Removal. *Water S.A.* 10, 121-126.
- Brinch, P.P., Rindel K. & Kalb, K. (1994). Upgrading to nutrient removal by means of internal carbon from sludge hydrolysis. *Wat. Sci. Tech.*, 29, 12, 31-40.
- Bécares, E. (1994). *Biología de una planta depuradora de fangos activados de doble etapa tratando aguas residuales de la industria farmacéutica*. Tesis Doctoral, Facultad de Biología, Área de Ecología de la Universidad de León, España.
- Bortone, G., Chech, J.S., Germirli, F., Bianchi, R. and Tilche, A. (1993). Experimental approaches for the characterisation of a nitrification/denitrification process on industrial waste water. Proc, 1<sup>st</sup> Int. Spec. Conf. on Microorganisms in Activated Sludge and Biofilm Processes, 129-136, Sept 27-28, Paris.
- Brdjanovic, D., Slamet, A., van Loosdrecht, M.C.M., Hooijmans, C.M., Alaerts, G.J. and Heijnen, J.J. (1998). Impact of excessive aeration on biological phosphorus removal from wastewater. *Wat. Res.*, 32, 1, 200-208.
- Brinch, P.P., Rindel, K. and Kalb, K. (1994). Upgrading to nutrient removal by means of internal carbon from sludge hydrolysis. *Wat. Sci. Tech.*, 29, 12, 31-40.
- Carlsson, H., Aspegren, H. and Hilmer, A. (1996). Interactions between wastewater quality and phosphorus release in the anaerobic reactor of the EBPR process. *Wat. Res.*, 30, 6, 1517-1527.
- Cecchi, F., Battistoni, P., Pavan, P., Fava, G. and Mata-Alvarez, J. (1994). Anaerobic digestion of OFMSW (organic fraction of municipal solid waste) and BNR (biological nutrient removal) processes: a possible integration- Preliminary Results. *Wat. Sci. Tech.*, 30, 8, 65-72.
- Charlton, J. (1994). Biological nutrient removal applied to weak sewage. *Wat. Sci. Tech.*, 29, 12, 41-48.
- Christensson, M. (1997). Enhanced biological phosphorus removal. Carbon source, nitrate as electron acceptor, and characterisation of the sludge community. Ph.D. Thesis, Department of Biotechnology, Lund University, Sweden.
- Christensson, M., lie, E., Jönsson, K., Johansson, P. and Welander, T. (1998). Hydrolysis and fermentation of sludge in primary clarifiers - A simple method to improve enhanced biological phosphorus removal. *Wat. Env. Res.*, 70, 2, 138-145.
- Chuang, S-H., Ouyang, C-F., Wang, Y-B. (1996). Kinetic Competition between Phosphorus Release and Denitrification on Sludge under Anoxic Condition. *Wat. Res.*, 30, 12, 2961-2968.
- Chudoba, J. (1985). Quantitative Estimation in COD Units of Refractory Organic Compounds Produced by Activated Sludge Microorganism. *Wat. Res.*, 19, 37-43.
- Cloete, T.E. and Steyn, P.L. (1988). A combined membrane filter-immunofluorescent technique for the in situ identification and enumeration of *Acinetobacter* in activated sludge. *Wat. Res.*, 22 (8), 961-969.
- Cloete, T.E. (1997). Introduction. In *Microbial Community Analysis: the Key to the Design of Biological Wastewater Treatment Systems* (eds. Cloete, T.E. and Muyima, N.Y.O.), Scientific and Technical Report, IAWQ, England.
- Cloete, T.E. and Muyima, N.Y.O. (1997). *Microbial Community Analysis: the Key to the Design of Biological Wastewater Treatment Systems*. Scientific and Technical Report, IAWQ, England.
- Collins, S.P., Pope, R.K., Scheetz, R.W., Ray, R.I., Wagner, P.A. & Little, B.J. (1993). Advantages of Environmental Scanning Electron Microscopy in studies of micro

- organisms. *Microscopy Research and Technique* 25: 398-405.
- Comeau, Y., Hall, K.J., Hancock, R.E.W. y Oldham, W.K. (1986). Biochemical model for enhanced biological phosphorus removal. *Wat. Res.*, 20, 1511-1521.
- Cortacáns, J.A. (1997). Investigación sobre la eliminación simultánea de nutrientes por vía biológica. Tesis doctoral por la Universidad Politécnica de Madrid, Madrid.
- Curtis, T.P. and Craine, N.G. (1998). The comparison of the diversity of activated sludge plants. *Wat. Sci. Tech.*, 37 (4/5), 71-78.
- Daigger, G.T., Robbins, M.H. and Marshall, B.R. (1985). The design of a selector to control low-F/M filamentous bulking. *Journal WPCF*, 57, 3, 220-226.
- Daigger et al., (1987). Factors affecting biological phosphorus removal for the VIP process, a high-rate University of Capetown type process. *Proc.IAWPRC Spec. Conf. Biological Phosphate Removal Wastewater*, Rome, Italy.
- Daigger, G.T., Waltrip, G.D., Romm, E.D. and Morales, L.M. (1988). Enhanced secondary treatment incorporating biological nutrient removal. *Journal WPCF*, 60, 10, 1833-1842.
- Daigger et al., (1990). Full-scale and pilot-scale experience with the VIP process. Paper presented at First Aust. Conf. Biol. Nutrient Removal, Bendigo, Aust.
- Danesh, S. and Oleszkiewicz, J.A. (1997). Use of a new anaerobic-aerobic reactor system to enhance biological phosphorus removal. *Wat. Sci. Technol.*, 35, 1, 137-144.
- Deakyne, C. W., Patel, M.A. and Krichten, D.J. 1(984). Pilot plant demonstration of biological phosphorus removal. *Journal WPCF*, 56, 7, 867-873.
- Dewisme, E. (1997). Características de las aguas residuales. En XV Curso sobre Tratamiento de Aguas Residuales y Explotación de Estaciones Depuradoras. CEDEX, Madrid.
- Directiva 91/271/CEE. DOCE L-35, 30-5-1991.
- Dold P.L., Ekama G.A. and Marais G.v.R., (1980). A general model for the activated sludge process. *Prog. Water Tech.*, 12, 47-77.
- Dold et al., (1986). Measurement of readily biodegradable cod fraction in municipal waste water by ultrafiltration. *UCT Report No. W57*, Dept. Civil Eng., Univ. of Cape Town, Rondebosch 7700, South Africa.
- Droste, R.L. (1997). *Theory and practice of water and wastewater treatment*. Jhon Wiley and Sons, Inc. Canada.
- Drysdale, G.D., Kasan, H.C. & Bux, F. (2000). Assessment of denitrification by the ordinary heterotrophic organisms in a NDBEPR activated sludge system. Proceedings of the 1st World Water Congress of the International Water Association (IWA), Paris 3-7 Jul 2000, 3:108-114.
- Eastman, J.A. and Ferguson, J.F., (1981). Solubilization of particulate organic carbon during the acid phase of anaerobic digestion. *Journal WPCF.*, 53, 352-366.
- Eikelboom, D.H. (1975). Filamentous organisms observed in activated sludge. *Wat. Res.*, 9, 365-388.
- Eikelboom, D.H. and van Buijsen, H.J.J. (1983). Microscopic sludge investigation manual. 20ed, TNO Research Institute for Environmental Hygiene, Delft, Holland.
- Eikelboom, D.H. (1991). *The role of competition between floc forming and filaments bacteria in bulking of activated sludge*. Biological Approach to Sewage Treatment Process: Current Status and Perspectives. Madoni, P.Perugia. Italy: Luiggi Bazzuchi Center, pp.143-149.
- Ekama, G.A., Marais G. v. R. (1977). The activated sludge process part ii. Dynamic behaviour. *Water SA*. 1, 18-50.
- Ekama, G.A. and Marais G. v. R. (1984). Nature of municipal wastewaters. Theory, design and operation of nutrient removal activated sludge process. Pretoria, South Africa, Water Research Commission.
- Ekama, G.A., Dold, P.L., Marais G.v.R. (1986). Procedures for determining influent COD fractions and the maximun specific growth rate of heterotrophs in activated sludge systems. *Wat. Sci. Technol.*, 18, 91-114.

- Ekama G.A. and Wentzel M.C., (1999). Difficulties and developments in biological nutrient removal technology and modelling. *Wat. Sci. Tech.*, 39, 6, 1-11.
- Elefsiniotis P. and Oldham W.K. (1994). Influence of pH on the acid-phase anaerobic digestion of primary sludge. *J. Chem. Tech. Biotechnol.*, 60, 89-96.
- EMASESA, (1997). *Microorganismos Filamentosos en el Fango Activado*. Varios autores. Ayuntamiento de Sevilla, España.
- EPA (1982). *Handbook for sampling and sample preservation of water and wastewater*. EPA-600/4-82-029. Environmental Monitoring and Support Lab, Cincinnati, Ohio, Estados Unidos, 402.
- EPA (1987). Design manual phosphorus removal. EPA/625/1-87/001.
- EPA (1992). *Wastewater Treatment/Disposal for Small Communities*. EPA/625/R-92/005. United States Environmental Protection Agency, Washington D.C., EEUU.
- EPA (1993). Manual nitrogen control. EPA/625/R-93/010.
- Escaler, I. (1997). Eliminació biològica de nutrients (nitrógen i fósfor) mitjançant el procé discontinu de fangs activats. Tesina de especialidad, ETSECCPB, UPC.
- Fernández-Galiano, D., Guinea, A., Serrano, S., Martín, M., Arregui, L., Rodríguez, B., Campos, I., Calvo, P. and Suárez, J. (1996). Guía práctica de identificación de protozoos ciliados en estaciones depuradoras de aguas residuales por lodos activados de la Comunidad Autónoma de Madrid. Dpto. Microbiología, Facultad de Biología, Universidad Complutense de Madrid.
- Fuhs, G. W. and Chen, M. (1975). Microbiological basis of phosphate removal in the activated sludge process for the treatment of wastewater. *Microb. Ecol.*, 2, 119-138.
- García J. (1996). *Eliminació de Matèria Orgànica i Nutrients en Llacunes d'Alt Rendiment*. Tesis doctoral, Departament de Productes Naturals, Biología Vegetal i Edafología de la Facultat de Farmàcia, Universitat de Barcelona.
- García, P.A. and Fernández-Polanco, F. (1996). Parámetros de operación en sistemas de eliminación de nutrientes. En: *2n Curs d'Enginyeria Ambiental. Eliminació de nutrients en aigües residuals*. Xavier Flotats. Lleida: Paperkite Editorial.
- Ginestet, P. Maisonnier, A. and Spérando, M. (2002). Wastewater COD characterization: biodegradability of physico-chemical fractions. *Wat. Sci. Tech.*, 45, 6, 89-97.
- Gerber, A., Mostert, E.S., Winter, C.T. and de Villiers, R.H. (1987). Interactions between phosphate, nitrate and organic substrate in biological nutrient removal process. *Wat. Sci. Tech.*, 19, 183-194.
- Goel, R., Mino, T., Satoh, H., Matsuo T. (1998). Intracellular Storage Compounds, Oxygen Uptake Rates and Biomass Yield with Readily and Slowly Degradable Substrates. *Wat. Sci. Tech.*, 38, 8-9, 85-93.
- Gonçalves, R.F., Charlier, A.C., Sammut, F. (1994). Primary fermentation of soluble and particulate organic matter for wastewater treatment. *Wat. Sci. Tech.*, 30, 6, 53-62.
- Grady, C.P.L. Jr. and Lim, H.C. (1980). Biological wastewater treatment, theory and applications. Marcel Dekker, New York., pp. 895.
- Gujer, W., Henze, M., Mino, T., Matsuo, T., Wentzel, M.C. and Marais, G.v.R. (1995). The activated sludge model No. 2: biological phosphorus removal. *Wat. Sci. Tech.*, 31, 2, 1-11.
- Gujer, W., Henze, M., Mino, T., and Van Loosdrecht, M. (1999). Activated sludge model No. 3. *Wat. Sci. Tech.*, 39, 1, 183-193.
- Hallin, S. and Pell, M. (1998). Metabolic properties of denitrifying bacteria adapting to methanol and ethanol in activated sludge. *Wat. Res.*, 32, 1, 13-18.
- Hart, M.A. and Melmed, L.N. (1982). Microbiology of nutrient removing activated sludge. *Wat. Sci. Tech.*, 14 (9/11), 1501-1502.
- Hatziconstantinou, G.J., Yannakopoulos, P. and Andreadakis, A. (1996). Primary sludge hydrolysis for biological nutrient removal. *Wat. Sci. Tech.*, 34 (1/2), 417-423.
- Helmer, C. and Kunst, S. (1998). Low temperature effects on phosphorus release and uptake by microorganisms in EBPR plants. *Wat. Sci. Tech.*, 37 (4/5), 531-539.

- Henze, M. (1986). Nitrate versus Oxygen Utilization Rates in Wastewater and Activated Sludge Systems, *Wat. Sci. Tech.*, 18, 6, 115.
- Henze, M., Grady, C.P.L. Jr., Gujer, W., Marais, G.v.R., Matsuo, T. (1987). Activated Sludge Model No 1, *IAWPRC Scientific and Technical Reports*, No 1, IAWPRC, Londres.
- Henze, M. (1991). Capabilities of biological nitrogen removal from wastewater. *Water Sci. Technol.*, 23, 4-6, pp. 669-679.
- Henze, M. et al. (1992). Characterization of Wastewater for Modelling of Activated Sludge Processes. *Water Sci. Technol.* 25, 6, 1-15.
- Henze, M., Gujer, W., Mino, T., Matsuo, T., Wentzel, M.C. and Marais, G.v.R. (1995a). Activated Sludge Model No. 2, *IAWPRC Scientific and Technical Reports*, No 3, IAWPRC, Londres.
- Henze, M., Gujer, W., Mino, T., Matsuo, Wentzel, M.C., Marais, G.v.R. (1995b). Wastewater and biomass characterization for the activated sludge model No. 2: Biological phosphorus removal. *Wat. Sci. Tech.*, 31, 2, 13-23.
- Henze, M., Harremoes, P., Cour Janse, J. and Arvin, E. (1997). *Wastewater Treatment: Biological and Chemical Processes*. 2ed. Springer-Verlag Berlin.
- Horan, N.J. (1993). Biological wastewater treatment systems: theory and operation. Jhon Wiley and Sons, Ltd, England.
- HRSD (1991). VIP, Virginie Initiative Plant. Informe de la Hampton Roads Sanitation District, Virginia.
- Hvitved-Jacobsen, T., Raunkjær, K. and Nielsen, P.H. (1995). Volatile fatty acids and sulfide in pressure mains. *Wat. Sci. Tech.*, 31, 7, 169-179.
- Irvine, R.L., Wilderer, P. A., Flemming, H.C. (1997). Controlled Unsteady State Processes and Technologies- an Overview. *Wat. Sci. Tech.*, 35, 1, 1-10.
- Iwema, A. and Meunier, A. (1985). Influence of nitrate on acetic acid induced biological phosphate removal. *Wat. Sci. Tech.*, 17, 11, 289-294.
- Janssen, P. M. (1994). Operating experiences on two full-scale plants, retrofitted for biological phosphorus removal. Nutrient removal from wastewaters. Horan, N. J., P. Lowe and E. Stentiford, Editors. Pennsylvania: Technomic Publishing Co., Inc.
- Jardin, N. and Pöpel, H.J. (1997). Waste activated sludge production of the enhanced biological phosphorus removal process. *Wat. Env. Res.*, 69, 3, 375-381.
- Jenkins, D., Richard, M.G. and Daigger, G.T. (1986). Manual on the causes and control of activated sludge bulking and foaming. Water Research Commission, Pretoria, Sur Africa.
- Jenkins, D. and Tandoi, V. (1991). The applied microbiology of enhanced biological phosphate removal- accomplishment and needs. *Wat. Res.*, 25, 12, 1471-1478.
- Jenkins, D. (1992). Towards a Comprehensive Model of Activated Sludge Bulking and Foaming. *Wat. Sci. Tech.*, 25, 215-230.
- Jenkins, D., Richard M. G. and Daigger, G. T. (1993). *Manual on the causes and Control of Activated Sludge Bulking and Foaming*, 2<sup>a</sup>. Ed. Lewis Publishers, Michigan, USA.
- Jorgensen, P.E. Bilogical Hydrolysis of Sludge from Primary Precipitation. En Hahn, H.H., Klute, R., editors, *Chemical Water and Wastewater Treatment*, 521-529. Springer-Verlag, Berlin, Heidelberg.
- Kappeler, J. and Gujer W. (1992). Estimation of Kinetic Parameters of Heterotrophic Biomass under Aerobic Conditions and Characterization of Wastewater for Activated Sludge Modelling. *Wat. Sci. Tech.*, 25, 6, 125-140.
- Karlsson, I. and Smith, G. (1990). Pre-precipitation facilitates nitrogen removal without tank expansion. *Wat. Sci. Tech.*, 22, (7/8), 85-92.
- Kaufman, L. and P. J. Rousseeuw 1990. Finding groups in data: an introduction to cluster analysis. New York: John Wiley & Sons.
- Kristensen, G.H., Jorgensen, P.E., Henze, M. (1992). Characterisation of Functional Microorganism Groups and Substrate in Activated Sludge and Wastewater by AUR, NUR and OUR. *Wat. Res.*, 25, 6, 43-57.

- Lee, S.E., Koopman, B.L., Jenkins, D., Lewis, R.F. (1982). The Effect of Aeration Basin Configuration on Activated Sludge Bulking at Low Organic Loading. *Wat. Sci. Tech.*, 14, 407-427.
- Lee, S.III, Koopman, B., Park, S. and Cadee, K. (1995). Effect of fermentation wastes on denitrification in activated sludge. *Wat. Env. Res.*, 67, 7, 1119-1122.
- Lee, N. M. (1996). *Parameters Affecting Microorganisms and the Process Performance in Biological Wastewater Treatment*. Tesis Doctoral. Department of Biotechnology, Lund University, Suecia.
- Lee, S.E., Kim, K.S., Ahn, J.H. and Kim, C.W. (1997). Comparison of phosphorus removal characteristics between various biological nutrient removal processes. *Wat. Sci. Tech.*, 36, 12, 61-68.
- Lesouef, A., Payraudeau, M. Rogalla, F., Kleiber, B. (1992). Optimizing Nitrogen Removal Reactor Configurations by On-site Calibration of the IAWPRC Activated Sludge Model. *Wat. Sci. Tech.*, 25, 6, 105-124.
- Levin, G.V. and Shapiro, J. (1965). Metabolic uptake of phosphorus by wastewater organisms. *Journal WPCF*, 36(6), 800-821.
- Lie, E. (1996). Limiting factors in biological nutrient removal from wastewater. Ph.D. Thesis, Department of Biotechnology, Lund University, Sweden.
- Lie, E. and Welander, T. (1997). A method for determination of the readily fermentable organic fraction in municipal wastewater. *Wat. Res.*, 31, 6, 1269-1274.
- Lo, C.K., Yu, C.W., Tam, N.F.Y. and Traynor, S. (1994). Enhanced nutrient removal by oxidation-reduction potential (ORP) controlled aeration in a laboratory scale extended aeration treatment system. *Wat. Res.*, 28, 10, 2087-2094.
- Lötter, L.H. and Murphy, M. (1985). The identification of heterotrophic bacteria in an activated sludge plant with particular reference to polyphosphate accumulation. *Wat. SA*, 11 (4), 179-184.
- Ludzack, F.L. and Ettinger, M.B. (1962). Controlling operation to minimize activated sludge effluent nitrogen. *J.Wat.Pol.Con.Fed.*, 34, 920.
- Mamais, D., Jenkins, D. and Pitt, P. (1993). A rapid physical-chemical method for the determination of readily biodegradable soluble COD in municipal wastewater. *Water Res.*, 27, 195-197.
- Marten, W.L. and Daigger, G.T. (1997). Full-scale evaluation of factors affecting the performance of anoxic selectors. *Water Envir. Res.*, 69, 7, 1272-1281.
- Martin Ruel, S. Comeau, Y., Héduit, A. Deronzier, G., Ginestet, P. And Audic, J.M. (2002). Operating conditions for the determination of the biochemical acidogenic potential of wastewater. *Wat. Res.*, 36, 2337-2341.
- Masdeu, M. (1997). La depuradora com a sistema biològic. Formació i estructura dels flocs. Anàlisi Microbiològica de Fangs Activats. Mas Carandell, Reus.
- Matia, L., Martín, J., Ventura, F. and Campos, C. (1999). La qualitat dels recursos no convencionals. En Xarxa Temática: Recursos d'aigua. (eds. M. Salgot, X. Sánchez y A. Torrens). Fundación Agbar.
- Metcalf-Eddy, (1991). *Wastewater Engineering, Treatment, Disposal and Reuse*. 3<sup>rd</sup> edition. McGraw-Hill Inc. New York.
- Metcalf and Eddy (1995). Ingeniería de aguas residuales: tratamiento, vertido y reutilización. 30 edición, McGraw-Hill Inc., Interamericana de España S.A.
- Mikawa, K., Emori, H., Takeshima, T., Ishiyama, E. and Tanaka, K. (1996). High rate and compact two-stage post-denitrification process with single-sludge pre-denitrification. *Water Sci. Technol.*, 34, 1-2, 467-475.
- Mino, T., Arun, V., Tsuzuki, Y., Matsuo, T. (1987) Effect of Phosphorus Accumulation on Acetate Metabolism in the Biological Phosphorus Removal Process. In: *Biological phosphate removal from wastewaters*, R. Ramadori, Pergamon Press, 27-38.
- Mino, T., San Pedro, C. and Delfin, Matsuo, T. (1995). Estimation of the Rate of Slowly Biodegradable COD (SBCOD) Hydrolysis under Anaerobic, Anoxic and Aerobic

- Conditions by Experiments using Starch as Model Substrate. *Water Sci. Tech.* 31, 2, 95-103.
- Monteith, H.D. and Stephenson, J.P. (1981). Mixing efficiencies in full-scale anaerobic digesters by tracer methods. *Journal WPCF*, 53, 1, 78-84.
- Montgomery, D.C., Runger, G.C. (1996). *Probabilidad y estadística aplicadas a la Ingeniería*. Mèxic D.F. (Mèxic): McGraw-Hill.
- Moriyama, K., Sato, K., Harada, Y., Washiyama, K., Okamoto, K. (1990). Simultaneous Biological Removal of Nitrogen and Phosphorus using Oxic-Anaerobic-Oxic Process. *Water Sci. Technol.*, 22, 7-8, 61-66.
- Mujeriego, R., Gaztelu, M., Jové, J.M. y Bravo, J.M. (1984). El tractament de les aigües residuals en els petits municipis de Barcelona. Diputació de Barcelona. 481 pp.
- Mujeriego, R. (1990). *Riego con Agua Residual Municipal Regenerada*. Mujeriego, R. ed. Universitat Politècnica de Catalunya. Barcelona. Diputació de Barcelona. pp. 145.
- Mujeriego, R. (2005). La fiabilidad de los procesos de regeneración del agua. *II Jornadas Técnicas de Gestión de Estaciones Depuradoras de Aguas Residuales*. Agencia Catalana del Agua, Generalitat de Catalunya, 18-20 enero 2005, Barcelona.
- Mullis, M.K., Schroeder E.D. (1971). A Rapid Biochemical Oxygen Demand Tets Suitable for Operational Control. *Journal WPCF*, 43, 2, 209-215.
- Münch, E. and Koch, A. (1999). A Survey of Prefermenter Design, Operation and Performance in Australia and Canada. *Water Sci. Technol.* 39, 6, 105-112.
- Muyima, N.Y., Momba, M.N.B. and Cloete, T.E. (1997). Biological methods for the treatment of wastewaters. *Microbial Community Analysis: The Key to the Design of Biological Wastewater Treatment Systems*. Scientific and Technical Report, IAWQ, Inglaterra.
- Narkis, N., Henefeld-Fourrier, S. and Rebhun, M. (1980). Volatile organic acids in raw wastewater and in physico-chemical treatment. *Wat. Res.*, 14, 9, 1215-1223.
- Neethling, J.B. (1995). Biological nutrient removal with activated sludge systems. HDR Engineering Inc., El Dorado Hills.
- Nicholls, H. A., Pitman A.R. and Osborn D.W. (1985). The readily biodegradable fraction of sewage: its influence on phosphorus removal and measurement. *Water Sci. Tech.* 17, Paris, 73-87.
- Nielsen, P.H., Andreasen, K., Wagner, M., Blackall, L.L., Helmer, H. and Seviour, R.J. (1998). Variability of type 021N in activated sludge as determined by *in situ* substrate uptake pattern and *in situ* hybridization with fluorescent rRNA targeted probes. *Wat. Sci. Tech.*, 37 (4/5), 423-430.
- Oldham, W.K. and Stevens, G.M. (1985). Operating Experiences with the Kelowna Pollution Control Centre. *Proceedings of the Seminar on Biological Phosphorus Removal in Municipal Wastewater Treatment*. Penticton, British Columbia, Canadá, Abril 17 y 18.
- Oliveira, M.A., Reis, E.M. and Nozaki, J. (2001). Biological treatment of wastewater from the cassava meal industry. *Environ. Res.*, Section A, 85, 177-183.
- Oliveras, J. (1998). *Els Microorganismes Filamentosos i els Protozoos en la Depuració d'Aigües del Sector Industrial Carni*. Tesis Doctoral. Universitat Politècnica de Catalunya (España).
- Orhon, D. and Artan, N. (1994). *Modelling of activated sludge systems*. Lancaster: Technomic Publ. Co.Inc.
- Orhon, D. Kabdasli, I., Ubay Çokgör, E., Tünay, O. and Artan, N. (2002). Experimental assessment of optimum operation strategy for large industrial wastewater treatment plants- A case study. *Env. Eng. Sci.*, 19, 1, 47-58.
- Osborn, D.W., Nicholls, H.A. (1978). Optimisation of the Activated Sludge Process for the Biological Removal of Phosphorus. *Progress in Water Technology*, 10, 261-277.
- Ostgaard, K., Christensson, M.. Lie E., Jönsson, K., Welander, T. (1997). Anoxic Biological Phosphorus Removal in a Full-Scale UCT Process. *Water Research*, 31, 11, 2719-2726.

- Ouyang, C.F., Chuang, S.H. and Su, J.L. (1999). Nitrogen and phosphorus removal in a combined activated sludge- RBC process. *Proc. Natl. Sci. Counc. ROC(A)*, 23, 2, 181-204.
- Park J.K., Wang J., and Novotny G., (1997). Wastewater characterisation for evaluation of biological phosphorus removal. Wisconsin Department of Natural Resources, Research Report, 174, 29 ([www.dnr.state.wi.us](http://www.dnr.state.wi.us)).
- Parody, D.F. (1997). Microorganismos filamentosos. Microorganismos Filamentosos en el Fango Activado. EMASESA, Ayuntamiento de Sevilla, 23-95.
- Pitman, A.R., Venter, S.L.V., Nicholls, H.A. (1983). Practical Experience with Biological Phosphorus Removal Plants in Johannesburg. *Water Sci. Technol.* 15, 3-4, 233-259.
- Pitman, A.R. (1991). Design Considerations for Nutrient Removal Activated Sludge Plants. *Water Sci. Tech.* 23, 781-790.
- Punrattanasin, W. (1997). *Investigation of the effects of COD/TP ratio on the performance of a biological nutrient removal system*. MSc. Thesis, Virginia Polytechnic Institute and State University, Blacksburg, Virginia, USA.
- Qasim, S.R., Chiang, W., Zhu, G. and Miller, R. (1996). Effect of biodegradable carbon on biological phosphorus removal. *J. Environ. Eng.*, September, 875-878.
- Quintela, M. and Ro, K.S. (2003). Atmospheric oxygen transfer during in situ oxygen uptake rate measurements. *J. Env. Eng.*, 129, 2, February 1, 183-186.
- Randall, C.W., Barnard, J.L. and Stensel, H.D. (1992). Design and retrofit of wastewater treatment plants for biological nutrient removal. Water Quality Management Library, Vol 5. Technomic, Inc., Pennsylvania, USA.
- Randall A.A. and Chapin, R.W. (1997). Acetic acid inhibition of biological phosphorus removal. *Wat. Env. Res.*, 69, 5, 955-960.
- Raunkjær, K., Hvítved-Jacobsen, T., Nielsen, P.H. (1995). Transformation of Organic Matter in a Gravity Sewer. *Wat. Env. Res.*, 67, 181-188.
- Rensink, J.H. and Donker, H.J.G.W. (1993). Management of bulking sludge control by selector. In prevention and control of bulking activated sludge, pp. 77-97. Luiggi Bazzuchi Center, Perugia, Italia.
- Richard, M. (1989). Activated Sludge Microbiology. WPCF, Virginia, USA.
- Rivas Mijares, G. (1978). Tratamiento de Aguas Residuales. 20 edición, Ediciones Vega, Venezuela.
- Rius, M.M. (2003). Caracterización del proceso de depuración de aguas residuales y dinámica de las comunidades de microorganismos en plantas de tipo SBR. Tesis doctoral. Universidad de Barcelona.
- Roeleveld, P.J. and van Loosdrecht, M.C.M. (2002). Experience with guidelines for wastewater characterization in The Netherlands. *Wat. Sci. Tech.*, 45, 6, 77-87.
- Romero, E. (1999). *Characterisation and thermo-hydro-mechanical behaviour of unsaturated Boom clay. An experimental study*. Tesis doctoral. Universitat Politècnica de Catalunya.
- Rössle, W.H. and Pretorius, W.A. (2001a). A review of characterisation requirements for in-line prefermenters Paper 1: Wastewater Characterisation. *Water SA*, 27, 3, 405-412.
- Rössle, W.H. and Pretorius, W.A. (2001b). A review of characterisation requirements for in-line prefermenters Paper 2: Process Characterisation. *Water SA*, 27, 3, 413-422.
- Sabalowsky, A.R. (1999). *An investigation of the feasibility of nitrification and denitrification of a complex industrial wastewater with high seasonal temperatures*. MSc. Thesis, Virginia Polytechnic Institute and State University, Blacksburg, Virginia, USA.
- Salgot, M. and Tapias, J.C. (1999). Aigües superficials. Xarxa Temática: Recursos d'aigües. (eds. M. Salgot, X. Sánchez y A. Torrens). Fundación Agbar.
- Salsky, D.S., Daigger, G.T. (1995). Wastewater Solids Fermentation for Volatile Acid Production and Enhanced Biological Phosphorus Removal. *Water Environ. Res.* 67, 2, 230-237.

- Salvadó, H. (1990). *Estudi de Protozoos Ciliats en Plantes de Tratament Biològic de les Aigües Residuials*. Tesis Doctoral. Universitat de Barcelona.
- Salvadó, H. (1999). Observación Microscópica del Fango (I): Identificación y Significación de Protozoos y Metazoos. En: *Curs d'Identificació i Control de Microorganismes Filamentosos en Fangs Activats*. Facultat de Farmacia, Universitat de Barcelona, España.
- Sanpedro, D.C., Mino, T., Matsuo, T. (1995). Evaluation of the Rate of Hydrolysis of Slowly Biodegradable COD (SBCOD) using as Substrate under Anaerobic, Anoxic and Aerobic Conditions. *Wat. Sci. Tech.*, 30, 11, 191-199.
- Sedlak, R. (1991). Phosphorus and nitrogen removal from municipal wastewater: principles and practice. 20 edición. Lewis Publishers, New York, USA.
- Sezgin, M., Jenkins, D., Parker, D.S. (1978). A Unified Theory of Filamentous Activated Sludge Bulking. *Journal WPCF*, **50**, 362-381
- Sharman, R. (1998). Water and Wastewater Technology, LBCC. <http://www.lbcc.cc.or.us/process110/8/97>
- Siebritz, I.P., Ekama, G.A., Marais, G. v. R. (1983) A Parametric Model for Biological Excess Phosphorus Removal. *Wat. Sci. Tech.*, **15**, 127-152.
- Smolders, G.J.F., van der Meij, J., van Loosdrecht, M.C.M., Heijnen, J.J. (1994). Model of the Anaerobic Metabolism of the Biological Phosphorus Removal Process: Stoichiometry and pH Influence. *Biotechnology and Bioengineering*, 43:461-470.
- Smolders, G.J.F., van Loosdrecht, M.C.M., Heijnen, J.J. (1995) A Metabolic Model for the Biological Phosphorus Removal. *Wat. Sci. Tech.*, **31**, 2, pp. 79-93.
- Sollfrank, U. and Gujer, W. (1991). Characterisation of domestic wastewater for mathematical modelling of the activated sludge process. *Wat. Sci. Tech.*, 23, Kyoto, 1057-1066.
- Spanjers, H. and Vanrolleghem, P. (1995). Respirometry as a tool for rapid characterization of wastewater and activated sludge. *Wat. Sci. Tech.* 31, 2, 105-114.
- Spérando, M. (1998). Développement d'une procédure de compartimentation d'une eau résiduaire urbaine et application à la modélisation dynamique de procédés à boues activées. *PhD Thesis* no. 488. INSA, Toulouse, France.
- Spérando, M. and Paul, E. (2000). Estimation of wastewater biodegradable COD fractions by combining respirometric experiments in various So/Xo ratios. *Wat. Res.*, 34, 4, 1233-1246.
- Stainer, R.Y., Ingraham, J.L., Wheelis, M.L., Painter, P.R. (1991). *Microbiología*. 2<sup>a</sup>. edición. Reverté: Barcelona.
- Standard Methods, (1995). Standard Methods for the Examination of Water and Wastewater. APHA – AWWA – WPCF, Washington, D.C.
- Stepan, S.F., Smith, J.F. and Reukers, J. (1978). Apparatus for on-site extraction of organic compounds from water. *Wat. Res.*, 12, 447-449.
- Stern, L.B., Marais G. v R. (1974). Sewage as Electron Donor in Biological Desnitritification. *Research Report W7*. Dept. Civil Eng., Univ. of Cape Town, Rondebosch 7700, South Africa.
- Suárez, J. (1997). La selección biológica como método de control del esponjamiento producido por microorganismos filamentosos. *Microorganismos Filamentosos en el Fango Activado*. EMASESA, Ayuntamiento de Sevilla, 97-125.
- SWCS (2000). *Sinopsis 1. Eutrophication of lakes and effects of global warming, acid deposition and UV-B penetration*. <http://www.chebucto.ns.ca/Science/SWCS/SWCS.html>
- Teira, M.R. (1996). Bases microbiológicas. Microorganismos nitrificantes, desnitrificantes y con capacidad para acumular fósforo. 21 Curso de Ingeniería Ambiental: Eliminación Biológica de Nutrientes en Aguas Residuales. Lérida 28-30 de octubre, 1-21.

- Torrijos, M., Cerro, RM., Capdeville, B., Zeghal, S., Payraudeau M. and Lesouef A. (1994). Sequencing Batch Reactor: Tool for Wastewater Characterization for the IAWPRC Model. *Water Sci. Technol.* 29, 7, 81-90.
- UPC (1985). Taules d'estadística. Escola Técnica Superior d'enginyers insustrials de Terrassa. Universitat Politècnica de Catalunya.
- Urbain, V., Mobarry, B., De Silva, V., Stahl, D.A., Rittmann, B.E. and Manem, J. (1998). Integration of performance, molecular biology and modelling to describe the activated sludge process. *Wat. Sci. Tech.*, 37(4/5), 223-229.
- Van Haandel, A.C. (1994). Influence of the digested COD concentration on the alkalinity requirement in anaerobic digesters. *Wat. Sci. Tech.*, 30, 8, 23-34.
- Wanner, J. (1994). *Activated Sludge Bulking and Foaming Control*. Technomic Publishing Co. Inc. Lancaster.
- Wanner, J. (1997). Microbial population dynamics in biological wastewater treatment plants. In *Microbial Community Analysis: The Key to the Design of Biological Wastewater Treatment Systems. Scientific and Technical Report*, IAWQ, Inglaterra.
- Wanner, J. (1998). Beginning of the activated sludge. *Acivated Sludge Pages*, Nº 1. <http://scitrav.com/wwater/asp1/begin.htm>
- Water Quality International (1996). Nutrient removal. 18th Biennial Conference of the IAWQ, 23-28 Junio, Singapore. Water Science and Technology.
- Wentzel, M.C., Dold, P.L., Ekama, G.A., Marais, G.v.R. (1985). Kinetics of Biological Phosphorus Release. *Water Sci. Technol.* 17, 57-71.
- Wentzel, M.C., Loewenthal, R.E., Ekama, G.A., Marais, G.v.R. (1989). Enhanced Polyphosphate Organism Cultures in Activated Sludge Systems. Part 1: Enhanced Culture Development. *Water SA*, 14, 81-92.
- Wentzel, M.C., Ekama, G.A., Dold, P.L., Marais, R. (1990). Biological Excess Phosphorus Removal –Steady State Process Desing. *Water SA*, 16, 1, pp. 29-48.
- Wentzel, M.C., Ekama, G.A. and Marais, G.v.R. (1991). Kinetics of nitrification denitrification biological excess phosphorus removal systems- A review. *Wat. Sci. Tech.*, 23, Kyoto, 555-565.
- Wentzel, M.C., Mbewe A. and Ekama, G.A. (1995). Batch test for measurement of readily biodegradable cod and active organism concentrations in municipal waste waters. *Water SA*, 21, 2, 177-123.
- Wetzel R.G., (1983). Limnology 2nd Edition, Saunders College Publishing, Philadelphia, pp. 679.
- Winkler, M.A. (1998). *Tratamiento Biológico de Aguas de Desecho*. Limusa SA (eds.) Mexico D.F.
- Witteborg, A., Last, A., Hamming, R., Hemmers, I. (1996). Respirometry for Determination of the Influent Ss- Concentration. *Water Sci. Technol.* 33, 1, 311-323.
- Ziglio, G., Andreottola, G., Foladori, P. and Ragazzi, M. (2001). Experimental validation of a single-OUR method for wastewater RBCOD characterisation. *Wat. Sci. Tech.*, 43, 11, 119-126.