



UNIVERSITAT DE BARCELONA



Receptores Monoaminérgicos en Corteza Prefrontal: Mecanismo de Acción de Fármacos Antipsicóticos

Tesis Doctoral presentada por

Noemí Santana Ramos

Barcelona, julio de 2007

VII. BIBLIOGRAFIA

- Abi-Dargham A, Mawlawi O, Lombardo I, Gil R, Martinez D, Huang Y, Hwang DR, Keilp J, Kochan L, Van Heertum R, Gorman JM, Laruelle M (2002) Prefrontal dopamine D1 receptors and working memory in schizophrenia. *J Neurosci* 22: 3708-3719.
- Acosta-Martinez M, Fiber JM, Brown RD, Etgen AM (1999) Localization of alpha(1B)-adrenergic receptor in female rat brain regions involved in stress and neuroendocrine function. *Neurochemistry International* 35: 383-391.
- Adell A, Artigas F (2004) The somatodendritic release of dopamine in the ventral tegmental area and its regulation by afferent transmitter systems. *Neurosci Biobehav Rev* 28: 415-431.
- Adell A, Myers RD (1995) Selective destruction of midbrain raphe nuclei by 5,7-DHT: is brain 5-HT involved in alcohol drinking in Sprague-Dawley rats? *Brain Res* 693: 70-79.
- Aghajanian G.K (1995) Electrophysiology of serotonin receptor subtypes and signal transduction mechanisms. In: Bloom FE, Kupfer DJ, editors. *Psychopharmacology: the fourth generation of progress*. New York: Raven Press. p 451-460.
- Aghajanian GK, Marek GJ (1997) Serotonin induces excitatory postsynaptic potentials in apical dendrites of neocortical pyramidal cells. *Neuropharmacology* 36: 589-599.
- Akbarian S, Kim JJ, Potkin SG, Hagman JO, Tafazzoli A, Bunney WE, Jr., Jones EG (1995) Gene expression for glutamic acid decarboxylase is reduced without loss of neurons in prefrontal cortex of schizophrenics. *Arch Gen Psychiatry* 52: 258-266.
- Albanese A, Minciacchi D (1983) Organization of the ascending projections from the ventral tegmental area: a multiple fluorescent retrograde tracer study in the rat. *J Comp Neurol* 216: 406-420.
- Amargos-Bosch M, Adell A, Bortolozzi A, Artigas F (2003) Stimulation of alpha(1)adrenoceptors in the rat medial prefrontal cortex increases the local in vivo 5-hydroxytryptamine release: reversal by antipsychotic drugs. *Journal of Neurochemistry* 87: 831-842.
- Amargos-Bosch M, Bortolozzi A, Puig MV, Serrats J, Adell A, Celada P, Toth M, Mengod G, Artigas F (2004) Co-expression and in vivo interaction of serotonin(1A) and serotonin(2A) receptors in pyramidal neurons of prefrontal cortex. *Cerebral Cortex* 14: 281-299.
- Araneda R, Andrade R (1991) 5-Hydroxytryptamine₂ and 5-Hydroxytryptamine_{1A} Receptors Mediate Opposing Responses on Membrane Excitability in Rat-Association Cortex. *Neuroscience* 40: 399-412.
- Ariano MA, Wang J, Noblett KL, Larson ER, Sibley DR (1997a) Cellular distribution of the rat D-1B receptor in central nervous system using anti-receptor antisera. *Brain Research* 746: 141-150.
- Ariano MA, Wang J, Noblett KL, Larson ER, Sibley DR (1997b) Cellular distribution of the rat D-4 dopamine receptor protein in the CNS using anti-receptor antisera. *Brain Research* 752: 26-34.
- Arnsten AF, Goldman-Rakic PS (1984) Selective prefrontal cortical projections to the region of the locus coeruleus and raphe nuclei in the rhesus monkey. *Brain Res* 306: 9-18.

- Arnsten AF, Goldman-Rakic PS (1998) Noise stress impairs prefrontal cortical cognitive function in monkeys: evidence for a hyperdopaminergic mechanism. *Arch Gen Psychiatry* 55: 362-368.
- Arnsten AF, Mathew R, Ubriani R, Taylor JR, Li BM (1999) Alpha-1 noradrenergic receptor stimulation impairs prefrontal cortical cognitive function. *Biol Psychiatry* 45: 26-31.
- Arnt J, Skarsfeldt T (1998) Do novel antipsychotics have similar pharmacological characteristics? A review of the evidence. *Neuropsychopharmacology* 18: 63-101.
- Ashby CR, Jr., Edwards E, Wang RY (1994) Electrophysiological evidence for a functional interaction between 5-HT_{1A} and 5-HT_{2A} receptors in the rat medial prefrontal cortex: an iontophoretic study. *Synapse* 17: 173-181.
- Ashby CR, Jr., Jiang LH, Kasser RJ, Wang RY (1990) Electrophysiological characterization of 5-hydroxytryptamine₂ receptors in the rat medial prefrontal cortex. *J Pharmacol Exp Ther* 252: 171-178.
- Aston-Jones G, Ennis M, Pieribone VA, Nickell WT, Shipley MT (1986) The brain nucleus locus coeruleus: restricted afferent control of a broad efferent network. *Science* 234: 734-737.
- Auclair A, Blanc G, Glowinski J, Tassin JP (2004) Role of serotonin(2A) receptors in the D-amphetamine-induced release of dopamine: comparison with previous data on alpha 1b-adrenergic receptors. *Journal of Neurochemistry* 91: 318-326.
- Audet MA, Descarries L, Doucet G (1989) Quantified regional and laminar distribution of the serotonin innervation in the anterior half of adult rat cerebral cortex. *J Chem Neuroanat* 2: 29-44.
- Azmitia EC, Gannon PJ, Kheck NM, WhitakerAzmitia PM (1996) Cellular localization of the 5-HT_{1A} receptor in primate brain neurons and glial cells. *Neuropsychopharmacology* 14: 35-46.
- Azmitia EC, Segal M (1978) An autoradiographic analysis of the differential ascending projections of the dorsal and median raphe nuclei in the rat. *J Comp Neurol* 179: 641-667.
- Aznar S, Qian ZX, Shah R, Rahbek B, Knudsen GM (2003) The 5-HT_{1A} serotonin receptor is located on calbindin- and parvalbumin-containing neurons in the rat brain. *Brain Research* 959: 58-67.
- Baldessarini RJ, Huston-Lyons D, Campbell A, Marsh E, Cohen BM (1992) Do central antiadrenergic actions contribute to the atypical properties of clozapine? *Br J Psychiatry Suppl* 12-16.
- Bartrup JT, Newberry NR (1994) 5-HT_{2A} receptor-mediated outward current in C6 glioma cells is mimicked by intracellular IP₃ release. *Neuroreport* 5: 1245-1248.
- Beaulieu C (1993) Numerical data on neocortical neurons in adult rat, with special reference to the GABA population. *Brain Res* 609: 284-292.
- Benes FM, Berretta S (2001) GABAergic interneurons: implications for understanding schizophrenia and bipolar disorder. *Neuropsychopharmacology* 25: 1-27.
- Benes FM, Vincent SL, Molloy R (1993) Dopamine-Immunoreactive Axon Varicosities Form Nonrandom Contacts with Gaba-Immunoreactive Neurons of Rat Medial Prefrontal Cortex. *Synapse* 15: 285-295.

- Berendse HW, Groenewegen HJ (1991) Restricted cortical termination fields of the midline and intralaminar thalamic nuclei in the rat. *Neuroscience* 42: 73-102.
- Berg KA, Maayani S, Goldfarb J, Scaramellini C, Leff P, Clarke WP (1998) Effector pathway-dependent relative efficacy at serotonin type 2A and 2C receptors: Evidence for agonist-directed trafficking of receptor stimulus. *Molecular Pharmacology* 54: 94-104.
- Berger B, Gaspar P, Verney C (1991) Dopaminergic innervation of the cerebral cortex: unexpected differences between rodents and primates. *Trends Neurosci* 14: 21-27.
- Birnbaum S, Gobeske KT, Auerbach J, Taylor JR, Arnsten AF (1999) A role for norepinephrine in stress-induced cognitive deficits: alpha-1-adrenoceptor mediation in the prefrontal cortex. *Biol Psychiatry* 46: 1266-1274.
- Birnbaum SG, Yuan PX, Wang M, Vijayraghavan S, Bloom AK, Davis DJ, Gobeske KT, Sweatt JD, Manji HK, Arnsten AF (2004) Protein kinase C overactivity impairs prefrontal cortical regulation of working memory. *Science* 306: 882-884.
- Björklund A, Lindvall O (1984) Dopamine-containing systems in the CNS. In: Björklund A, Hökfelt T, editors. *Handbook of chemical neuroanatomy, Vol 2: Classical transmitters in the CNS*. Amsterdam: Elsevier.
- Blue ME, Yagaloff KA, Mamounas LA, Hartig PR, Molliver ME (1988) Correspondence between 5-HT₂ receptors and serotonergic axons in rat neocortex. *Brain Res* 453: 315-328.
- Bortolozzi A, Amargos-Bosch M, Adell A, Diaz-Mataix L, Serrats J, Pons S, Artigas F (2003) In vivo modulation of 5-hydroxytryptamine release in mouse prefrontal cortex by local 5-HT_{2A} receptors: effect of antipsychotic drugs. *European Journal of Neuroscience* 18: 1235-1246.
- Bouthenet ML, Souil E, Martres MP, Sokoloff P, Giros B, Schwartz JC (1991) Localization of Dopamine-D3 Receptor Messenger-Rna in the Rat-Brain Using In situ Hybridization Histochemistry - Comparison with Dopamine-D2 Receptor Messenger-Rna. *Brain Research* 564: 203-219.
- Brodmann K (1909) *Vergleichende Lokalisationslehre der Grosshirnrinde in ihren Prinzipien dargestellt auf Grund des Zellenbaues*. Barth, Leipzig.
- Bromet EJ, Fennig S (1999) Epidemiology and natural history of schizophrenia. *Biological Psychiatry* 46: 871-881.
- Brown VJ, Bowman EM (2002) Rodent models of prefrontal cortical function. *Trends Neurosci* 25: 340-343.
- Brozoski TJ, Brown RM, Rosvold HE, Goldman PS (1979) Cognitive Deficit Caused by Regional Depletion of Dopamine in Prefrontal Cortex of Rhesus-Monkey. *Science* 205: 929-932.
- Bymaster FP, HemrickLuecke SK, Perry KW, Fuller RW (1996) Neurochemical evidence for antagonism by olanzapine of dopamine, serotonin, alpha(1)-adrenergic and muscarinic receptors in vivo in rats. *Psychopharmacology* 124: 87-94.
- Cannon TD, Rosso IM, Bearden CE, Sanchez LE, Hadley T (1999) A prospective cohort study of neurodevelopmental processes in the genesis and epigenesis of schizophrenia. *Development and Psychopathology* 11: 467-485.

- Carli M, Robbins TW, Evenden JL, Everitt BJ (1983) Effects of lesions to ascending noradrenergic neurones on performance of a 5-choice serial reaction task in rats; implications for theories of dorsal noradrenergic bundle function based on selective attention and arousal. *Behav Brain Res* 9: 361-380.
- Carlsson A (1988) The Current Status of the Dopamine Hypothesis of Schizophrenia. *Neuropsychopharmacology* 1: 179-186.
- Carlsson A, Waters N, Holm-Waters S, Tedroff J, Nilsson M, Carlsson ML (2001) Interactions between monoamines, glutamate, and GABA in schizophrenia: new evidence. *Annu Rev Pharmacol Toxicol* 41: 237-260.
- Carlsson A, Waters N, Waters S, Carlsson ML (2000) Network interactions in schizophrenia - therapeutic implications. *Brain Research Reviews* 31: 342-349.
- Carlsson M, Carlsson A (1990) Interactions between glutamatergic and monoaminergic systems within the basal ganglia--implications for schizophrenia and Parkinson's disease. *Trends Neurosci* 13: 272-276.
- Carr DB, Sesack SR (2000) Projections from the rat prefrontal cortex to the ventral tegmental area: target specificity in the synaptic associations with mesoaccumbens and mesocortical neurons. *J Neurosci* 20: 3864-3873.
- Celada P, Puig MV, Casanovas JM, Guillazo G, Artigas F (2001) Control of dorsal raphe serotonergic neurons by the medial prefrontal cortex: Involvement of serotonin-1A, GABA(A), and glutamate receptors. *J Neurosci* 21: 9917-9929.
- Ciliax BJ, Nash N, Heilman C, Sunahara R, Hartney A, Tiberi M, Rye DB, Caron MG, Niznik HB, Levey AI (2000) Dopamine D(5) receptor immunolocalization in rat and monkey brain. *Synapse* 37: 125-145.
- Claro E, Fain JN, Picatoste F (1993) Noradrenaline Stimulation Unbalances the Phosphoinositide Cycle in Rat Cerebral Cortical Slices. *Journal of Neurochemistry* 60: 2078-2086.
- Cole BJ, Robbins TW (1992) Forebrain norepinephrine: role in controlled information processing in the rat. *Neuropsychopharmacology* 7: 129-142.
- Conde F, Audinat E, Maire-Lepoivre E, Crepel F (1990) Afferent connections of the medial frontal cortex of the rat. A study using retrograde transport of fluorescent dyes. I. Thalamic afferents. *Brain Res Bull* 24: 341-354.
- Conde F, Maire-Lepoivre E, Audinat E, Crepel F (1995) Afferent connections of the medial frontal cortex of the rat. II. Cortical and subcortical afferents. *J Comp Neurol* 352: 567-593.
- Cornea-Hebert V, Riad M, Wu C, Singh SK, Descarries L (1999) Cellular and subcellular distribution of the serotonin 5-HT_{2A} receptor in the central nervous system of adult rat. *Journal of Comparative Neurology* 409: 187-209.
- Creese I, Burt DR, Snyder SH (1976) Dopamine Receptor-Binding Predicts Clinical and Pharmacological Potencies of Anti-Schizophrenic Drugs. *Science* 192: 481-483.

- Dalley JW, Cardinal RN, Robbins TW (2004) Prefrontal executive and cognitive functions in rodents: neural and neurochemical substrates. *Neurosci Biobehav Rev* 28: 771-784.
- Daniel DG, Zimbroff DL, Potkin SG, Reeves KR, Harrigan EP, Lakshminarayanan M (1999) Ziprasidone 80 mg/day and 160 mg/day in the acute exacerbation of schizophrenia and schizoaffective disorder: a 6-week placebo-controlled trial. Ziprasidone Study Group. *Neuropsychopharmacology* 20: 491-505.
- Davidoff SA, Benes FM (1998) High-resolution scatchard analysis shows D1 receptor binding on pyramidal and nonpyramidal neurons. *Synapse* 28: 83-90.
- Davis KL, Kahn RS, Ko G, Davidson M (1991) Dopamine in schizophrenia: a review and reconceptualization. *Am J Psychiatry* 148: 1474-1486.
- Day HE, Campeau S, Watson SJ, Jr., Akil H (1997) Distribution of alpha 1a-, alpha 1b- and alpha 1d-adrenergic receptor mRNA in the rat brain and spinal cord. *J Chem Neuroanat* 13: 115-139.
- de Lima AD, Morrison JH (1989) Ultrastructural analysis of somatostatin-immunoreactive neurons and synapses in the temporal and occipital cortex of the macaque monkey. *J Comp Neurol* 283: 212-227.
- Defagot MC, Antonelli MC (1997) Autoradiographic localization of the putative D4 dopamine receptor in rat brain. *Neurochemical Research* 22: 401-407.
- Defagot MC, Malchiodi EL, Villar MJ, Antonelli MC (1997) Distribution of D4 dopamine receptor in rat brain with sequence-specific antibodies. *Molecular Brain Research* 45: 1-12.
- DeFelipe J (2002) Cortical interneurons: from Cajal to 2001. *Prog Brain Res* 136: 215-238.
- DeFelipe J, Arellano JI, Gomez A, Azmitia EC, Munoz A (2001) Pyramidal cell axons show a local specialization for GABA and 5-HT inputs in monkey and human cerebral cortex. *Journal of Comparative Neurology* 433: 148-155.
- DeFelipe J, Hendry SH, Hashikawa T, Molinari M, Jones EG (1990) A microcolumnar structure of monkey cerebral cortex revealed by immunocytochemical studies of double bouquet cell axons. *Neuroscience* 37: 655-673.
- DeFelipe J, Hendry SH, Jones EG (1989) Synapses of double bouquet cells in monkey cerebral cortex visualized by calbindin immunoreactivity. *Brain Res* 503: 49-54.
- Del Arco A, Mora F (2001) Dopamine release in the prefrontal cortex during stress is reduced by the local activation of glutamate receptors. *Brain Res Bull* 56: 125-130.
- del Rio MR, DeFelipe J (1997) Synaptic connections of calretinin-immunoreactive neurons in the human neocortex. *J Neurosci* 17: 5143-5154.
- Delatour B, Gisquet-Verrier P (1999b) Lesions of the prelimbic-infralimbic cortices in rats do not disrupt response selection processes but induce delay-dependent deficits: evidence for a role in working memory? *Behav Neurosci* 113: 941-955.

- Delatour B, Gisquet-Verrier P (1999a) Lesions of the prelimbic-infralimbic cortices in rats do not disrupt response selection processes but induce delay-dependent deficits: evidence for a role in working memory? *Behav Neurosci* 113: 941-955.
- Delatour B, Gisquet-Verrier P (2000) Functional role of rat prelimbic-infralimbic cortices in spatial memory: evidence for their involvement in attention and behavioural flexibility. *Behav Brain Res* 109: 113-128.
- DeLeon A, Patel NC, Crismon ML (2004) Aripiprazole: a comprehensive review of its pharmacology, clinical efficacy, and tolerability. *Clin Ther* 26: 649-666.
- Derkach V, Surprenant A, North RA (1989) 5-HT₃ Receptors Are Membrane Ion Channels. *Nature* 339: 706-709.
- Deutch AY, Moghaddam B, Innis RB, Krystal JH, Aghajanian GK, Bunney BS, Charney DS (1991) Mechanisms of action of atypical antipsychotic drugs. Implications for novel therapeutic strategies for schizophrenia. *Schizophr Res* 4: 121-156.
- Diaz J, Levesque D, Lammers CH, Griffon N, Martres MP, Schwartz JC, Sokoloff P (1995) Phenotypical characterization of neurons expressing the dopamine D₃ receptor in the rat brain. *Neuroscience* 65: 731-745.
- Diaz J, Pilon C, Le Foll B, Gros C, Triller A, Schwartz JC, Sokoloff P (2000) Dopamine D-3 receptors expressed by all mesencephalic dopamine neurons. *Journal of Neuroscience* 20: 8677-8684.
- Domyancic AV, Morilak DA (1997) Distribution of alpha_{1A} adrenergic receptor mRNA in the rat brain visualized by in situ hybridization. *J Comp Neurol* 386: 358-378.
- Duncan G, Miyamoto S, Gu H, Lieberman J, Koller B, Snouwaert J (2002) Alterations in regional brain metabolism in genetic and pharmacological models of reduced NMDA receptor function. *Brain Res* 951: 166-176.
- Dursun SM, Handley SL (1996) Similarities in the pharmacology of spontaneous and DOI-induced head-shakes suggest 5HT_{2A} receptors are active under physiological conditions. *Psychopharmacology (Berl)* 128: 198-205.
- Fallon JH, Opole IO, Potkin SG (2003) The neuroanatomy of schizophrenia: circuitry and neurotransmitter systems. *Clinical Neuroscience Research* 3: 77-107.
- Ferezou I, Cauli B, Hill EL, Rossier J, Hamel E, Lambolez B (2002) 5-HT₃ receptors mediate serotonergic fast synaptic excitation of neocortical vasoactive intestinal peptide/cholecystokinin interneurons. *J Neurosci* 22: 7389-7397.
- Finlay JM, Zigmond MJ, Abercrombie ED (1995) Increased dopamine and norepinephrine release in medial prefrontal cortex induced by acute and chronic stress: effects of diazepam. *Neuroscience* 64: 619-628.
- Fleischhacker WW (1995) New Drugs for the Treatment of Schizophrenic-Patients. *Acta Psychiatrica Scandinavica* 91: 24-30.
- Floresco SB, Magyar O (2006) Mesocortical dopamine modulation of executive functions: beyond working memory. *Psychopharmacology (Berl)* 188: 567-585.

- Floresco SB, Seamans JK, Phillips AG (1997) Selective roles for hippocampal, prefrontal cortical, and ventral striatal circuits in radial-arm maze tasks with or without a delay. *J Neurosci* 17: 1880-1890.
- Freedman R (2003) Schizophrenia. *New England Journal of Medicine* 349: 1738-1749.
- Freneau RT, Duncan GE, Fornaretto MG, Dearry A, Gingrich JA, Breese GR, Caron MG (1991) Localization of D1 Dopamine Receptor Messenger-Rna in Brain Supports A Role in Cognitive, Affective, and Neuroendocrine Aspects of Dopaminergic Neurotransmission. *Proceedings of the National Academy of Sciences of the United States of America* 88: 3772-3776.
- Freund TF, Buzsaki G (1996) Interneurons of the hippocampus. *Hippocampus* 6: 347-470.
- Fuster JM (1997) *The Prefrontal Cortex. Anatomy, Physiology, and Neuropsychology of the Frontal Lobe.* Philadelphia-New York: Lipincott-Raven.
- Fuster JM (1973) Unit activity in prefrontal cortex during delayed-response performance: neuronal correlates of transient memory. *J Neurophysiol* 36: 61-78.
- Gaspar P, Bloch B, Le Moine C (1995) D1 and D2 receptor gene expression in the rat frontal cortex: cellular localization in different classes of efferent neurons. *Eur J Neurosci* 7: 1050-1063.
- Gehlert DR, Gackenheim SL, Wong DT, Robertson DW (1991) Localization of 5-Ht3 Receptors in the Rat-Brain Using [H-3] Ly278584. *Brain Research* 553: 149-154.
- Gellman RL, Aghajanian GK (1993) Pyramidal cells in piriform cortex receive a convergence of inputs from monoamine activated GABAergic interneurons. *Brain Res* 600: 63-73.
- Gibson JR, Beierlein M, Connors BW (1999) Two networks of electrically coupled inhibitory neurons in neocortex. *Nature* 402: 75-79.
- Gilmore JH, Jarskog LF (1997) Exposure to infection and brain development: cytokines in the pathogenesis of schizophrenia. *Schizophr Res* 24: 365-367.
- Gisquet-Verrier P, Winocur G, Delatour B (2000) Functional dissociation between dorsal and ventral regions of the medial prefrontal cortex in rats. *Psychobiology* 28: 248-260.
- Goldman-Rakic PS (1995) Cellular basis of working memory. *Neuron* 14: 477-485.
- Goldman-Rakic PS, Leranth C, Williams SM, Mons N, Geffard M (1989) Dopamine synaptic complex with pyramidal neurons in primate cerebral cortex. *Proc Natl Acad Sci U S A* 86: 9015-9019.
- Goldman-Rakic PS, Muly EC, III, Williams GV (2000) D(1) receptors in prefrontal cells and circuits. *Brain Res Brain Res Rev* 31: 295-301.
- Goldstein LE, Rasmusson AM, Bunney BS, Roth RH (1996) Role of the amygdala in the coordination of behavioral, neuroendocrine, and prefrontal cortical monoamine responses to psychological stress in the rat. *J Neurosci* 16: 4787-4798.
- Goldstein M, Deutch AY (1992) Dopaminergic mechanisms in the pathogenesis of schizophrenia. *FASEB J* 6: 2413-2421.

- Gorelova N, Seamans JK, Yang CR (2002) Mechanisms of dopamine activation of fast-spiking interneurons that exert inhibition in rat prefrontal cortex. *J Neurophysiol* 88: 3150-3166.
- Gottesman II (1991) *Schizophrenia Genesis: The Origins of Madness*. New York: Freeman.
- Gottesman II, Shields J (1982) *Schizophrenia: The Epigenetic Puzzle*. Cambridge, UK: Cambridge University Press.
- Grady RM, Zhou H, Cunningham JM, Henry MD, Campbell KP, Sanes JR (2000) Maturation and maintenance of the neuromuscular synapse: genetic evidence for roles of the dystrophin-glycoprotein complex. *Neuron* 25: 279-293.
- Granon S, Passetti F, Thomas KL, Dalley JW, Everitt BJ, Robbins TW (2000) Enhanced and impaired attentional performance after infusion of D1 dopaminergic receptor agents into rat prefrontal cortex. *J Neurosci* 20: 1208-1215.
- Greenshaw AJ (1993) Behavioural pharmacology of 5-HT₃ receptor antagonists: a critical update on therapeutic potential. *Trends Pharmacol Sci* 14: 265-270.
- Groenewegen HJ, Uylings HB (2000) The prefrontal cortex and the integration of sensory, limbic and autonomic information. *Prog Brain Res* 126: 3-28.
- Groenewegen HJ, Wright CI, Uylings HB (1997) The anatomical relationships of the prefrontal cortex with limbic structures and the basal ganglia. *J Psychopharmacol* 11: 99-106.
- Guidotti A, Auta J, Davis JM, Giorgi-Gerevini V, Dwivedi Y, Grayson DR, Impagnatiello F, Pandey G, Pesold C, Sharma R, Uzunov D, Costa E (2000) Decrease in reelin and glutamic acid decarboxylase67 (GAD67) expression in schizophrenia and bipolar disorder: a postmortem brain study. *Arch Gen Psychiatry* 57: 1061-1069.
- Hague C, Lee SE, Chen ZJ, Prinster SC, Hall RA, Minneman KP (2006) Heterodimers of alpha1B and alpha1D-adrenergic receptors form a single functional entity. *Faseb Journal* 20: A253-A254.
- Hajos M, Richards CD, Szekely AD, Sharp T (1998) An electrophysiological and neuroanatomical study of the medial prefrontal cortical projection to the midbrain raphe nuclei in the rat. *Neuroscience* 87: 95-108.
- Harder JA, Ridley RM (2000) The 5-HT_{1A} antagonist, WAY 100 635, alleviates cognitive impairments induced by dizocilpine (MK-801) in monkeys. *Neuropharmacology* 39: 547-552.
- Harrison PJ (1999) Neurochemical alterations in schizophrenia affecting the putative receptor targets of atypical antipsychotics - Focus on dopamine (D-1, D-3, D-4) and 5-HT_{2a} receptors. *British Journal of Psychiatry* 174: 12-22.
- Hashimoto T, Bergen SE, Nguyen QL, Xu B, Monteggia LM, Pierri JN, Sun Z, Sampson AR, Lewis DA (2005) Relationship of brain-derived neurotrophic factor and its receptor TrkB to altered inhibitory prefrontal circuitry in schizophrenia. *J Neurosci* 25: 372-383.
- Hashimoto T, Volk DW, Eggan SM, Mirnics K, Pierri JN, Sun Z, Sampson AR, Lewis DA (2003) Gene expression deficits in a subclass of GABA neurons in the prefrontal cortex of subjects with schizophrenia. *J Neurosci* 23: 6315-6326.

- Heidbreder CA, Groenewegen HJ (2003) The medial prefrontal cortex in the rat: evidence for a dorso-ventral distinction based upon functional and anatomical characteristics. *Neurosci Biobehav Rev* 27: 555-579.
- Heisler LK, Chu HM, Brennan TJ, Danao JA, Bajwa P, Parsons LH, Tecott LH (1998) Elevated anxiety and antidepressant-like responses in serotonin 5-HT_{1A} receptor mutant mice. *Proceedings of the National Academy of Sciences of the United States of America* 95: 15049-15054.
- Herrick-Davis K, Grinde E, Harrigan TJ, Mazurkiewicz JE (2005) Inhibition of serotonin 5-hydroxytryptamine_{2C} receptor function through heterodimerization - Receptor dimers bind two molecules of ligand and one G-protein. *Journal of Biological Chemistry* 280: 40144-40151.
- Huang Q, Zhou D, Chase K, Gusella JF, Aronin N, DiFiglia M (1992) Immunohistochemical localization of the D₁ dopamine receptor in rat brain reveals its axonal transport, pre- and postsynaptic localization, and prevalence in the basal ganglia, limbic system, and thalamic reticular nucleus. *Proc Natl Acad Sci U S A* 89: 11988-11992.
- Ichikawa J, Ishii H, Bonaccorso S, Fowler WL, O'Laughlin IA, Meltzer HY (2001) 5-HT_{2A} and D₂ receptor blockade increases cortical DA release via 5-HT_{1A} receptor activation: a possible mechanism of atypical antipsychotic-induced cortical dopamine release. *J Neurochem* 76: 1521-1531.
- Jakab RL, Goldman-Rakic PS (1998) 5-Hydroxytryptamine_{2A} serotonin receptors in the primate cerebral cortex: possible site of action of hallucinogenic and antipsychotic drugs in pyramidal cell apical dendrites. *Proc Natl Acad Sci U S A* 95: 735-740.
- Jakab RL, Goldman-Rakic PS (2000) Segregation of serotonin 5-HT_{2A} and 5-HT₃ receptors in inhibitory circuits of the primate cerebral cortex. *J Comp Neurol* 417: 337-348.
- Jarskog LF, Gilmore JH, Selinger ES, Lieberman JA (2000) Cortical bcl-2 protein expression and apoptotic regulation in schizophrenia. *Biol Psychiatry* 48: 641-650.
- Jaskiw GE, Weinberger DR (1987) The prefrontal cortex-accumbens circuit: Who's in charge? *Behav Brain Sci* 10: 217-218.
- Javitt DC, Zukin SR (1991) Recent advances in the phencyclidine model of schizophrenia. *Am J Psychiatry* 148: 1301-1308.
- Jones EG (2000) Microcolumns in the cerebral cortex. *Proceedings of the National Academy of Sciences of the United States of America* 97: 5019-5021.
- Jones LS, Miller G, Gauger LL, Davis JN (1985) Regional distribution of rat brain alpha 1-adrenergic receptors: correlation between (125I)-heat membrane binding and in vitro autoradiography. *Life Sci* 36: 45-51.
- Jones P (1997) The early origins of schizophrenia. *Br Med Bull* 53: 135-155.
- Kandel E, Schwartz J, Jessell T (2000) *Principles of Neural Science*. McGraw-Hill.
- Kapur S, Seeman P (2002a) Atypical antipsychotics, cortical D₂ receptors and sensitivity to endogenous dopamine. *Br J Psychiatry* 180: 465-466.

- Kapur S, Seeman P (2002b) NMDA receptor antagonists ketamine and PCP have direct effects on the dopamine D(2) and serotonin 5-HT(2) receptors-implications for models of schizophrenia. *Mol Psychiatry* 7: 837-844.
- Kawa K (1994) Distribution and Functional-Properties of 5-Ht₃ Receptors in the Rat Hippocampal Dentate Gyrus - A Patch-Clamp Study. *Journal of Neurophysiology* 71: 1935-1947.
- Kawaguchi Y, Kubota Y (1996) Physiological and morphological identification of somatostatin- or vasoactive intestinal polypeptide-containing cells among GABAergic cell subtypes in rat frontal cortex. *J Neurosci* 16: 2701-2715.
- Kesner RP (2000) Subregional analysis of mnemonic functions of the prefrontal cortex in the rat. *Psychobiology* 28: 219-228.
- Khan ZU, Gutierrez A, Martin R, Penafiel A, Rivera A, de la Calle A (1998) Differential regional and cellular distribution of dopamine D₂-like receptors: An immunocytochemical study of subtype-specific antibodies in rat and human brain. *Journal of Comparative Neurology* 402: 353-371.
- Khan ZU, Gutierrez A, Martin R, Penafiel A, Rivera A, de la Calle A (2000) Dopamine D₅ receptors of rat and human brain. *Neuroscience* 100: 689-699.
- Kia HK, Brisorgueil MJ, Hamon M, Calas A, Verge D (1996a) Ultrastructural localization of 5-hydroxytryptamine_{1A} receptors in the rat brain. *J Neurosci Res* 46: 697-708.
- Kia HK, Miquel MC, Brisorgueil MJ, Daval G, Riad M, El Mestikawy S, Hamon M, Verge D (1996b) Immunocytochemical localization of serotonin_{1A} receptors in the rat central nervous system. *J Comp Neurol* 365: 289-305.
- Knable MB, Barci BM, Bartko JJ, Webster MJ, Torrey EF (2002) Molecular abnormalities in the major psychiatric illnesses: Classification and Regression Tree (CRT) analysis of post-mortem prefrontal markers. *Mol Psychiatry* 7: 392-404.
- Kosofsky BE, Molliver ME (1987) The serotonergic innervation of cerebral cortex: different classes of axon terminals arise from dorsal and median raphe nuclei. *Synapse* 1: 153-168.
- Krimer LS, Jakab RL, Goldman-Rakic PS (1997) Quantitative three-dimensional analysis of the catecholaminergic innervation of identified neurons in the macaque prefrontal cortex. *J Neurosci* 17: 7450-7461.
- Kroeze WK, Roth BL (1998) The molecular biology of serotonin receptors: therapeutic implications for the interface of mood and psychosis. *Biol Psychiatry* 44: 1128-1142.
- Krystal JH, Karper LP, Seibyl JP, Freeman GK, Delaney R, Bremner JD, Heninger GR, Bowers MB, Charney DS (1994) Subanesthetic Effects of the Noncompetitive Nmda Antagonist, Ketamine, in Humans - Psychotomimetic, Perceptual, Cognitive, and Neuroendocrine Responses. *Archives of General Psychiatry* 51: 199-214.
- Kuhar MJ, Ritz MC, Boja JW (1991) The dopamine hypothesis of the reinforcing properties of cocaine. *Trends Neurosci* 14: 299-302.
- Kuroda M, Yokofujita J, Murakami K (1998) An ultrastructural study of the neural circuit between the prefrontal cortex and the mediodorsal nucleus of the thalamus. *Prog Neurobiol* 54: 417-458.

- Lambe EK, Goldman-Rakic PS, Aghajanian GK (2000) Serotonin induces EPSCs preferentially in layer V pyramidal neurons of the frontal cortex in the rat. *Cereb Cortex* 10: 974-980.
- Landwehrmeyer B, Mengod G, Palacios JM (1993) Dopamine-D3 Receptor Messenger-Rna and Binding-Sites in Human Brain. *Molecular Brain Research* 18: 187-192.
- Le Moal M, Simon H (1991) Mesocorticolimbic dopaminergic network: functional and regulatory roles. *Physiol Rev* 71: 155-234.
- Le Moine C, Gaspar P (1998) Subpopulations of cortical GABAergic interneurons differ by their expression of D1 and D2 dopamine receptor subtypes. *Brain Res Mol Brain Res* 58: 231-236.
- Lejeune F, Millan MJ (1998) Induction of burst firing in ventral tegmental area dopaminergic neurons by activation of serotonin (5-HT)1A receptors: WAY 100,635-reversible actions of the highly selective ligands, flesinoxan and S 15535. *Synapse* 30: 172-180.
- Levesque D, Diaz J, Pilon C, Martres MP, Giros B, Souil E, Schott D, Morgat JL, Schwartz JC, Sokoloff P (1992) Identification, Characterization, and Localization of the Dopamine-D3 Receptor in Rat-Brain Using 7-[H-3]Hydroxy-N,N-Di-Normal-Propyl-2-Aminotetralin. *Proceedings of the National Academy of Sciences of the United States of America* 89: 8155-8159.
- Lewis DA (1997) Development of the prefrontal cortex during adolescence: Insights into vulnerable neural circuits in schizophrenia. *Neuropsychopharmacology* 16: 385-398.
- Lewis DA, Hashimoto T, Volk DW (2005) Cortical inhibitory neurons and schizophrenia. *Nat Rev Neurosci* 6: 312-324.
- Lewis DA, Morrison JH (1989) Noradrenergic innervation of monkey prefrontal cortex: a dopamine-beta-hydroxylase immunohistochemical study. *J Comp Neurol* 282: 317-330.
- Lidow MS, Goldmanrakic PS, Gallager DW, Rakic P (1991) Distribution of Dopaminergic Receptors in the Primate Cerebral-Cortex - Quantitative Autoradiographic Analysis Using [H-3] Raclopride, [H-3] Spiperone and [H-3] Sch23390. *Neuroscience* 40: 657-671.
- Lidow MS, Wang F, Cao Y, Goldman-Rakic PS (1998) Layer V neurons bear the majority of mRNAs encoding the five distinct dopamine receptor subtypes in the primate prefrontal cortex. *Synapse* 28: 10-20.
- Lieberman JA, Mailman RB, Duncan G, Sikich L, Chakos M, Nichols DE, Kraus JE (1998) Serotonergic basis of antipsychotic drug effects in schizophrenia. *Biol Psychiatry* 44: 1099-1117.
- Lipska BK (2004) Using animal models to test a neurodevelopmental hypothesis of schizophrenia. *Journal of Psychiatry & Neuroscience* 29: 282-286.
- Lipska BK, Weinberger DR (1993) Delayed effects of neonatal hippocampal damage on haloperidol-induced catalepsy and apomorphine-induced stereotypic behaviors in the rat. *Brain Res Dev Brain Res* 75: 213-222.
- Lipska BK, Weinberger DR (1995) Genetic variation in vulnerability to the behavioral effects of neonatal hippocampal damage in rats. *Proc Natl Acad Sci U S A* 92: 8906-8910.

- Lorente de Nó R (1938) Architectonics and structure of the cerebral cortex. In *Physiology of the Nervous System* (edited by FULTON, J. F.) pp. 291-330. New York: Oxford University Press.
- Malhotra AK, Adler CM, Kennison SD, Elman I, Pickar D, Breier A (1997) Clozapine blunts N-methyl-D-aspartate antagonist-induced psychosis: A study with ketamine. *Biological Psychiatry* 42: 664-668.
- Mansour A, Meadorwoodruff JH, Bunzow JR, Civelli O, Akil H, Watson SJ (1990) Localization of Dopamine-D2 Receptor Messenger-Rna and D1 and D2 Receptor-Binding in the Rat-Brain and Pituitary - An Insitu Hybridization-Receptor Autoradiographic Analysis. *Journal of Neuroscience* 10: 2587-2600.
- Mansour A, Meadorwoodruff JH, Zhou Q, Civelli O, Akil H, Watson SJ (1992b) A Comparison of D1-Receptor Binding and Messenger-Rna in Rat-Brain Using Receptor Autoradiographic and Insitu Hybridization Techniques. *Neuroscience* 46: 959-971.
- Mansour A, Meadorwoodruff JH, Zhou Q, Civelli O, Akil H, Watson SJ (1992c) A Comparison of D1-Receptor Binding and Messenger-Rna in Rat-Brain Using Receptor Autoradiographic and Insitu Hybridization Techniques. *Neuroscience* 46: 959-971.
- Mansour A, Meadorwoodruff JH, Zhou Q, Civelli O, Akil H, Watson SJ (1992a) A Comparison of D1-Receptor Binding and Messenger-Rna in Rat-Brain Using Receptor Autoradiographic and Insitu Hybridization Techniques. *Neuroscience* 46: 959-971.
- Mansour A, Meadorwoodruff JH, Zhou QY, Civelli O, Akil H, Watson SJ (1991) A Comparison of D1-Receptor Binding and Messenger-Rna in Rat-Brain Using Receptor Autoradiographic and Insitu Hybridization Techniques. *Neuroscience* 45: 359-371.
- Mao ZM, Arnsten AF, Li BM (1999) Local infusion of an alpha-1 adrenergic agonist into the prefrontal cortex impairs spatial working memory performance in monkeys. *Biol Psychiatry* 46: 1259-1265.
- Marcelis M, van Os J, Sham P, Jones P, Gilvarry C, Cannon M, McKenzie K, Murray R (1998) Obstetric complications and familial morbid risk of psychiatric disorders. *American Journal of Medical Genetics* 81: 29-36.
- Marek GJ, Aghajanian GK (1996) LSD and the phenethylamine hallucinogen DOI are potent partial agonists at 5-HT_{2A} receptors on interneurons in rat piriform cortex. *J Pharmacol Exp Ther* 278: 1373-1382.
- Marek GJ, Aghajanian GK (1999) 5-HT_{2A} receptor or alpha1-adrenoceptor activation induces excitatory postsynaptic currents in layer V pyramidal cells of the medial prefrontal cortex. *Eur J Pharmacol* 367: 197-206.
- Marenco S, Weinberger DR (2000) The neurodevelopmental hypothesis of schizophrenia: Following a trail of evidence from cradle to grave. *Development and Psychopathology* 12: 501-527.
- Maricq AV, Peterson AS, Brake AJ, Myers RM, Julius D (1991) Primary Structure and Functional Expression of the 5HT₃ Receptor, A Serotonin-Gated Ion Channel. *Science* 254: 432-437.
- Martin-Ruiz R, Puig MV, Celada P, Shapiro DA, Roth BL, Mengod G, Artigas F (2001) Control of serotonergic function in medial prefrontal cortex by serotonin-2A receptors through a glutamate-dependent mechanism. *Journal of Neuroscience* 21: 9856-9866.

- Marx CE, Jarskog LF, Lauder JM, Lieberman JA, Gilmore JH (2001) Cytokine effects on cortical neuron MAP-2 immunoreactivity: implications for schizophrenia. *Biol Psychiatry* 50: 743-749.
- Matsuoka T, Nishizaki T, Ikeuchi Y, Okada Y, Sumino K (1997) Differential effects of the serotonin receptors on cultured rat cerebral cortical neurons. *Cellular and Molecular Life Sciences* 53: 233-236.
- Mattay VS, Tessitore A, Callicott JH, Bertolino A, Goldberg TE, Chase TN, Hyde TM, Weinberger DR (2002) Dopaminergic modulation of cortical function in patients with Parkinson's disease. *Ann Neurol* 51: 156-164.
- McCormick DA, Connors BW, Lighthall JW, Prince DA (1985) Comparative electrophysiology of pyramidal and sparsely spiny stellate neurons of the neocortex. *J Neurophysiol* 54: 782-806.
- McCune SK, Voigt MM, Hill JM (1993) Expression of multiple alpha adrenergic receptor subtype messenger RNAs in the adult rat brain. *Neuroscience* 57: 143-151.
- McQuade R, Sharp T (1997) Functional mapping of dorsal and median raphe 5-hydroxytryptamine pathways in forebrain of the rat using microdialysis. *J Neurochem* 69: 791-796.
- Meadorwoodruff JH, Mansour A, Grandy DK, Damask SP, Civelli O, Watson SJ (1992) Distribution of D(5) Dopamine Receptor Messenger-Rna in Rat-Brain. *Neuroscience Letters* 145: 209-212.
- Mehta MA, Riedel WJ (2006) Dopaminergic enhancement of cognitive function. *Curr Pharm Des* 12: 2487-2500.
- Mello NK, Negus SS (2001) Effects of indatraline and buprenorphine on self-administration of speedball combinations of cocaine and heroin by rhesus monkeys. *Neuropsychopharmacology* 25: 104-117.
- Meltzer CC, Smith G, DeKosky ST, Pollock BG, Mathis CA, Moore RY, Kupfer DJ, Reynolds CF (1998) Serotonin in aging, late-life depression, and Alzheimer's disease: The emerging role of functional imaging. *Neuropsychopharmacology* 18: 407-430.
- Meltzer HY (1999) The role of serotonin in antipsychotic drug action. *Neuropsychopharmacology* 21: 106S-115S.
- Mengod G, Vilaro MT, Niznik HB, Sunahara RK, Seeman P, Odowd BF, Palacios JM (1991) Visualization of A Dopamine-D1 Receptor Messenger-Rna in Human and Rat-Brain. *Molecular Brain Research* 10: 185-191.
- Mengod G, Vilaro MT, Landwehrmeyer GB, Martinezmir MI, Niznik HB, Sunahara RK, Seeman P, Odowd BF, Probst A, Palacios JM (1992) Visualization of Dopamine D1, D2 and D3 Receptor Messenger-Rnas in Human and Rat-Brain. *Neurochemistry International* 20: S33-S43.
- Michel MC, Erdbrugger W, Bischoff A, Brodde OE (1993) Differential Phospholipase-A(2) and Phospholipase-C Coupling for Rat Renal Angiotensin-li, Neuropeptide-y and Alpha(1)-Adrenergic Receptors. *Faseb Journal* 7: A467.
- Milad MR, Quirk GJ (2002) Neurons in medial prefrontal cortex signal memory for fear extinction. *Nature* 420: 70-74.

- Millan MJ (2000) Improving the treatment of schizophrenia: focus on serotonin (5-HT)(1A) receptors. *J Pharmacol Exp Ther* 295: 853-861.
- Millar JK, Wilson-Annan JC, Anderson S, Christie S, Taylor MS, Semple CA, Devon RS, Clair DM, Muir WJ, Blackwood DH, Porteous DJ (2000) Disruption of two novel genes by a translocation co-segregating with schizophrenia. *Hum Mol Genet* 9: 1415-1423.
- Miller EK, Cohen JD (2001) An integrative theory of prefrontal cortex function. *Annu Rev Neurosci* 24: 167-202.
- Mirnics K, Middleton FA, Marquez A, Lewis DA, Levitt P (2000) Molecular characterization of schizophrenia viewed by microarray analysis of gene expression in prefrontal cortex. *Neuron* 28: 53-67.
- Misane I, Ogren SO (2003) Selective 5-HT_{1A} antagonists WAY 100635 and NAD-299 attenuate the impairment of passive avoidance caused by scopolamine in the rat. *Neuropsychopharmacology* 28: 253-264.
- Miyamoto S, Duncan GE, Mailman RB (2000) Developing novel antipsychotic drugs: strategies and goals. *Curr Opin CPNS Invest Drugs* 2: 25-39.
- Miyamoto S, Lieberman JA, Fleischhacker WW, et al. (2003) Antipsychotic drugs. In *Psychiatry*, ed. A Tasman, J Kay, JA Lieberman, pp. 1928–1964. Chichester, UK: JohnWiley & Sons.
- Miyamoto S, Duncan GE, Marx CE, Lieberman JA (2005) Treatments for schizophrenia: a critical review of pharmacology and mechanisms of action of antipsychotic drugs. *Molecular Psychiatry* 10: 79-104.
- Miyamoto S, LaMantia AS, Duncan GE, Sullivan P, Gilmore JH, Lieberman JA (2003) Recent advances in the neurobiology of schizophrenia. *Mol Interv* 3: 27-39.
- Moghaddam B (2003) Bringing order to the glutamate chaos in schizophrenia. *Neuron* 40: 881-884.
- Moghaddam B, Bunney BS (1989) Differential effect of cocaine on extracellular dopamine levels in rat medial prefrontal cortex and nucleus accumbens: comparison to amphetamine. *Synapse* 4: 156-161.
- Mohn AR, Gainetdinov RR, Caron MG, Koller BH (1999) Mice with reduced NMDA receptor expression display behaviors related to schizophrenia. *Cell* 98: 427-436.
- Molinoff PB (1984) Alpha- and beta-adrenergic receptor subtypes properties, distribution and regulation. *Drugs* 28 Suppl 2: 1-15.
- Morales M, Battenberg E, deLecea L, Bloom FE (1996) The type 3 serotonin receptor is expressed in a subpopulation of GABAergic neurons in the rat neocortex and hippocampus. *Brain Research* 731: 199-202.
- Morales M, Bloom FE (1997) The 5-HT₃ receptor is present in different subpopulations of GABAergic neurons in the rat telencephalon. *J Neurosci* 17: 3157-3167.

- Morgan MA, Schulkin J, LeDoux JE (2003) Ventral medial prefrontal cortex and emotional perseveration: the memory for prior extinction training. *Behav Brain Res* 146: 121-130.
- Morrison JH, Molliver ME, Grzanna R (1979) Noradrenergic Innervation of Cerebral-Cortex - Widespread Effects of Local Cortical-Lesions. *Science* 205: 313-316.
- Moser PC, Hitchcock JM, Lister S, Moran PM (2000) The pharmacology of latent inhibition as an animal model of schizophrenia. *Brain Res Brain Res Rev* 33: 275-307.
- Mountcastle VB, Davies PW, Berman AL (1957) Response properties of neurons of cat's somatic sensory cortex to peripheral stimuli. *J Neurophysiol* 20: 374-407.
- Mrzljak L, Bergson C, Pappy M, Huff R, Levenson R, Goldman-Rakic PS (1996) Localization of dopamine D4 receptors in GABAergic neurons of the primate brain. *Nature* 381: 245-248.
- Murphy BL, Arnsten AFT, Goldman-Rakic PS, Roth RH (1996) Increased dopamine turnover in the prefrontal cortex impairs spatial working memory performance in rats and monkeys. *Proceedings of the National Academy of Sciences of the United States of America* 93: 1325-1329.
- Neafsey EJ (1990) Prefrontal cortical control of the autonomic nervous system: anatomical and physiological observations. *Prog Brain Res* 85: 147-165.
- Newcomer JW, Farber NB, Jevtovic-Todorovic V, Selke G, Melson AK, Hershey T, Craft S, Olney JW (1999) Ketamine-induced NMDA receptor hypofunction as a model of memory impairment and psychosis. *Neuropsychopharmacology* 20: 106-118.
- Noain D, Avale ME, Wedemeyer C, Calvo D, Peper M, Rubinstein M (2006) Identification of brain neurons expressing the dopamine D4 receptor gene using BAC transgenic mice. *European Journal of Neuroscience* 24: 2429-2438.
- O'Donnell P (2003) Dopamine gating of forebrain neural ensembles. *European Journal of Neuroscience* 17: 429-435.
- Oleskevich S, Descarries L (1990) Quantified distribution of the serotonin innervation in adult rat hippocampus. *Neuroscience* 34: 19-33.
- Palacios JM, Hoyer D, Cortes R (1987) Alpha-1-Adrenoceptors in the Mammalian Brain - Similar Pharmacology But Different Distribution in Rodents and Primates. *Brain Research* 419: 65-75.
- Pandya DN, Yeterian EH (1990) Prefrontal cortex in relation to other cortical areas in rhesus monkey: architecture and connections. *Prog Brain Res* 85: 63-94.
- Parks CL, Robinson PS, Sibille E, Shenk T, Toth M (1998) Increased anxiety of mice lacking the serotonin(1A) receptor. *Proceedings of the National Academy of Sciences of the United States of America* 95: 10734-10739.
- Pazos A, Cortes R, Palacios JM (1985) Quantitative autoradiographic mapping of serotonin receptors in the rat brain. II. Serotonin-2 receptors. *Brain Res* 346: 231-249.
- Pazos A, Palacios JM (1985) Quantitative autoradiographic mapping of serotonin receptors in the rat brain. I. Serotonin-1 receptors. *Brain Res* 346: 205-230.

- Peyron C, Petit JM, Rampon C, Jouvet M, Luppi PH (1998) Forebrain afferents to the rat dorsal raphe nucleus demonstrated by retrograde and anterograde tracing methods. *Neuroscience* 82: 443-468.
- Pieribone VA, Nicholas AP, Dagerlind A, Hokfelt T (1994) Distribution of alpha 1 adrenoceptors in rat brain revealed by in situ hybridization experiments utilizing subtype-specific probes. *J Neurosci* 14: 4252-4268.
- Pompeiano M, Palacios JM, Mengod G (1992) Distribution and Cellular-Localization of Messenger-Rna Coding for 5-HT_{1A} Receptor in the Rat-Brain - Correlation with Receptor-Binding. *Journal of Neuroscience* 12: 440-453.
- Pompeiano M, Palacios JM, Mengod G (1994) Distribution of the serotonin 5-HT₂ receptor family mRNAs: comparison between 5-HT_{2A} and 5-HT_{2C} receptors. *Brain Res Mol Brain Res* 23: 163-178.
- Porrino LJ, Goldman-Rakic PS (1982) Brainstem innervation of prefrontal and anterior cingulate cortex in the rhesus monkey revealed by retrograde transport of HRP. *J Comp Neurol* 205: 63-76.
- Porter RH, Benwell KR, Lamb H, Malcolm CS, Allen NH, Revell DF, Adams DR, Sheardown MJ (1999) Functional characterization of agonists at recombinant human 5-HT_{2A}, 5-HT_{2B} and 5-HT_{2C} receptors in CHO-K1 cells. *Br J Pharmacol* 128: 13-20.
- Preston GC, Milson DS, Ceuppens PR, Warburton DM (1992) Effects of the 5-HT₃ receptor antagonist GR 68755 on a scopolamine induced cognitive deficit in healthy subjects. *Br J Clin Pharmacol* 33: 546.
- Preuss TM (1995) Do Rats Have Prefrontal Cortex - the Rose-Woolsey-Akert Program Reconsidered. *Journal of Cognitive Neuroscience* 7: 1-24.
- Puig MV, Celada P, Diaz-Mataix L, Artigas F (2003) In vivo modulation of the activity of pyramidal neurons in the rat medial prefrontal cortex by 5-HT_{2A} receptors: relationship to thalamocortical afferents. *Cereb Cortex* 13: 870-882.
- Puig MV, Santana N, Celada P, Mengod G, Artigas F (2004) In vivo excitation of GABA interneurons in the medial prefrontal cortex through 5-HT₃ receptors. *Cereb Cortex* 14: 1365-1375.
- Rajkowska G, Goldman-Rakic PS (1995) Cytoarchitectonic definition of prefrontal areas in the normal human cortex: I. Remapping of areas 9 and 46 using quantitative criteria. *Cereb Cortex* 5: 307-322.
- Rall W (1970) Dendritic neuron theory and dendrodendritic synapsis in a simple cortical system. In Schmitt FO (ed), *Neurosciences: Second Study Program*. New York: Rockefeller Press.
- Raymond JR, Mukhin YV, Gettys TW, Garnovskaya MN (1999) The recombinant 5-HT_{1A} receptor: G protein coupling and signalling pathways. *British Journal of Pharmacology* 127: 1751-1764.
- Riad N, Garcia S, Watkins KC, Jodoin N, Doucet E, Langlois X, El Mestikawy S, Hamon M, Descarries L (2000) Somatodendritic localization of 5-HT_{1A} and preterminal axonal localization of 5-HT_{1B} serotonin receptors in adult rat brain. *Journal of Comparative Neurology* 417: 181-194.
- Roerig B, Katz LC (1997) Modulation of intrinsic circuits by serotonin 5-HT₃ receptors in developing ferret visual cortex. *J Neurosci* 17: 8324-8338.

- Ropert N, Guy N (1991) Serotonin facilitates GABAergic transmission in the CA1 region of rat hippocampus in vitro. *J Physiol* 441: 121-136.
- Rose JE, Woolsey CN (1947) The Orbitofrontal Cortex and Its Connections with the Mediodorsal Nucleus in Rabbit, Sheep and Cat. *Research Publications-Association for Research in Nervous and Mental Disease* 27: 210-232.
- Sahakian BJ, Sarna GS, Kantamaneni BD, Jackson A, Hutson PH, Curzon G (1985) Association Between Learning and Cortical Catecholamines in Non-Drug-Treated Rats. *Psychopharmacology* 86: 339-343.
- Santana N, Bortolozzi A, Serrats J, Mengod G, Artigas F (2004) Expression of serotonin1A and serotonin2A receptors in pyramidal and GABAergic neurons of the rat prefrontal cortex. *Cereb Cortex* 14: 1100-1109.
- Saper CB, Sawchenko PE (2003) Magic peptides, magic antibodies: guidelines for appropriate controls for immunohistochemistry. *J Comp Neurol* 465: 161-163.
- Sara SJ, Herve-Minvielle A (1995) Inhibitory influence of frontal cortex on locus coeruleus neurons. *Proc Natl Acad Sci U S A* 92: 6032-6036.
- Sawa A, Snyder SH (2002) Schizophrenia: Diverse approaches to a complex disease. *Science* 296: 692-695.
- Sawaguchi T, Goldman-Rakic PS (1994) The role of D1-dopamine receptor in working memory: local injections of dopamine antagonists into the prefrontal cortex of rhesus monkeys performing an oculomotor delayed-response task. *J Neurophysiol* 71: 515-528.
- Sawaguchi T, Matsumura M, Kubota K (1988) Dopamine Enhances the Neuronal-Activity of Spatial Short-Term-Memory Task in the Primate Prefrontal Cortex. *Neuroscience Research* 5: 465-473.
- Schmidt CJ, Sorensen SM, Kehne JH, Carr AA, Palfreyman MG (1995) The role of 5-HT_{2A} receptors in antipsychotic activity. *Life Sci* 56: 2209-2222.
- Schmitz D, Gloveli T, Empson RM, Heinemann U (1998) Serotonin reduces polysynaptic inhibition via 5-HT_{1A} receptors in the superficial entorhinal cortex. *J Neurophysiol* 80: 1116-1121.
- Schreiber R, Brocco M, Audinot V, Gobert A, Veiga S, Millan MJ (1995) (1-(2,5-Dimethoxy-4 iodophenyl)-2-Aminopropane)-Induced Head-Twitches in the Rat Are Mediated by 5-Hydroxytryptamine (5-Ht)(_{2A}) Receptors - Modulation by Novel 5-Ht_{2A/2C} Antagonists, D-1 Antagonists and 5-Ht_{1A} Agonists. *Journal of Pharmacology and Experimental Therapeutics* 273: 101-112.
- Schultz W (1998) Predictive reward signal of dopamine neurons. *J Neurophysiol* 80: 1-27.
- Seamans JK, Yang CR (2004a) The principal features and mechanisms of dopamine modulation in the prefrontal cortex. *Progress in Neurobiology* 74: 1-57.
- Seamans JK, Yang CR (2004b) The principal features and mechanisms of dopamine modulation in the prefrontal cortex. *Prog Neurobiol* 74: 1-58.

- Seeger TF, Seymour PA, Schmidt AW, Zorn SH, Schulz DW, Lebel LA, McLean S, Guanowsky V, Howard HR, Lowe JA, III, . (1995) Ziprasidone (CP-88,059): a new antipsychotic with combined dopamine and serotonin receptor antagonist activity. *J Pharmacol Exp Ther* 275: 101-113.
- Seeman P (1992) Dopamine Receptor Sequences - Therapeutic Levels of Neuroleptics Occupy D2-Receptors, Clozapine Occupies D4. *Neuropsychopharmacology* 7: 261-284.
- Segal DS, Geyer MA, Schuckit MA (1981) Stimulant-induced psychosis: an evaluation of animal methods. *Essays Neurochem Neuropharmacol* 5: 95-129.
- Sesack SR, Carr DB, Omelchenko N, Pinto A (2003) Anatomical substrates for glutamate-dopamine interactions: evidence for specificity of connections and extrasynaptic actions. *Ann N Y Acad Sci* 1003: 36-52.
- Sesack SR, Deutch AY, Roth RH, Bunney BS (1989) Topographical organization of the efferent projections of the medial prefrontal cortex in the rat: an anterograde tract-tracing study with Phaseolus vulgaris leucoagglutinin. *J Comp Neurol* 290: 213-242.
- Sesack SR, Snyder CL, Lewis DA (1995) Axon terminals immunolabeled for dopamine or tyrosine hydroxylase synapse on GABA-immunoreactive dendrites in rat and monkey cortex. *J Comp Neurol* 363: 264-280.
- Sheldon PW, Aghajanian GK (1990) Serotonin (5-HT) induces IPSPs in pyramidal layer cells of rat piriform cortex: evidence for the involvement of a 5-HT2-activated interneuron. *Brain Res* 506: 62-69.
- Simon H, Scatton B, Moal ML (1980) Dopaminergic A10 neurones are involved in cognitive functions. *Nature* 286: 150-151.
- Somogyi P (1979) An interneurone making synapses specifically on the axon initial segment of pyramidal cells in the cerebral cortex of the cat [proceedings]. *J Physiol* 296: 18P-19P.
- Somogyi P, Cowey A (1981) Combined Golgi and electron microscopic study on the synapses formed by double bouquet cells in the visual cortex of the cat and monkey. *J Comp Neurol* 195: 547-566.
- Somogyi P, Kisvarday ZF, Freund TF, Cowey A (1984) Characterization by Golgi impregnation of neurons that accumulate 3H-GABA in the visual cortex of monkey. *Exp Brain Res* 53: 295-303.
- Somogyi P, Takagi H, Richards JG, Mohler H (1989) Subcellular localization of benzodiazepine/GABAA receptors in the cerebellum of rat, cat, and monkey using monoclonal antibodies. *J Neurosci* 9: 2197-2209.
- Somogyi P, Tamas G, Lujan R, Buhl EH (1998) Salient features of synaptic organisation in the cerebral cortex. *Brain Res Brain Res Rev* 26: 113-135.
- Sorg BA, Kalivas PW (1993) Effects of cocaine and footshock stress on extracellular dopamine levels in the medial prefrontal cortex. *Neuroscience* 53: 695-703.
- Sotelo C, Cholley B, El Mestikawy S, Gozlan H, Hamon M (1990) Direct Immunohistochemical Evidence of the Existence of 5-HT1A Autoreceptors on Serotonergic Neurons in the Midbrain Raphe Nuclei. *Eur J Neurosci* 2: 1144-1154.

- Spanagel R, Weiss F (1999) The dopamine hypothesis of reward: past and current status. *Trends Neurosci* 22: 521-527.
- Stanwood GD, Artymyshyn RP, Kung MP, Kung HF, Lucki I, McGonigle P (2000) Quantitative autoradiographic mapping of rat brain dopamine D3 binding with [*l*-125]7-OH-PIPAT: Evidence for the presence of D3 receptors on dopaminergic and nondopaminergic cell bodies and terminals. *Journal of Pharmacology and Experimental Therapeutics* 295: 1223-1231.
- Staubli U, Xu FB (1995) Effects of 5-HT₃ receptor antagonism on hippocampal theta rhythm, memory, and LTP induction in the freely moving rat. *J Neurosci* 15: 2445-2452.
- Stefansson H, Sigurdsson E, Steinthorsdottir V, Bjornsdottir S, Sigmundsson T, Ghosh S, Brynjolfsson J, Gunnarsdottir S, Ivarsson O, Chou TT, Hjaltason O, Birgisdottir B, Jonsson H, Gudnadottir VG, Gudmundsdottir E, Bjornsson A, Ingvarsson B, Ingason A, Sigfusson S, Hardardottir H, Harvey RP, Lai D, Zhou M, Brunner D, Mutel V, Gonzalo A, Lemke G, Sainz J, Johannesson G, Andresson T, Gudbjartsson D, Manolescu A, Frigge ML, Gurney ME, Kong A, Gulcher JR, Petursson H, Stefansson K (2002) Neuregulin 1 and susceptibility to schizophrenia. *Am J Hum Genet* 71: 877-892.
- Straub RE, Jiang Y, MacLean CJ, Ma Y, Webb BT, Myakishev MV, Harris-Kerr C, Wormley B, Sadek H, Kadambi B, Cesare AJ, Gibberman A, Wang X, O'Neill FA, Walsh D, Kendler KS (2002) Genetic variation in the 6p22.3 gene DTNBP1, the human ortholog of the mouse dysbindin gene, is associated with schizophrenia. *Am J Hum Genet* 71: 337-348.
- Swanson LW (1998) *Brain maps: structure of the rat brain*. Amsterdam: Elsevier.
- Tecott LH, Maricq AV, Julius D (1993) Nervous-System Distribution of the Serotonin 5-Ht₃ Receptor Messenger-Rna. *Proceedings of the National Academy of Sciences of the United States of America* 90: 1430-1434.
- Terreberry RR, Neafsey EJ (1983) Rat medial frontal cortex: a visceral motor region with a direct projection to the solitary nucleus. *Brain Res* 278: 245-249.
- Thaker GK, Carpenter WT, Jr. (2001) Advances in schizophrenia. *Nat Med* 7: 667-671.
- Thierry AM, Deniau JM, Chevalier G, Ferron A, Glowinski J (1983) An electrophysiological analysis of some afferent and efferent pathways of the rat prefrontal cortex. *Prog Brain Res* 58: 257-261.
- Tiberi M, Jarvie KR, Silvia C, Falardeau P, Gingrich JA, Godinot N, Bertrand L, Yang-Feng TL, Fremeau RT, Jr., Caron MG (1991) Cloning, molecular characterization, and chromosomal assignment of a gene encoding a second D1 dopamine receptor subtype: differential expression pattern in rat brain compared with the D1A receptor. *Proc Natl Acad Sci U S A* 88: 7491-7495.
- Tsuang M (2000) Schizophrenia: genes and environment. *Biol Psychiatry* 47: 210-220.
- Tzschentke TM (2001) Pharmacology and behavioral pharmacology of the mesocortical dopamine system. *Prog Neurobiol* 63: 241-320.
- Uberti MA, Hague C, Oller H, Minneman KP, Hall RA (2005) Heterodimerization with beta2-adrenergic receptors promotes surface expression and functional activity of alpha1D-adrenergic receptors. *J Pharmacol Exp Ther* 313: 16-23.

- Urakubo A, Jarskog LF, Lieberman JA, Gilmore JH (2001) Prenatal exposure to maternal infection alters cytokine expression in the placenta, amniotic fluid, and fetal brain. *Schizophr Res* 47: 27-36.
- Uylings HB, Groenewegen HJ, Kolb B (2003) Do rats have a prefrontal cortex? *Behav Brain Res* 146: 3-17.
- Uylings HB, van Eden CG (1990) Qualitative and quantitative comparison of the prefrontal cortex in rat and in primates, including humans. *Prog Brain Res* 85: 31-62.
- van Eden CG, Hoorneman EM, Buijs RM, Matthijssen MA, Geffard M, Uylings HB (1987) Immunocytochemical localization of dopamine in the prefrontal cortex of the rat at the light and electron microscopical level. *Neuroscience* 22: 849-862.
- Vawter MP, Crook JM, Hyde TM, Kleinman JE, Weinberger DR, Becker KG, Freed WJ (2002) Microarray analysis of gene expression in the prefrontal cortex in schizophrenia: a preliminary study. *Schizophr Res* 58: 11-20.
- Verberne AJ, Lewis SJ, Worland PJ, Beart PM, Jarrott B, Christie MJ, Louis WJ (1987) Medial prefrontal cortical lesions modulate baroreflex sensitivity in the rat. *Brain Res* 426: 243-249.
- Verney C, Alvarez C, Geffard M, Berger B (1990) Ultrastructural Double-Labeling Study of Dopamine Terminals and Gaba-Containing Neurons in Rat Anteromedial Cerebral-Cortex. *European Journal of Neuroscience* 2: 960-972.
- Vertes RP (2004) Differential projections of the infralimbic and prelimbic cortex in the rat. *Synapse* 51: 32-58.
- Vertes RP, Hoover WB, Do Valle AC, Sherman A, Rodriguez JJ (2006) Efferent projections of reuniens and rhomboid nuclei of the thalamus in the rat. *Journal of Comparative Neurology* 499: 768-796.
- Vijayraghavan S, Wang M, Birnbaum SG, Williams GV, Arnsten AF (2007) Inverted-U dopamine D1 receptor actions on prefrontal neurons engaged in working memory. *Nat Neurosci* 10: 376-384.
- Vincent SL, Khan Y, Benes FM (1993) Cellular distribution of dopamine D1 and D2 receptors in rat medial prefrontal cortex. *J Neurosci* 13: 2551-2564.
- Vincent SL, Khan Y, Benes FM (1995) Cellular Colocalization of Dopamine D-1 and D-2 Receptors in Rat Medial Prefrontal Cortex. *Synapse* 19: 112-120.
- Volk DW, Austin MC, Pierri JN, Sampson AR, Lewis DA (2000) Decreased glutamic acid decarboxylase67 messenger RNA expression in a subset of prefrontal cortical gamma-aminobutyric acid neurons in subjects with schizophrenia. *Arch Gen Psychiatry* 57: 237-245.
- Volk DW, Pierri JN, Fritschy JM, Auh S, Sampson AR, Lewis DA (2002) Reciprocal alterations in pre- and postsynaptic inhibitory markers at chandelier cell inputs to pyramidal neurons in schizophrenia. *Cereb Cortex* 12: 1063-1070.
- Volkow ND, Fowler JS, Wang GJ, Goldstein RZ (2002) Role of dopamine, the frontal cortex and memory circuits in drug addiction: insight from imaging studies. *Neurobiol Learn Mem* 78: 610-624.

- Von Huben SN, Davis SA, Lay CC, Katner SN, Crean RD, Taffe MA (2006) Differential contributions of dopaminergic D1- and D2-like receptors to cognitive function in rhesus monkeys. *Psychopharmacology (Berl)* 188: 586-596.
- Wan FJ, Geyer MA, Swerdlow NR (1995) Presynaptic dopamine-glutamate interactions in the nucleus accumbens regulate sensorimotor gating. *Psychopharmacology (Berl)* 120: 433-441.
- Wang M, Vijayraghavan S, Goldman-Rakic PS (2004) Selective D2 receptor actions on the functional circuitry of working memory. *Science* 303: 853-856.
- Wedzony K, Chocyk A, Mackowiak M, Fijał K, Czyrak A (2000) Cortical localization of dopamine D4 receptors in the rat brain - Immunocytochemical study. *Journal of Physiology and Pharmacology* 51: 205-221.
- Weinberger DR (1999) Schizophrenia: New phenes and new genes. *Biological Psychiatry* 46: 3-7.
- Weiner DM, Levey AI, Sunahara RK, Niznik HB, Odowd BF, Seeman P, Brann MR (1991) D1 and D2 Dopamine Receptor Messenger-Rna in Rat-Brain. *Proceedings of the National Academy of Sciences of the United States of America* 88: 1859-1863.
- Williams GV, Goldman-Rakic PS (1995) Modulation of memory fields by dopamine D1 receptors in prefrontal cortex. *Nature* 376: 572-575.
- Williams GV, Rao SG, Goldman-Rakic PS (2002) The physiological role of 5-HT2A receptors in working memory. *J Neurosci* 22: 2843-2854.
- Williams SM, Goldman-Rakic PS (1998) Widespread origin of the primate mesofrontal dopamine system. *Cereb Cortex* 8: 321-345.
- Willins DL, Deutch AY, Roth BL (1997) Serotonin 5-HT2A receptors are expressed on pyramidal cells and interneurons in the rat cortex. *Synapse* 27: 79-82.
- Woo TU, Whitehead RE, Melchitzky DS, Lewis DA (1998) A subclass of prefrontal gamma-aminobutyric acid axon terminals are selectively altered in schizophrenia. *Proc Natl Acad Sci U S A* 95: 5341-5346.
- Xie Z, Lee SP, O'Dowd BF, George SR (1999) Serotonin 5-HT1B and 5-HT1D receptors form homodimers when expressed alone and heterodimers when co-expressed. *FEBS Lett* 456: 63-67.
- Xu T, Pandey SC (2000) Cellular localization of serotonin(2A) (5HT(2A)) receptors in the rat brain. *Brain Res Bull* 51: 499-505.
- Zahrt J, Taylor JR, Mathew RG, Arnsten AFT (1997) Supranormal stimulation of D-1 dopamine receptors in the rodent prefrontal cortex impairs spatial working memory performance. *Journal of Neuroscience* 17: 8528-8535.
- Zelenin S, Aperia A, Heijtz RD (2002) Calcyon in the rat brain: Cloning of cDNA and expression of mRNA. *Journal of Comparative Neurology* 446: 37-45.
- Zhou FM, Hablitz JJ (1999) Activation of serotonin receptors modulates synaptic transmission in rat cerebral cortex. *J Neurophysiol* 82: 2989-2999.

